

BINDING CONSTRUCTS AND METHODS FOR USE THEREOF

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a Continuation-in-Part of U.S. Ser. No. 10/053,530, filed Jan. 17, 2002, which is incorporated herein by reference in its entirety and which
5 claims the benefit of priority of U.S. Provisional Application Ser. No. 60/367,358 (formerly U.S. Ser. No. 09/765,208, filed Jan. 17, 2001), also incorporated herein by reference in its entirety and for which a Petition to Convert a Non-Provisional Application to a Provisional Application was filed on Jan. 16, 2002, and granted on Jun. 6, 2002. This application also claims the benefit of priority of U.S. Provisional
10 Application No. 60/385,691 filed on Jun. 3, 2002, and entitled "Recombinant Signaling Receptors for Tumor Gene Therapy," which is also incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates generally to compounds having various
15 utilities including uses for research, diagnostics, and therapy, for example, immunotherapy. Compounds of the invention include immunologically active proteins and protein conjugates. Such proteins include recombinant or engineered binding proteins such as, for example, binding domain-immunoglobulin fusion proteins, which may include single chain Fv-immunoglobulin fusion proteins and
20 compounds containing single chain Fv-immunoglobulins. The present invention also relates to compositions and methods for treating conditions, diseases and disorders that would improved, eased, or lessened from the administration of, for example, polypeptide and/or nucleic acid constructs of the invention, including, for example, malignant conditions and B cell disorders, including diseases characterized by
25 autoantibody production and/or inflammation.

BACKGROUND OF THE INVENTION

The immune system is one of the most complex of the body's many intricate systems. A vast and complicated arrangement made up of many different types of cells and involving many different kinds of molecules, the human immune
30 system allows the body to respond to foreign invaders such as bacteria, viruses, and other infectious agents, as well as foreign material such as pollen. In general, the human immune system is divided into two main parts, antibody-mediated immunity

(also called "humoral" or "circulating" immunity) and cell-mediated immunity, both of which are managed by lymphocytes. Lymphocytes are one of the five kinds of white blood cells (leukocytes) circulating in the blood. There are several kinds of lymphocytes, each with different functions to perform. The most common types of lymphocytes are B lymphocytes (B cells), which are responsible for making antibodies, and T lymphocytes (T cells). Cells of the immune system not only include T cells and B cells, but also Natural Killer Cells, granulocytes (or polymorphonuclear (PMN) leukocytes), macrophages, and dendritic cells. The humoral system is managed by B cells with help from T cells and deals with infectious agents in the blood and tissues of the body. The cell-mediated system is managed by T cells and deals with cells of the body that have been infected.

An antigen is a substance, usually macromolecular, that stimulates or induces an immune response. Because of its complex macromolecular structure, a single microorganism consists of multiple antigens (*e.g.*, surface structures such as cell wall components, fimbriae, flagella, *etc.*, or extracellular proteins, such as toxins or enzymes produced by the microorganism). The coat proteins and some of the envelope proteins of animal viruses are also usually antigenic. A host is generally able to respond specifically to antigens that come into contact with components of its immune system. Both the antibody-mediated immunity and cell-mediated immunity systems involve complex interrelationships that allow them to mount immune reactions to almost any antigen. In other words, the immune system is able to recognize foreign substances (antigens) that stimulate the system to produce antibody-mediated immunity, cell-mediated immunity, or both.

The immune system complex is constituted by a variety of different cell types and organs disseminated throughout the body. These include the primary lymphoid organs, *i.e.*, the bone marrow and the thymus. All the cells of the immune system are initially derived from the bone marrow in a process called hematopoiesis. During hematopoiesis bone marrow-derived stem cells differentiate into either mature cells of the immune system ("B" cells) or into precursors of cells that migrate out of the bone marrow to mature in the thymus ("T" cells). In addition to red blood cells, platelets, and B cells, the bone marrow also produces Natural Killer cells, granulocytes, and immature thymocytes. The function of the thymus is to produce mature T cells. Immature thymocytes, also known as prothymocytes, leave the bone

marrow and migrate into the thymus where they mature and are then released into the bloodstream

The immune system complex also includes secondary lymphoid organs, *e.g.*, the spleen, the lymph nodes, *etc.*, as well as a circulatory system that is separate from blood vessels. The spleen, made up of B cells, T cells, macrophages, dendritic cells, Natural Killer cells, and red blood cells, is an immunologic filter of the blood. In addition to capturing antigens from the blood that passes through the spleen, migratory macrophages and dendritic cells bring antigens to the spleen via the bloodstream. An immune response is initiated in the spleen when macrophages or dendritic cells present the antigen to the appropriate B or T cells, and B cells become activated and produce large amounts of antibody. Lymphatic vessels and lymph nodes are the parts of a special circulatory system that carries lymph, a transparent fluid containing white blood cells, chiefly lymphocytes. Lymph bathes the tissues of the body, and is then collected in lymphatic vessels. Lymph nodes dot the network of lymphatic vessels and function as an immunologic filter for lymph when afferent lymph ducts bring lymph-containing antigens into the node. Composed mostly of T cells, B cells, dendritic cells, and macrophages, the lymph nodes drain fluid from most tissues. Antigens are filtered out of the lymph in the lymph node before the lymph is returned to the circulation. Macrophages and dendritic cells that capture antigens also present these foreign materials to T and B cells in the lymph nodes, resulting in the stimulation of B cells to develop there into antibody-secreting plasma cells. Antibodies leave the lymph node by the efferent ducts that empty into the blood stream. Lymphocytes can also leave the node by the efferent duct and travel to other sites in the lymphatic system or enter into the blood circulation. A single lymphocyte completes a circuit through the circulating blood and lymphatic systems once every 24 hours.

Tonsils, adenoids, Peyer's patches, and the appendix are also lymphoid tissues. Peyer's patches (masses of lymphocytes) are similar to the tonsils and are found throughout the body, especially in the mucous linings of the digestive and respiratory tracts. It is the function of the phagocytic cells found in Peyer's patches and other lymphatic aggregate follicles to defend the body against, for example, inadequately digested food particles crossing the gut wall and entering the blood, and to attack unwanted foreign invaders while they are still in the bowel.

The major function of B cells is the production of antibodies in response to foreign proteins of bacteria, viruses, and tumor cells. T cells are usually divided into two major groups, namely, the cytotoxic T lymphocytes ("Tc" cells or CTLs) and the helper T cells ("Th" cells or T helper cells). Th cells, also referred to as CD4+ T cells, function to augment or potentiate immune responses by the secretion of specialized factors that activate other white blood cells to fight off infection. They enhance the production of antibodies by B cells. Tc cells, also called CD8+ T cells, can directly kill certain tumor cells, viral-infected cells, and sometimes parasites. Tc cells are also important in down-regulation of immune responses. Both types of T cells often depend on the secondary lymphoid organs (the lymph nodes and spleen) as sites where activation occurs, but they are also found in other tissues of the body, including the liver, lung, blood, and intestinal and reproductive tracts.

Natural Killer cells, often referred to as NK cells, represent another type of lymphocyte and are similar to the Tc cell subset. They function as effector cells that directly kill certain tumors such as melanomas and lymphomas, and viral-infected cells. They are called "natural" killers because, unlike cytotoxic T cells, they do not need to recognize a specific antigen before carrying out their function. While NK cells, unlike the Tc cells, kill their targets without prior activation in the lymphoid organs, NK cells activated by Th cell secretions will kill tumor or viral-infected targets more effectively. NK cells target tumor cells and protect against a wide variety of infectious microbes. In several immunodeficiency diseases, including AIDS, Natural Killer cell function is abnormal. Natural Killer cells may also contribute to immunoregulation by secreting high levels of influential lymphokines. Some NK cells have surface receptors (FcγRIII, also called CD16) for the Fc portion of the IgG antibody. They bind to target cells through receptors for the Fc portion of an antibody that has reacted with antigen on a target cell. This type of cell-mediated immunity is called antibody-dependent cell-mediated cytotoxicity (ADCC). NK cells may also have receptors for the C3 component of complement, another immune defense system, and therefore recognize cells that are coated with C3 as targets. ADCC is thought to be an important defense against a variety of parasitic infections caused, for example, by protozoa and helminths.

Although small lymphocytes look identical, they can be distinguished by molecules carried on their cell surface. Not only do such markers distinguish

between B cells and T cells, they distinguish among various subsets of cells that behave differently. Every mature T cell, for instance, carries a marker known as T3 (or CD3). In addition, most helper T cells carry a T4 (CD4) marker, a molecule that recognizes Class II major histocompatibility complex ("MHC") antigens. A molecule
5 known as T8 (CD8), which recognizes Class I MHC antigens, is found on many suppressor/cytotoxic T cells.

Another group of white blood cells collectively referred to as granulocytes, or polymorphonuclear leukocytes (PMNs), is composed of three cell types. These cells, neutrophils, eosinophils, and basophils are important in the
10 removal of bacteria and parasites from the body. Neutrophils migrate through capillary walls and into infected tissue where they kill invaders (*e.g.*, bacteria) and then engulf the remnants by phagocytosis. Eosinophils are cytotoxic, releasing the contents of their granules on an invader. Basophils leave the blood and accumulate at the site of an infection or other inflammation and discharge the contents of their
15 granules, releasing a variety of mediators such as histamine, serotonin, prostaglandins and leukotrienes that, for example, increase blood flow to the area. Mediators released by basophils also play an important part in some allergic responses such as hay fever and anaphylactic responses to insect stings. . .

Monocytes are large phagocytic white blood cells released from the
20 bone marrow into the blood circulation. When a monocyte enters tissue, it develops into a macrophage. Macrophages are also large, phagocytic cells that engulf foreign material (antigens) that enter the body, as well as dead and dying cells of the body. Macrophages are important in the regulation of immune responses, and are often referred to as scavengers, or antigen-presenting cells (APCs) because they pick up and
25 ingest foreign materials and present these antigens to other cells of the immune system such as T cells and B cells. This is one of the important first steps in the initiation of an immune response. Stimulated macrophages exhibit increased levels of phagocytosis and also secrete Interleukin-1 (IL-1), a product that helps to activate B cells and T cells.

30 Dendritic cells also originate in the bone marrow and function as APCs. They are usually found in the structural compartment of lymphoid organs such as the thymus, lymph nodes and spleen, but are also found in the bloodstream and

other tissues. It is believed that dendritic cells capture antigen or bring it to the lymphoid organs where an immune response is initiated.

Important features of the immunological system relevant to host defense and/or immunity to pathogenic microorganisms include specificity, memory, and tolerance. It is understood, for example, that an antibody or reactive T cell will react specifically with the antigen that induced its formation; it will not react with other antigens. Generally, this specificity is of the same order as that of enzyme-substrate specificity or receptor-ligand specificity, although cross-reactivity is possible. The specificity of the immune response is explained by clonal selection. During the primary immune response, a specific antigen selects a pre-existing clone of specific lymphocytes and stimulates its activation, proliferation and differentiation. It is also understood that once the immune system has responded to produce a specific type of antibody or reactive T cell, it is capable of producing more of the antibody or activated T cell more rapidly and in larger amounts; this is called the secondary (or memory) response. It is also recognized that an animal generally does not undergo an immunological response to its own (potentially-antigenic) components. The animal is said to be tolerant, or unable to react to its own potentially antigenic components. This ensures that under normal conditions, an immune response to "self" antigens (called an autoimmune response) does not occur. Tolerance is brought about in a number of ways, but in essence the immune system is able to distinguish "self" components from "non-self" (foreign) antigens; it will respond to "non-self" but not to "self". Sometimes in an animal, tolerance can be "broken", which may result in an autoimmune disease.

The biological activities of the antibody-mediated and cell-mediated immune responses are different and vary from one type of infection to another. There are several classes or types of antibodies (and subclasses of various types) involved in antibody-mediated immunity. All of the classes of antibodies that are produced in response to a specific antigen react stereochemically with that antigen and not with other (different) antigens. The host has the genetic capacity to produce specific antibodies to thousands of different antigens, but does not do so until there is an appropriate (specific) antigenic stimulus. Due to clonal selection, the host produces only the homologous antibodies that will react with that antigen which, as noted above, are found in blood (plasma), lymph, and many extravascular tissues. Once the

antibody-mediated immunity response occurs following interaction of B lymphocytes with antigen and their differentiation into antibody-secreting plasma cells, the secreted antibody binds to the antigen which, in turn, results in its neutralization or elimination from the body.

5 Cell-mediated immunity, on the other hand, is mediated by specific subpopulations of T-lymphocytes called effector T cells that exist in precursor form as "resting T cells" (pT cells). These cells bear receptors for specific antigens and recognize these antigens on the surfaces of other cells. Stimulation with that antigen results in T cell activation. T cells enlarge, enter into a mitotic cycle, reproduce and
10 develop into effector T cells whose activities are responsible for this type of immunity. They also develop into clones of identical reactive T cells called memory T cells. As noted above, most of the T cells in the body belong to one of two subsets and are distinguished by the presence on their surface of one or the other of two glycoproteins designated CD4 and CD8. Which of these molecules is present
15 determines the types of cells to which the T cell can bind. T cells bearing CD8 ($CD8^+$ T cells) always recognize antigen in association with Class I MHC proteins and typically function as cytotoxic T cells. Almost all the cells of the body express Class I MHC molecules. T cells bearing CD4 ($CD4^+$ T cells) always recognize antigens in association with Class II MHC proteins on the surfaces of other cells. Only
20 specialized antigen-presenting cells express Class II MHC molecules, including dendritic cells, phagocytic cells such as macrophages, and B cells. $CD4^+$ T lymphocytes generally function as T helper cells.

 T helper cells, which include Th1 cells and Th2 cells, respond to antigen with the production of lymphokines. Th1 and Th2 cells can be distinguished
25 based on their lymphokine profiles. Like all T cells, Th cells arise in the thymus. When they are presented with an antigen by antigen-presenting dendritic cells they begin to proliferate and become activated. There are two kinds of dendritic cell, DC1 cells (descended from monocytes) and DC2 cells (which appear to be derived from lymphocytes).

30 Th1 cells (inflammatory Th1 cells involved in the elimination of pathogens residing intracellularly in vesicular compartments) are produced when DC1-type dendritic cells present antigen to the T cell receptor for antigen (TCR) and secrete Interleukin 12 (IL-12). This paracrine stimulation activates Th1 cells to secrete

their own lymphokines, in particular, Tumor-Necrosis Factor-beta (TNF- β) (also known as lymphotoxin) and Interferon-gamma (IFN- γ). These lymphokines stimulate macrophages to kill bacteria they have engulfed by phagocytosis and they recruit other leukocytes to the site producing inflammation. Th1 cells are essential for cell-mediated immunity and for controlling intracellular pathogens such as, for example, *Listeria* and *Mycobacterium tuberculosis*.

Th2 cells ("true" helper Th2 cells, which are required for antibody production by B cells) are produced when DC2-type dendritic cells present antigen to the T cell receptor for antigen and, presumably, one or more paracrine stimulants. The major lymphokines secreted by Th2 cells are Interleukin 4 (IL-4), which stimulates class-switching in B cells and promotes their synthesis of IgE antibodies, acts as a positive-feedback device promoting more pre-Th cells to enter the Th2 pathway, and blocks expression of the IL-12 receptor thereby inhibiting pre-Th cells in the thymus from entering the Th1 pathway. IL-4 also causes B cells to proliferate and differentiate into antibody-secreting plasma cells and memory B cells. IL-4 activates only B cells in the vicinity which themselves have bound the antigen, and not others, so as to sustain the specificity of the immune response. Th2 cells also produce Interleukin 5 (IL-5, which attracts and activates eosinophils), Interleukin 10 (IL-10, which inhibits IL-12 production by DCs and prevents maturation of pre-Th cells to Th1 cells), and Interleukin 13 (IL-13, which also promotes the synthesis of IgE antibodies).

Activation of the Th2 cell also causes it to begin to produce Interleukin 2 (IL-2), and to express a membrane receptor for IL-2. The secreted IL-2 autostimulates proliferation of Th2 cells. For example, IL-2 binds IL-2 receptors on other T cells (which have bound the antigen) and stimulates their proliferation. In addition IL-2, stimulated Th2 cells also produce IFN- γ and Interleukin 6 (IL-6), which mediate various aspects of the immune response. IFN- γ activates Natural Killer cells to their full cytolytic potential, and is an activator of macrophages and thus increases their antitumor activities. If the macrophages are infected by intracellular parasites, it activates macrophages which in turn destroy the parasites. IFN- γ also reinforces the antitumor activities of the cytotoxic lymphocytes, increases the nonspecific activities of NK-cells, and is one of the factors that controls the differentiation of B cells and increases the secretion of immunoglobins. IL-6

stimulates several types of leukocytes, as well as the production of Acute Phase Proteins in the liver. It is particularly important in inducing B cells to differentiate into antibody forming (plasma) cells. Thus, Th2 cells provide help for B cells and are essential for antibody-mediated immunity.

5 Cytotoxic T lymphocytes are able to kill cells that show a new or foreign antigen on their surface (for example, virus-infected cells, or tumor cells, or transplanted tissue cells). The CD8⁺ CTLs also come in two subsets: Tc1 that, like Th1 cells, secrete IFN- γ , and Tc2 that, like Th2 cells, secrete IL-4.

10 The cell-mediated immunity response also plays a role in destruction of tumor cells and in rejection of tissue transplants in animals. A major problem in tissue transplantation is rejection, which is often based on cell-mediated immunity response to "foreign" cells (because they are not a perfect antigenic match). Because tumor cells contain specific antigens not seen on normal cells they also may be recognized as foreign and destroyed by the forces of cell-mediated immunity. If tumor
15 cells develop on a regular basis in animals, it may be cell-mediated immunity that eliminates them or holds them in check. The increase in the incidence of many types of cancer (tumors) in humans with advancement of age may be correlated with a decline in the peak efficiency of the immune system that occurs about 25 years of age.

20 A summary of the types of cells involved in the expression of cell-mediated immunity follows. Tc lymphocytes kill cells bearing foreign antigen on surface in association with Class I MHC and can kill cells that are harboring intracellular parasites (either bacteria or viruses) as long as the infected cell is displaying a microbial antigen on its surface. Tc cells kill tumor cells and account for rejection of transplanted cells. Tc cells recognize antigen-Class I MHC complexes on
25 target cells, contact them, and release the contents of granules directly into the target cell membrane which lyses the cell. Th lymphocytes produce lymphokines that are "helper" factors for development of B cells into antibody-secreting plasma cells. They also produce certain lymphokines that stimulate the differentiation of effector T lymphocytes and the activity of macrophages. Th1 cells recognize antigen on
30 macrophages in association with Class II MHC and become activated (by IL-1) to produce lymphokines including IFN- γ that activates macrophages and NK cells. These cells mediate various aspects of the cell-mediated immunity response including delayed-type hypersensitivity reactions. Th2 cells recognize antigen in association

with Class II MHC on an APC and then produce interleukins and other substances that stimulate specific B cell and T cell proliferation and activity. Macrophages are important as antigen-presenting cells that initiate T cell interactions, development, and proliferation. Macrophages are also involved in expression of cell-mediated immunity because they become activated by IFN- γ produced in a cell-mediated immunity response. Activated macrophages have increased phagocytic potential and release soluble substances that cause inflammation and destroy many bacteria and other cells. Natural Killer cells are cytotoxic cells that lyse cells bearing new antigen regardless of their MHC type and even lyse some cells that bear no MHC proteins. NK cells are defined by their ability to kill cells displaying a foreign antigen (*e.g.*, tumor cells) regardless of MHC type and regardless of previous sensitization (exposure) to the antigen. NK cells can be activated by IL-2 and IFN- γ , and lyse cells in the same manner as cytotoxic T lymphocytes. Some NK cells have receptors for the Fc domain of the IgG antibody and are thus able to bind to the Fc portion of IgG on the surface of a target cell and release cytolytic components that kill the target cell via antibody-dependent cell-mediated cytotoxicity.

Extracellular factors that affect cell proliferation and differentiation have been defined as cytokines. These include the lymphokines, which are proteins produced by T-lymphocytes that have effects on the differentiation, proliferation, and activity of various cells involved in the expression of cell-mediated immunity. In general, lymphokines function by (1) focusing circulating leukocytes and lymphocytes into the site of immunological encounter; (2) stimulating the development and proliferation of B cells and T cells; (3) stimulating and preparing macrophages for their phagocytic tasks; (4) stimulating Natural Killer cells; and (5) providing antiviral cover and activity. A summary of various important lymphokines follows. Initially referred to as lymphocyte activation factor, IL-1 is mainly a product of macrophages, and has a variety of effects on various types of cells. It acts as a growth regulator of T cells and B cells, and it induces other cells such as hepatocytes to produce proteins relevant to host defense. IL-1 forms a chemotactic gradient for neutrophils and serves as an endogenous pyrogen which produces fever. Thus, IL-1 plays an important role in both the immune responses and in the inflammatory response. IL-2 stimulates the proliferation of T cells and activates NK cells. IL-3 regulates the proliferation of stem cells and the differentiation of mast cells. IL-4

causes B cell proliferation and enhanced antibody synthesis. IL-6 (also referred to as Interferon-beta2, hybridoma growth factor, B-cell differentiation factor, and hepatocyte stimulatory factor) has effects on B cell differentiation and on antibody production and on T cell activation, growth, and differentiation, and probably has a major role in the mediation of the inflammatory and immune responses initiated by infection or injury. IL-8 is a chemotactic attractant for neutrophils. IL-13 shares many of the properties of IL-4, and is a potent regulator of inflammatory and immune responses. IFN- γ is produced by T cells and may be considered a lymphokine. It is sometimes called "immune interferon" (Interferon-alpha being referred to as "leukocyte interferon" and Interferon-beta being referred to as "fibroblast interferon"). IFN- γ has several antiviral effects including inhibition of viral protein synthesis in infected cells. It also activates macrophages and NK cells, and stimulates IL-1, IL-2, and antibody production. Lymphotoxins include the Tumor Necrosis Factors. TNF-beta is produced by T cells, while TNF-alpha is produced by T cells as well as other types of cells. TNFs function to kill cells, including tumor cells (at a distance). There are several Colony Stimulating Factors (CSFs), including granulocyte macrophage colony stimulating factor (GMCSF), which cause phagocytic white cells of all types to differentiate and divide.

The nature of the membrane receptors for antigen on B cells and T cells is fairly well understood. Each B cell has approximately 10^5 membrane-bound antibody molecules (IgD or IgM) that correspond in specificity to the antibody the cell is programmed to produce (these receptors being referred to as BCRs). CD32 (Fc γ RII) on B cells are receptors for the Fc region of IgG. CD21 and CD35 on B cells are receptors for complement components. Each T cell has about 10^5 molecules of a specific antigen-binding T cell receptor (a TCR) exposed on its surface. The TCR is similar, but not identical, to an antibody. There are two types of T cells that differ in their TCRs, alpha/beta ($\alpha\beta$) T cells and gamma/delta ($\gamma\delta$) T cells. The TCR of alpha/beta T cells binds a bimolecular complex displayed by a Class I MHC molecule at the surface of an antigen-presenting cell. As noted above, most Th cells express CD4, whereas most Tc cells express CD8.

Both BCRs and TCRs are similar in that they are integral membrane proteins, they are present in thousands of identical copies exposed at the cell surface, they are made before the cell ever encounters an antigen, they are encoded by genes

assembled by the recombination of segments of DNA, they have a unique binding site that binds through non-covalent forces to a portion of the antigen called an epitope (or antigenic determinant) that depends on complementarity of the surface of the receptor and the surface of the epitope, and successful binding of the antigen receptor to the epitope, if accompanied by additional signals, results in stimulation of the cell to leave G_0 and enter the cell cycle and repeated mitosis that leads to the development of a clone of cells bearing the same antigen receptor, *i.e.*, a clone of cells of the identical specificity. BCRs and TCRs differ in their structure, the genes that encode them, and the type of epitope to which they bind.

Induction of a primary immune response begins when an antigen penetrates epithelial surfaces. It will eventually come into contact with macrophages or certain other classes of antigen presenting cells, including B cells, monocytes, dendritic cells, Langerhans cells, and endothelial cells. Antigens, such as bacterial cells, are internalized by endocytosis and "processed" by APCs, then "presented" to immunocompetent lymphocytes to initiate the early steps of the immunological response. Processing by a macrophage (for example) results in attaching antigenic materials to the surface of the membrane in association with Class II MHC molecules on the surface of the cell. The antigen-class II MHC complex is presented to a T-helper (Th2) cell which is able to recognize processed antigen associated with a Class II MHC molecule on the membrane of the macrophage. This interaction, together with stimulation by IL-1 from secreted by the macrophage, will activate the Th2 cell.

As indicated above, B cells themselves behave as APCs. Cross-linked antigens bound to antibody receptors on the surface of a B cell cause internalization of some of the antigen and expression on the B cell membrane together with Class II MHC molecules. The Th2 cell recognizes the antigen together with the Class II MHC molecules, and secretes the various lymphokines that activate the B cells to become antibody-secreting plasma cells and memory B cells. Even if the antigen cannot cross-link the receptor, it may be endocytosed by the B cell, processed, and returned to the surface in association with Class II MHC where it can be recognized by specific Th2 cells which will become activated to initiate B cell differentiation and proliferation. In any case, the overall B cell response leads to antibody-mediated immunity.

The antigen receptors on B cell surfaces are thought to be the specific types of antibodies that they are genetically-programmed to produce. Hence, there are

thousands of sub-populations of B cells distinguished by their ability to produce a unique antibody molecule. B cells can also react with a homologous antigen on the surface of the macrophage, or with soluble antigens. When a B cell is bound to antigen, and simultaneously is stimulated by IL-4 produced by a nearby Th2 cell, the B cell is stimulated to grow and divide to form a clone of identical B cells, each capable of producing identical antibody molecules. The activated B cells further differentiate into plasma cells which synthesize and secrete large amounts of antibody, and into memory B cells. The antibodies produced and secreted by the plasma cells will react specifically with the homologous antigen that induced their formation. Many of these reactions lead to host defense and to prevention of reinfection by pathogens. Memory cells play a role in secondary immune responses. Plasma cells are relatively short-lived (about one week) but produce large amounts of antibody during this period. Memory cells, on the other hand, are relatively long-lived and upon subsequent exposure to antigen they become quickly transformed into antibody-producing plasma cells.

Generation of cell mediated immunity begins when, for example, a cytotoxic T cell recognizes a processed antigen associated with Class I MHC on the membrane of a cell (usually an altered self cell, but possibly a transplanted tissue cell, or a eukaryotic parasite). Under stimulation by IL-2 produced by Th2 cells, the Tc cell becomes activated to become a cytotoxic T lymphocyte capable of lysing the cell which is showing the new foreign antigen on its surface, a primary manifestation of cell-mediated immunity. The interaction between an antigen-presenting macrophage and a Th cell stimulates the macrophage to produce and secrete a Interleukin-1 that acts locally on the Th cell, stimulating the Th-cell to differentiate and produce its own cytokines (which may here be called lymphokines because they arise from a lymphocyte). These lymphokines have various functions. IL-4 has an immediate effect on nearby B cells. IL-2 has an immediate effect on T cells as described above.

Leucocytes also express adhesion promoting receptors that mediate cell-cell and cell-matrix interactions. These adhesive interactions are crucial to the regulation of haemopoiesis and thymocyte maturation, the direction and control of leucocyte traffic and migration through tissues, and the development of immune and non-immune inflammatory responses. Several families of adhesion receptors have been identified. The leucocyte integrin family comprises three alpha-beta

heterodimeric membrane glycoproteins that share a common beta subunit, designated CD18. The alpha subunits of each of the three members, lymphocyte function associated antigen-1 (LFA-1), macrophage antigen-1 (Mac-1) and p150,95 are designated CD11a, b, and c respectively. These adhesion molecules play a critical part in the immune and inflammatory responses of leucocytes. The leucocyte integrin family is, in turn, part of the integrin superfamily, members of which are evolutionally, structurally and functionally related. Another Integrin subfamily found on leucocytes is the VLA group, so-called because the "very late activation antigens" VLA-1 and VLA-2 were originally found to appear late in T-cell activation. Members of this family function mainly as extracellular matrix adhesion receptors and are found both on haemopoietic and non-haemopoietic cells. They play a part in diverse cellular functions including tissue organisation, lymphocyte recirculation and T-cell immune responses. Another integrin subfamily, the cytoadhesins, are receptors on platelets and endothelial cells that bind extracellular matrix proteins. A second family of adhesion receptors is the immunoglobulin superfamily, members of which include CD2, LFA-3, and ICAM-1, which participate in T-cell adhesive interactions, and the antigen-specific receptors of T and B cells, CD4, CD8, and the MHC Class I and II molecules. Another recognized family of adhesion receptors is the selectins, characterized by a common lectin domain. Leucocyte adhesion molecule-1 (LAM-1), which is the human homologue of the murine homing receptor, MEL-14, is expressed on leucocytes, while endothelial leucocyte adhesion molecule-1 (ELAM-1) and granule membrane protein (GMP-140) are expressed on stimulated endothelial cells and activated platelets.

Activation of an immune response requires physical cell-cell contact in addition to cytokines. Thus, for example, development of B and T cell precursors require intimate contact with stromal cells. At least three critical cell-cell contact events are required for the generation of immune responses. The first is initial contact of a specific antigen with a naive T cell. Because of the requirement for MHC presentation, this is an obligate cell contact event. In normal situations the critical antigen presenting cell is the dendritic cell. In addition to the MHC/peptide-TCR interaction there are other non-antigen specific membrane bound ligand-receptor pairs which are important for the dendritic cell-T cell interaction. The principal one is the association of the CD28 molecule on the T cell with either of two ligands, B7.1

(CD80) and B7.2 (CD86), on the dendritic cell. These molecules are termed accessory molecules and it is understood that the CD28 molecule delivers an essential second signal to the T cell without which the T cell does not become activated.

5 A second essential cell-cell contact is between the activated T cell and an antigen-specific B cell. Most antigens are T cell-dependent, that is, an antibody response to the antigen absolutely requires T cell help. This help is delivered both by cytokines and by cell-cell contact. Cells bind specific antigen via surface Ig, then internalize, process, and present it on Class II MHC molecules. This enables them to be recognized by T cells specific for helper epitopes from the specific antigen. This
10 cell-cell interaction also requires CD28 binding to B7 on the B cell. In addition, a molecule called CD40 ligand or CD154, the expression of which is induced upon T cell activation, binds to CD40 on B cells. CD40 crosslinking promotes B cell proliferation, prevents apoptosis of germinal-center B cells, and promotes immunoglobulin isotype switching. The CD28-B7 and CD40-CD40L receptor ligand
15 interactions are both essential for the dialogue between B and T cells that causes their mutual activation.

A third cell-cell interaction that is essential in immune responses is the binding of activated B cells (which have migrated into a specialised structure in lymphoid organs called germinal centers) to follicular dendritic cells (FDCs). FDCs
20 are specialized stromal cells that hold intact, *i.e.*, unprocessed, antigen on their surface in the form of long-lived immune complexes. Among other molecules, FDCs express CD23, which binds to germinal center B cells via a CR2 receptor and stimulates differentiation to plasma cells. Time is required before a primary immune response is effective as a host defense. Antigens have to be recognized, taken up, digested,
25 processed, and presented by APCs. A few select Th cells must react with antigen and respond; preexisting B or T lymphocytes must encounter the antigen and proliferate and differentiate into effector cells (plasma cells or Tc cells). In the case of antibody-mediated immunity, antibody level has to build up to an effective physiological concentration to render its host resistant. It may take several days or weeks to reach a
30 level of effective immunity, even though this immunity may persist for many months, or years, or even a lifetime, due to the presence of the antibodies. In natural infections, the inoculum is small, and even though the antigenic stimulus increases during microbial replication, only small amounts of antibody are formed within the

first few days, and circulating antibody is not detectable until about a week after infection.

With regard to induction of a secondary immune response, it is understood that on re-exposure to microbial antigens (secondary exposure to antigen), there is an accelerated immunological response, *i.e.*, the secondary or memory response. Larger amounts of antibodies are formed in only 1-2 days. This is due to the activities of specific memory B cells or memory T cells which were formed during the primary immune response. These memory cells, when stimulated by homologous antigen, "remember" having previously seen the antigen, and are able to rapidly divide and differentiate into effector cells. Stimulating memory cells to rapidly produce very high (effective) levels of persistent circulating antibodies is the basis for giving "booster"-type vaccinations to humans and pets. Thus, following the first exposure to an antigen the immune response (as evidenced by following the concentration of specific antibody in the serum) develops gradually over a period of days, reaches a low plateau within 2-3 weeks, and usually begins to decline in a relatively short period of time. When the antigen is encountered a second time, a secondary (memory) response causes a rapid rise in the concentration of antibody, reaching a much higher level in the serum, which may persist for a relatively long period of time. This is not to say that a protective level of antibody may not be reached by primary exposure alone, but usually to ensure a high level of protective antibody that persists over a long period of time, it is necessary to have repeated antigenic stimulation of the immune system.

An immunoglobulin molecule (abbreviated Ig), is a multimeric protein composed of two identical light chain polypeptides and two identical heavy chain polypeptides (H_2L_2) that are joined into a macromolecular complex by interchain disulfide bonds, *i.e.*, covalent bonds between the sulfhydryl groups of neighboring cysteine residues. There are various classes of human antibody proteins, each of which is produced by a specific clone of plasma cells. Five human immunoglobulin classes are defined on the basis of their heavy chain composition, and are named IgG, IgM, IgA, IgE, and IgD. The IgG-class and IgA-class antibodies are further divided into subclasses, namely, IgG1, IgG2, IgG3, and IgG4, and IgA1 and IgA2. Intrachain disulfide bonds join different areas of the same polypeptide chain, which results in the formation of loops that, along with adjacent amino acids, constitute the

immunoglobulin domains. At the amino-terminal portion (also called the "NE₂-terminus" or the "N-terminus"), each light chain and each heavy chain has a single variable region that shows considerable variation in amino acid composition from one antibody to another. The light chain variable region, V_L, associates with the variable
5 region of a heavy chain, V_H, to form the antigen binding site of the immunoglobulin, called the Fv.

In addition to variable regions, each of the antibody chains have one or more constant regions. Light chains have a single constant region domain. Thus, light chains have one variable region and one constant region. Heavy chains have several
10 constant region domains. The heavy chains in IgG, IgA, and IgD antibodies have three constant region domains, which are designated CH1, CH2, and CH3, and the heavy chains in IgM and IgE antibodies have four constant region domains, CH1, CH2, CH3 and CH4. Thus, heavy chains have one variable region and three or four constant regions. Immunoglobulin structure and function are reviewed, for example,
15 in Harlow *et al.*, Eds., *Antibodies: A Laboratory Manual*, Chapter 14, Cold Spring Harbor Laboratory, Cold Spring Harbor (1988).

The heavy chains of immunoglobulins can also be divided into three functional regions: the Fd region (a fragment comprising V_H and CH1, *i.e.*, the two N-terminal domains of the heavy chain), the hinge region, and the Fc region (the
20 "fragment crystallizable" region, derived from constant regions and formed after pepsin digestion). The Fd region in combination with the light chain forms an Fab (the "fragment antigen-binding"). Because an antigen will react stereochemically with the antigen-binding region at the amino terminus of each Fab the IgG molecule is divalent, *i.e.*, it can bind to two antigen molecules. The Fc contains the domains that
25 interact with immunoglobulin receptors on cells and with the initial elements of the complement cascade. Thus, the Fc fragment is generally considered responsible for the effector functions of an immunoglobulin, such as complement fixation and binding to Fc receptors. Pepsin sometimes also cleaves before the third constant domain (CH3) of the heavy chain to give a large fragment F(abc) and a small
30 fragment pFcb. These terms are also used for analogous regions of the other immunoglobulins. The hinge region, found in IgG, IgA, and IgD class antibodies, acts as a flexible spacer, allowing the Fab portion to move freely in space. In contrast to

the constant regions, the hinge domains are structurally diverse, varying in both sequence and length among immunoglobulin classes and subclasses.

For example, the length and flexibility of the hinge region varies among the IgG subclasses. The hinge region of IgG1 encompasses amino acids 216-231 and because it is freely flexible, the Fab fragments can rotate about their axes of symmetry and move within a sphere centered at the first of two inter-heavy chain disulfide bridges. IgG2 has a shorter hinge than IgG1, with 12 amino acid residues and four disulfide bridges. The hinge region of IgG2 lacks a glycine residue, it is relatively short and contains a rigid poly-proline double helix, stabilised by extra inter-heavy chain disulfide bridges. These properties restrict the flexibility of the IgG2 molecule. IgG3 differs from the other subclasses by its unique extended hinge region (about four times as long as the IgG1 hinge), containing 62 amino acids (including 21 prolines and 11 cysteines), forming an inflexible poly-proline double helix. In IgG3 the Fab fragments are relatively far away from the Fc fragment, giving the molecule a greater flexibility. The elongated hinge in IgG3 is also responsible for its higher molecular weight compared to the other subclasses. The hinge region of IgG4 is shorter than that of IgG1 and its flexibility is intermediate between that of IgG1 and IgG2. The flexibility of the hinge region reportedly decreases in the order IgG3>IgG1>IgG4>IgG2. The four IgG subclasses also differ from each other with respect to their effector functions. This difference is related to differences in structure, including with respect to the interaction between the variable region, Fab fragments, and the constant Fc fragment. Nevertheless, aside from glycosylation within the CH2 region, for example, in spite of this knowledge there are no set rules or conventions regarding means or methods to change features, including sequences, of these subclasses of molecule to change, control, add, or remove different functions, for example, ADCC, CDC, and other functions.

According to crystallographic studies, the immunoglobulin hinge region can be further subdivided functionally into three regions: the upper hinge region, the core region, and the lower hinge region. Shin *et al.*, 1992 *Immunological Reviews* 130: 87. The upper hinge region includes amino acids from the carboxyl end of CH1 to the first residue in the hinge that restricts motion, generally the first cysteine residue that forms an interchain disulfide bond between the two heavy chains. The length of the upper hinge region correlates with the segmental flexibility

of the antibody. The core hinge region contains the inter-heavy chain disulfide bridges, and the lower hinge region joins the amino terminal end of the CH2 domain and includes residues in CH2. *Id.* The core hinge region of human IgG1 contains the sequence Cys-Pro-Pro-Cys (SEQ ID NO: 515) which, when dimerized by disulfide bond formation, results in a cyclic octapeptide believed to act as a pivot, thus conferring flexibility. The hinge region may also contain one or more glycosylation sites, which include a number of structurally distinct types of sites for carbohydrate attachment. For example, IgA1 contains five glycosylation sites within a 17 amino acid segment of the hinge region, conferring resistance of the hinge region polypeptide to intestinal proteases, considered an advantageous property for a secretory immunoglobulin.

Conformational changes permitted by the structure and flexibility of the immunoglobulin hinge region polypeptide sequence may also affect the effector functions of the Fc portion of the antibody. Three general categories of effector functions associated with the Fc region include (1) activation of the classical complement cascade, (2) interaction with effector cells, and (3) compartmentalization of immunoglobulins. The different human IgG subclasses vary in the relative efficacies with which they fix complement, or activate and amplify the steps of the complement cascade. *See, e.g., Kirschfink, 2001 Immunol. Rev.* **180**: 177; Chakraborti *et al.*, 2000 *Cell Signal* **12**: 607; Kohl *et al.*, 1999 *Mol. Immunol.* **36**: 893; Marsh *et al.*, 1999 *Curr. Opin. Nephrol. Hypertens.* **8**: 557; Speth *et al.*, 1999 *Wien Klin. Wochenschr.* **111**: 378.

Complement-dependent cytotoxicity (CDC) is believed to be a significant mechanism for clearance of specific target cells such as tumor cells. CDC is a stream of events that consists of a series of enzymes that become activated by each other in a cascade fashion. Complement has an important role in clearing antigen, accomplished by its four major functions: (1) local vasodilation; (2) attraction of immune cells, especially phagocytes (chemotaxis); (3) tagging of foreign organisms for phagocytosis (opsonization); and (4) destruction of invading organisms by the membrane attack complex (MAC attack). The central molecule is the C3 protein. It is an enzyme that is split into two fragments by components of either the classical pathway or the alternative pathway. The classical pathway is induced by antibodies, especially IgG and IgM, while the alternative pathway is nonspecifically

stimulated by bacterial products like lipopolysaccharide (LPS). Briefly, the products of the C3 split include a small peptide C3a which is chemotactic for phagocytic immune cells and results in local vasodilation by causing the release of C5a fragment from C5. The other part of C3, C3b coats antigens on the surface of foreign organisms and acts to opsonize the organism for destruction. C3b also reacts with other components of the complement system to form an MAC consisting of C5b, C6, C7, C8 and C9.

In general, IgG1 and IgG3 most effectively fix complement, IgG2 is less effective, and IgG4 does not activate complement. Complement activation is initiated by binding of C1q, a subunit of the first component C1 in the cascade, to an antigen-antibody complex. Even though the binding site for C1q is located in the CH2 domain of the antibody, the hinge region influences the ability of the antibody to activate the cascade. For example, recombinant immunoglobulins lacking a hinge region are unable to activate complement. Shin *et al.*, 1992. Without the flexibility conferred by the hinge region, the Fab portion of the antibody bound to the antigen may not be able to adopt the conformation required to permit C1q to bind to CH2. *See id.* Hinge length and segmental flexibility have been reported to correlate with complement activation; however, the correlation is not absolute. Human IgG3 molecules with altered hinge regions that are as rigid as IgG4, for example, can still effectively activate the cascade.

The absence of a hinge region, or a lack of a functional hinge region, can also affect the ability of certain human IgG immunoglobulins to bind Fc receptors on immune effector cells. Binding of an immunoglobulin to an Fc receptor facilitates antibody-dependent cell-mediated cytotoxicity, which as noted above is presumed to be an important mechanism for the elimination of tumor cells. The human IgG Fc receptor (FcR) family is divided into three groups, FcγRI (CD64), which is capable of binding IgG with high affinity, and FcγRII (CD32) and FcγRIII (CD16), both of which are lower affinity receptors. The molecular interaction between each of the three receptors and an immunoglobulin has not been defined precisely, but experimental evidence indicates that residues in the hinge proximal region of the CH2 domain may be important to the specificity of the interaction between the antibody and the Fc receptor. IgG1 myeloma proteins and recombinant IgG3 chimeric antibodies that lack a hinge region are reportedly unable to bind FcγRI, perhaps

because accessibility to CH2 is decreased. Shin *et al.*, 1993 *Intern. Rev. Immunol.* **10**: 177, 178-79.

Unusual and apparently evolutionarily unrelated exceptions to the H₂L₂ structure of conventional antibodies occur in some isotypes of the immunoglobulins found in camelids (camels, dromedaries and llamas; Hamers-Casterman *et al.*, 1993 *Nature* **363**: 446; Nguyen *et al.*, 1998 *J. Mol. Biol.* **275**: 413), nurse sharks (Roux *et al.*, 1998 *Proc. Nat. Acad. Sci. USA* **95**: 11804), and in the spotted ratfish (Nguyen, *et al.*, "Heavy-chain antibodies in Camelidae; a case of evolutionary innovation," 2002 *Immunogenetics* **54**(1): 39-47). These antibodies can apparently form antigen-binding regions using only heavy chain variable region, *i.e.*, these functional antibodies are homodimers of heavy chains only (referred to as "heavy-chain antibodies" or "HCAbs"). In both species, these variable regions often contain an extended third complementarity determining region (CDR3) that may help compensate for the lack of a light chain variable region, and there are frequent disulfide bonds between CDR regions that presumably help to stabilize the binding site. Muyldermans *et al.*, 1994 *Prot. Engineer.* **7**: 1129; Roux *et al.*, 1998. However, the precise function of the heavy chain-only "antibodies" is unknown, and the evolutionary pressure leading to their formation has not been identified. See, *e.g.*, Nguyen, *et al.*, 2002, *supra*. Camelids, including camels, llamas, and alpacas, also express conventional H₂L₂ antibodies, and the heavy chain-only antibodies thus do not appear to be present in these animals simply as an alternative antibody structure.

Variable regions (V_HH) of the camelid heavy chain-only immunoglobulins and conventional (H₂L₂) heavy chain variable regions contain amino acid differences, including differences at several positions that may be involved in the interface between conventional V_H and V_L domains. Nguyen *et al.*, 1998 *J. Mol. Biol.* **275**: 413; Muyldermans *et al.*, 1994 *Prot. Engineer.* **7**: 1129. Camelid V_HH reportedly recombines with IgG2 and IgG3 constant regions that contain hinge, CH2, and CH3 domains and lack a CH1 domain. Hamers-Casterman *et al.*, 1993 *Nature* **363**: 446. Interestingly, V_HH are encoded by a chromosomal locus distinct from the V_H locus (Nguyen *et al.*, 1998, *supra*), indicating that camelid B cells have evolved complex mechanisms of antigen recognition and differentiation. Thus, for example, llama IgG1 is a conventional (H₂L₂) antibody isotype in which V_H recombines with a constant region that contains hinge, CH1, CH2 and CH3 domains,

whereas the llama IgG2 and IgG3 are heavy chain-only isotypes that lack CH1 domains and that contain no light chains.

The classes of immunoglobulins have different physical and chemical characteristics and they exhibit unique biological properties. Their synthesis occurs at different stages and rates during an immune response and/or during the course of an infection. Their importance and functions in host resistance (immunity) are different.

Immunoglobulin G (IgG), a protein with a molecular weight of about 150,000 daltons (150 kD), is the predominant Ig in the serum. It makes up about 80% of the total antibody found in an animal at any given time, being 75% of the total serum antibody. It can diffuse out of the blood stream into the extravascular spaces and it is the most common Ig found there. Its concentration in tissue fluids is increased during inflammation, and it is particularly effective at the neutralization of bacterial extracellular toxins and viruses. It also has opsonizing ability and complement-fixing ability. The polypeptide composition of the Fc region of all IgG1 antibody molecules is relatively constant regardless of antibody specificity; however, as noted above, the Fab regions always differ in their exact amino acid sequences depending upon their antigenic specificity. Specific amino acid regions of the Fc portion of the molecule are recognized by receptors on phagocytes and certain other cells, and the Fc domain contains a peptide region that will bind to and activate complement, which is often required for the manifestation of antibody-mediated immunity. Because the IgG molecule is divalent, it can cross-link antigen molecules, which may lead to precipitation or agglutination of antigens; if IgG is bound to antigen on a microbial cell or surface, its Fc region may provide an extrinsic ligand that will be recognized by specific receptors on phagocytes. Microbial cells or viruses coated with IgG molecules are opsonized for phagocytosis, and opsonized pathogens are taken up and destroyed much more readily by phagocytes than their non-opsonized counterparts. IgG, as well as IgM and IgA, will neutralize the activity of toxins, including bacterial exotoxins. Furthermore, cross-linked IgG molecules on the surface of a cell can bind and activate complement from the serum and set off a cascade of reactions that can lead to destruction of the cell.

IgM is the first immunoglobulin to appear in the blood stream during the course of an infection. It is mainly confined to the bloodstream and provides protection against blood-borne pathogens. IgM makes up about 10% serum

immunoglobulins, and is arranged to resemble a pentamer of five immunoglobulin molecules (having a molecular weight of about 900 kD) tethered together at by their Fc domains. In addition to covalent linkages between the monomeric subunits, the pentamer is stabilized by a 1 Skd polypeptide called J chain. IgM, therefore, has a theoretical "valence" of ten (*i.e.*, it has ten exposed Fab domains). Probably, the most important role of IgM is its ability to function early in the immune responses against blood-borne pathogens given its efficiency in agglutinating particulate antigens. IgM binds also complement strongly and IgM antibodies bound to a microbial surface act as opsonins, rendering the microbe more susceptible to phagocytosis. In the presence of complement and IgM whole microbial cells may be killed and lysed. As noted above, IgM also appears on the surfaces of mature B cells as a transmembranous monomer where it functions as an antigen receptor, capable of activating B cells when bound to antigen.

Gene rearrangement at the immunoglobulin loci during lymphoid development generates a repertoire of B lymphocytes that express a diversity of antigen receptors. The gene rearrangement, which is catalysed by the rearrangement-activating gene ("RAG") recombinase, integrates the immunoglobulin V, D and J gene segments to yield productively rearranged immunoglobulin genes that encode the heavy and light chains of IgM antibodies. The diversity of IgM antibodies in this primary repertoire is achieved through combinatorial mechanisms (the choice of V, D and J gene segments utilized in a particular antibody), as well as junctional diversity. The joining of V, D and J gene segments is somewhat imprecise, and nucleotides may be inserted at the junction in a non-templated manner. There is therefore a very high degree of resultant diversity at the V-D-J borders. This contributes in a major way to the structural diversity of the third complementarity determining region of the antibody, a region that often plays a critical role in antigen recognition. This primary repertoire of IgM antibodies comprises a few million different structures. The size of this repertoire means that any incoming antigen is likely to encounter an antibody that recognizes it with acceptable affinity. A high- affinity binding site is unlikely to be available for most incoming antigens (the repertoire is not large enough), but the affinity of the available IgM antibodies in the primary repertoire will vary from antigen to antigen. If an epitope is re-iterated at high density on the surface of the antigen (*e.g.*, a repeated structure on the surface of a virus or bacterium), then an IgM

antibody may nevertheless be effective in mediating clearance of the organism, despite the low affinity of the individual interaction between antigenic epitope and immunoglobulin combining site. The density of the epitopes may allow multivalent interactions with IgM, leading to a high-avidity interaction, providing that a suitable spacing of antigenic epitopes can occur. Nevertheless, to ensure an effective and specific response, especially when the concentration of antigen is low (as may occur when the body is faced with a very small number of infecting viral particles), it would be preferable if high-affinity antibodies were available for neutralizing, for example, an infecting organism. The size of the primary repertoire mitigates against the likelihood of such high-affinity antibodies being present in this repertoire. The immune system therefore operates using a two-stage strategy. The primary repertoire of IgM antibodies is generated by a process of gene rearrangement and takes place prior to antigen encounter during early lymphocyte development. However, once foreign antigen has been encountered, those B cells in the primary repertoire that encode suitable (albeit low-affinity) antibodies are selectively expanded and subjected to an iterative alternation of directed hypermutation and antigen-mediated selection. This allows a significant maturation in affinity of the antigen-specific antibodies that are produced. Antigen triggering also drives isotype switch recombination. Thus, in the absence of external antigen stimulation and any maternally derived immunoglobulin, the serum will only contain a diversity of unmutated IgM molecules that have been generated by gene rearrangement. This repertoire shifts with age as a result of continuous antigen exposure, such that the majority of the serum immunoglobulin in older animals is composed of mutated IgG (and IgA) molecules whose specificities have developed as a consequence of antigen selection.

IgA exists as a H_2L_2 monomer of about 160 kD in serum and, in secretions, as a dimer of the H_2L_2 monomer of about 400 kD. As with IgM, polymerization (dimerization) is via a J-chain. IgA has two subclasses based on different heavy chains, IgA1 and IgA2. IgA1 is produced in bone marrow and makes up most of the serum IgA. Both IgA1 and IgA2 are synthesized in GALT (gut associated lymphoid tissues) to be secreted onto the mucosal surfaces. Because IgA may be synthesized locally and secreted in the seromucous secretions of the body, it is sometimes referred to as secretory antibody or sIgA. Quantitatively, IgA is synthesized in amounts greater than IgG, but it has a short half life in serum (6 days),

and it is lost in secretory products. The concentration of IgA in serum is about 15% of the total antibody. Secretion of dimeric IgA is mediated by a 100 kD glycoprotein called secretory component. It is the addition of the secretory piece to IgA molecules that accounts for their ability to exit the body to mucosal surfaces via the exocrine glands. IgM can be transported similarly and makes up a small proportion of secretory antibodies. Secretory IgA is the predominant immunoglobulin present in gastrointestinal fluids, nasal secretions, saliva, tears and other mucous secretions of the body. IgA antibodies are important in resistance to infection of the mucosal surfaces of the body, particularly the respiratory, intestinal and urogenital tracts. IgA acts as a protective coating for the mucous surfaces against microbial adherence or initial colonization. It can also neutralize toxin activity on mucosal surfaces. Fc receptors for IgA-coated microorganisms found on monocytes and neutrophils derived from the respiratory mucosa suggest that IgA may have a role in the lung, at least, in opsonization of pathogens. Secretory IgA is also transferred via the milk, i.e., the colostrum, from a nursing mother to a newborn, which provides passive immunity to many pathogens, especially those that enter by way of the GI tract.

IgE is a immunoglobulin of about 190 kD that accounts for about 0.002% of the total serum immunoglobulins. It is produced by plasma cells below the respiratory and intestinal epithelia. The majority of IgE is bound to tissue cells, especially mast cells. If an infectious agent succeeds in penetrating the IgA barrier, it comes up against the next line of defense, the MALT (mucosa-associated lymphoid tissues) system which is managed by IgE. IgE is bound very firmly to specific IgE Fc receptors on mast cells. Contact with antigen leads to release of mediators of inflammation from the mast cells, which effectively recruits various agents of the immune response including complement, chemotactic factors for phagocytes, T cells, *etc.* Although a well-known manifestation of this reaction is a type of immediate hypersensitivity reaction called atopic allergy (*e.g.*, hives, asthma, hay fever, *etc.*), the MALT responses act as a defense mechanism because they amplify the inflammatory response and may facilitate rejection of a pathogen.

IgD is a molecule of about 175 kd that resembles IgG in its monomeric form. IgD antibodies are found for the most part on the surfaces of B lymphocytes. The same cells may also carry IgM antibody. As noted above, it is thought that IgD

and IgM function as mutually-interacting antigen receptors for control of B cell activation and suppression. Hence, IgD may have an immunoregulatory function.

In addition to opsonization, activation of complement, and ADCC, antibodies have other functions in host defense including steric hindrance, toxin neutralization, agglutination, and precipitation. With regard to steric hindrance, it is understood that antibodies combine with the surfaces of microorganisms and may block or prevent their attachment to susceptible cells or mucosal surfaces. Antibody against a viral component can block attachment of the virus to susceptible host cells and thereby reduce infectivity. Secretory IgA can block attachment of pathogens to mucosal surfaces. Toxin-neutralizing antibodies (antitoxins) can also react with a soluble bacterial toxin and block the interaction of the toxin with its specific target cell or substrate. Antibodies can also combine with the surfaces of microorganisms or soluble antigens and cause them to agglutinate or precipitate. This reduces the number of separate infectious units and makes them more readily phagocytosed because the clump of particles is larger in size. Floccules or aggregates of neutralized toxin may be removed by phagocytes.

Antibodies have been proposed for use in therapy. Animals, including humans and mice have the ability to make antibodies able to recognize (by binding to) virtually any antigenic determinant and to discriminate between similar epitopes. Not only does this provide the basis for protection against disease organisms, but it makes antibodies attractive candidates to target other types of molecules found in the body, such as receptors or other proteins present on the surface of normal cells and molecules present uniquely on the surface of cancer cells. Thus the remarkable specificity of antibodies makes them promising agents for human therapy.

Initial antibody preparations available for use, such as intravenous gammaglobulins, included animal and human antisera that were used *in vivo* to destroy bacteria (tetanus, pneumococcus) and neutralize virus (hepatitis A and B, rabies, cytomegalovirus, and varicella zoster) in the blood of infected individuals. Possibly the most important early application was the use and endotoxins. However, there are problems associated with the use of antibodies in human therapy because the response of the immune system to any antigen, even the simplest, is "polyclonal," *i.e.*, the system manufactures antibodies of a great range of structures both in their binding regions as well as in their effector regions. Polyclonal antibody treatment was also

associated with unwanted side effects. In addition to the polyclonal nature of these antibody preparations, there was the risk of infection from contaminating viruses. Serum sickness, anaphylaxis, and were also considered limiting factors. Furthermore, even if one were to isolate a single antibody-secreting cell, and place it in culture, it would die out after a few generations because of the limited growth potential of all normal somatic cells.

Until the late 1970s, polyclonal antibodies obtained from the blood serum of immunized animals, provided the only source of antibodies for research or treatment of disease. Isolation of specific antibodies was essentially impossible until Kohler and Milstein discovered how to make "monoclonal antibodies" that would have a single specificity, that would all be alike due to manufacture by a single clone of plasma cells, and that could be grown indefinitely. This technique was described in a 1975 publication (*Nature* **256**: 495-97), and Köhler and Milstein received the 1984 Nobel Prize in Medicine for their work.

The first step in Kohler and Milstein's technique for production of monoclonal antibodies involves immunizing an experimental animal with the antigen of interest. In most of their experiments, Kohler and Milstein injected a mouse with sheep red blood cells. The mouse's body initiates an immune response and begins producing antibodies specific to the antigen. The mouse's spleen is then removed and B cells producing the antibody of interest are isolated. Tumor-producing cells that have been grown in culture are then fused with the B lymphocytes using polyethylene glycol in order to produce a "hybridoma." Only hybridomas resulting from the fusion will survive. The spleen lymphocyte has a limited life span, so any B cells that did not fuse with a myeloma will die in the culture. As well, those cells that lack the antibody-producing aspect of the B cell will not secrete the enzyme HGPRT, which is required for growth in the HAT medium. The hypoxanthine-aminopterinthymidine (HAT) medium, on which the cells are grown, inhibits the pathway for nucleotide synthesis. Cells which produce HGPRT can bypass this pathway and continue to grow. By placing the fused cells in a HAT medium, the true hybridomas can be isolated (McKay, Raff, Reichardt 1981). The isolated hybridoma cells are then screened for specificity to the desired antigen. Because each hybridoma descends from one B cell, it makes copies of only one antibody. The hybridoma that produces the antibody of interest is grown in culture to produce large amounts of monoclonal

antibodies, which are then isolated for further use. The technique is called somatic cell hybridization, and the resulting hybridoma (selected for both immortality and production of the specific antibody of interest) may be cultured indefinitely, *i.e.*, it is a potentially immortal cell line.

5 Monoclonal antibodies are now widely used as diagnostic and research reagents. However, their introduction into human therapy has been much slower. One principal difficulty is that mouse antibodies are "seen" by the human immune system as foreign, and the human patient mounts an immune response against them, producing HAMA ("human anti-mouse antibodies"). These not only cause the
10 therapeutic antibodies to be quickly eliminated from the host, but also form immune complexes that cause damage to the kidneys.

Two approaches have been used in an attempt to reduce the problem of HAMA. The first is the production of chimeric antibodies in which the antigen-binding part (variable regions) of a mouse monoclonal antibody is fused to the
15 effector part (constant region) of a human antibody using genetic engineering. In a second approach, rodent antibodies have been altered through a technique known as complementarity determining region (CDR) grafting or "humanization." In this process, the antigen binding sites, which are formed by three CDRs of the heavy chain and three CDRs of the light chain, are excised from cells secreting rodent mAb and
20 grafted into the DNA coding for the framework of the human antibody. Because only the antigen-binding site CDRs, rather than the entire variable domain of the rodent antibody are transplanted, the resulting humanized antibody (a second generation or "hyperchimeric" antibody) is reportedly less immunogenic than a first generation chimeric antibody. This process has been further improved to include changes
25 referred to as "reshaping" (Verhoeyen, *et al.*, "Reshaping human antibodies: grafting an anti-lysozyme activity," 1988 *Science* **239**: 1534-1536; Riechmann, *et al.*, "Reshaping human antibodies for therapy," 1988 *Nature* **332**: 323--337; Tempest, *et al.*, "Reshaping human monoclonal antibody to inhibit respiratory syncytial virus infection in vivo," *Bio/Technol* 1991 **9**: 266-271), "hyperchimerization" (Queen, *et al.*, "A humanized antibody that binds to the human interleukin 2 receptor," 1989
30 *Proc Natl Acad Sci USA* **86**: 10029-10033; Co, *et al.*, "Humanized antibodies for antiviral therapy," 1991 *Proc Natl Acad Sci USA* **88**: 2869-2873; Co, *et al.*, "Chimeric and humanized antibodies with specificity for the CD33 antigen," 1992 *J Immunol*

148: 1149-1154), and "veneering" (Mark, *et al.*, "Derivation of therapeutically active humanized and veneered anti-CD18 antibodies. In: Metcalf BW, Dalton BJ, eds. Cellular adhesion: molecular definition to therapeutic potential. New York: Plenum Press, 1994: 291-312). In the reshaping process on the basis of homology, the rodent variable region is compared with the consensus sequence of the protein sequence subgroup to which it belongs. Similarly, the selected human constant region accepting framework is compared with its family consensus sequence. Gussowal, *et al.*, "Humanization of monoclonal antibodies," 1991 *Meth Enzymol* **203**: 99-121. The sequence analyses identify residues, which may have undergone mutation during the affinity maturation procedure and may therefore be idiosyncratic. Inclusion of the more common human residues is said to lessen immunogenicity problems by replacing human acceptor idiosyncratic residues. Further, the reshaping process is said to allow comparison of human and rodent consensus sequences to identify any systematic "species" differences. RSV19 antibodies were humanized by employing this procedure. Taylor *et al.*, "Humanized monoclonal antibody to respiratory syncytial virus," 1991 *Lancet* **337**: 1411-1412; Tempest, *et al.*, "Reshaping a human monoclonal antibody to inhibit human respiratory syncytial virus infection in vivo," 1991 *Bio/Technol* **9**: 266-271. Hyperchimerization is an alternative method of identifying residues outside CDR regions that are likely to be involved in the reconstitution of binding activity. In this method, the human sequences are compared with murine variable region sequences and the one with highest homology is selected as the acceptor framework. As in the reshaping procedure, the "idiosyncratic" residues are replaced by more commonly occurring human residues. The non-CDR residues that may be interacting with the CDR sequences are identified. Finally, it is determined which one of these residues is to be included in the variable region framework. Humanized antibodies against CD33 antigen were reportedly developed by this method. Co, *et al.*, "Chimeric and humanized antibodies with specificity for the CD33 antigen," 1992 *J Immunol* **148**: 1149-154. See also Carter, *et al.*, "Humanization of an anti-p185 HER2 antibody for human cancer therapy," 1992 *Proc Natl Acad Sci USA* **89**: 4285-4289. The displayed surface of the protein is the primary determinant of its immunogenicity. A humanized murine antibody can thus be made less immunogenic by replacing exposed residues that differ from those commonly found in human antibodies. This method of humanization is referred to as "veneering." Appropriate replacement of the outer residues may have little or no

impact on the inner domains or interdomain framework. Veneering is a two-step process. In the first step, the most homologous human variable regions are selected and compared by each single residue to the corresponding mouse variable regions. In the second step, the mouse framework residues, which differ from its human
5 homologue, are replaced by the residues present in the human homologue. This replacement involves only those residues that are on the surface and at least partially exposed.

Nevertheless, it took more than a quarter century of research for monoclonal antibody technology and genetic engineering methods to result in the
10 development of immunoglobulin molecules for treatment of human diseases. Indeed, it was not until the past five years that monoclonal antibodies became as an expanding class of therapeutics. See Glennie M J and van de Winkel J G, *Drug Discov Today* 2003 Jun. 1;8(11): 503-10; Souriau C and Hudson P J, "Recombinant antibodies for cancer diagnosis and therapy," 2003 *Expert Opin Biol Ther.* 3(2): 305-18. See also
15 Pendley C, *et al.*, "Immunogenicity of therapeutic monoclonal antibodies," 2003 *Curr Opin Mol Ther.* 5(2): 172-9.

All the same, an average of less than one therapeutic antibody per year has been introduced to the market beginning in 1986, eleven years after the publication of monoclonal antibodies. Five murine monoclonal antibodies were
20 introduced into human medicine over a ten year period from 1986-1995, including "muromonab-CD3" (OrthoClone OKT3®), which binds to a molecule on the surface of T cells and was launched in 1986 to prevent acute rejection of organ transplants; "edrecolomab" (PANOREX®), launched in 1995 for treatment of colorectal cancer; "odulimomab" (ANTILFA®), launched in 1997 for use in transplant rejection; and,
25 "ibritumomab" (ZEVALIN® yixetan), launched in 2002 for use in non-Hodgkin's lymphoma. Additionally, one monoclonal Fab, "abciximab" (REOPRO®), was launched in 1995. It inhibits the clumping of platelets by binding the receptors on their surface that normally are linked by fibrinogen and may be helpful in preventing reclogging of the coronary arteries in patients who have undergone angioplasty. Three
30 chimeric monoclonal antibodies were also launched: "rituximab" (RITUXAN®), in 1997, which binds to the CD20 molecule found on most B cells and is used to treat B cell lymphomas; "basiliximab" (SIMULECT®), in 1998 for transplant rejection; and "infliximab" (REMICADE®) which binds to tumor necrosis factor-alpha (TNF-a), in

1998 for treatment of as rheumatoid arthritis and Crohn's disease. Additionally, "abciximab" (REOPRO®), a 47.6 kD Fab fragment of the chimeric human-murine monoclonal antibody 7E3 that binds to the glycoprotein (GP) IIb/IIIa receptor of human platelets, was launched in 1995 as an adjunct to percutaneous coronary intervention for the prevention of cardiac ischemic complications in patients undergoing percutaneous coronary intervention. Finally, seven "humanized" monoclonals were launched from 1997-2003: "daclizumab" (ZENAPAX®) in 1997, which binds to part of the IL-2 receptor produced at the surface of activated T cells and is used to prevent acute rejection of transplanted kidneys; "palivizumab" (SYNAGIS®) in 1998 for RSV; "trastuzumab" (HERCEPTIN®) in 1998, which binds HER-2, a growth factor receptor found on breast cancers cells; "gemtuzumab" (MYLOTARG®) in 2000, which is a conjugate of a monoclonal antibody that binds CD33, a cell-surface molecule expressed-by the cancerous cells in acute myelogenous leukemia (AML) but not found on the normal stem cells needed to repopulate the bone marrow; and "alemtuzumab" (MABCAMPATH®) in 2001, which binds to CD52, a molecule found on white blood cells and has produced temporary remission of chronic lymphocytic leukemia; "adalimumab" (HUMIRA® (D2E7)), a human anti-TNF monoclonal containing human-derived heavy chain and light chain variable regions and human IgG: kappa. constant regions was launched in 2002 for the treatment of rheumatoid arthritis; and, "omalizumab" (XOLAIR®), which binds to IgE and prevents it from binding to mast cells was approved in 2003 for the treatment of adults and adolescents over 12 years of age with moderate to severe persistent asthma who have a positive skin test or *in vitro* reactivity to a perennial aeroallergen and whose symptoms are inadequately controlled with inhaled corticosteroids.

Thus, protein engineering has been applied in an effort to diminish problems related to immunogenicity of administered recombinant immunoglobulin polypeptides and to try to alter antibody effector functions. However, problems remain. For example, the majority of cancer patients treated with rituximab relapse, generally within about 6-12 months, and fatal infusion reactions within 24 hours of rituximab infusion have been reported. These fatal reactions followed an infusion reaction complex that included hypoxia, pulmonary infiltrates, acute respiratory distress syndrome, myocardial infarction, ventricular fibrillation or cardiogenic shock. Acute renal failure requiring dialysis with instances of fatal outcome has also been

reported in the setting of tumor lysis syndrome following treatment with rituximab, as have severe mucocutaneous reactions, some with fatal outcome. Additionally, high doses of rituximab are required for intravenous injection because the molecule is large, approximately 150 kDa, and diffusion is limited into the lymphoid tissues where many tumor cells reside.

Trastuzumab administration can result in the development of ventricular dysfunction and congestive heart failure, and the incidence and severity of cardiac dysfunction has been reported to be particularly high in patients who received trastuzumab in combination with anthracyclines and cyclophosphamide. Trastuzumab administration can also result in severe hypersensitivity reactions (including anaphylaxis), infusion reactions, and pulmonary events.

Patients receiving daclizumab immunosuppressive therapy are at increased risk for developing lymphoproliferative disorders and opportunistic infections, and it is not known whether daclizumab use will have a long-term effect on the ability of the immune system to respond to antigens first encountered during daclizumab-induced immunosuppression.

Hepatotoxicity, including severe hepatic veno-occlusive disease (VOD), has also been reported in association with the use of gemtuzumab as a single agent, as part of a combination chemotherapy regimen, and in patients without a history of liver disease or hematopoietic stem-cell transplant (HSCT). Patients who receive gemtuzumab either before or after HSCT, patients with underlying hepatic disease or abnormal liver function, and patients receiving gemtuzumab in combinations with other chemotherapy may be at increased risk for developing severe VOD. Death from liver failure and from VOD has been reported in patients who received gemtuzumab, and it has been cautioned that even careful monitoring may not identify all patients at risk or prevent the complications of hepatotoxicity.

Hepatotoxicity was also reported in patients receiving alemtuzumab. Serious and, in some rare instances fatal, pancytopenia/marrow hypoplasia, autoimmune idiopathic thrombocytopenia, and autoimmune hemolytic anemia have occurred in patients receiving alemtuzumab therapy. Alemtuzumab can also result in serious infusion reactions as well as opportunistic infections.

In patients treated with adalimumab, serious infections and sepsis, including fatalities, have been reported, as has the exacerbation of clinical symptoms and/or radiographic evidence of demyelinating disease, and patients treated with adalimumab in clinical trials had a higher incidence of lymphoma than the expected rate in the general population.

Serious adverse reactions in clinical studies with omalizumab have included malignancies and anaphylaxis, in which the observed incidence of malignancy among omalizumab-treated patients (0.5%) was numerically higher than among patients in control groups (0.2%).

Smaller immunoglobulin molecules have been constructed in an effort to overcome various problems associated with whole immunoglobulin therapy. Single chain immunoglobulin variable region fragment polypeptides (scFvs) are made of an immunoglobulin heavy chain variable domain joined via a short linker peptide to an immunoglobulin light chain variable domain. Huston *et al.*, 1988 *Proc. Natl. Acad. Sci. USA*, **85**: 5879-83. It has been suggested that the smaller size of scFv molecules may lead to more rapid clearance from plasma and more effective penetration into tissues than whole immunoglobulins. See, e.g., Jain, 1990 *Cancer Res.* **50**: 814s-819s. An anti-tumor scFv was reported to show more rapid tumor penetration and more even distribution through the tumor mass than the corresponding chimeric antibody. Yokota *et al.*, *Cancer Res.* **52**: 3402-08 (1992).

Despite advantages that scFv molecules may have with regard to serotherapy, drawbacks to this therapeutic approach also exist. For example, rapid clearance of scFv may prevent delivery of a minimum effective dose to the target tissue. Additionally, manufacturing adequate amounts of scFv for administration to patients has been challenging due to difficulties in expression and isolation of scFv that adversely affect yields. During expression, scFv molecules lack stability and often aggregate due to pairing of variable regions from different molecules. Furthermore, production levels of scFv molecules in mammalian expression systems are reportedly low, which may limit the potential for efficient manufacturing of scFv molecules for therapy. Davis *et al.*, 1990 *J. Biol. Chem.* **265**: 10410-18; Traunecker *et al.*, 1991 *EMBO J.* **10**: 3655-59. Strategies for means to improve production have been explored, and reportedly include the addition of glycosylation sites to variable regions. See, e.g., U.S. Pat. No. 5,888,773; Jost *et al.*, 1994 *J. Biol. Chem.* **269**:

26267-73. Another disadvantage to the use of scFvs for therapy is the lack of effector function. An scFv that lacks the cytolytic functions, ADCC, and complement dependent-cytotoxicity may be less effective or ineffective for treating disease. Even though development of scFv technology began over 12 years ago, there are currently
5 no scFv products approved for therapy.

Alternatively, it has been proposed that fusion of an scFv to another molecule, such as a toxin, could take advantage of the specific antigen-binding activity and the small size of an scFv to deliver the toxin to a target tissue. Chaudary
10 *et al.*, 1989 *Nature* **339**: 394; Batra *et al.*, 1991 *Mol. Cell. Biol.* **11**: 2200. Conjugation or fusion of toxins to scFvs has thus been offered as an alternative strategy to provide potent, antigen-specific molecules, but dosing with such conjugates or chimeras can be limited by excessive and/or non-specific toxicity due to the toxin moiety of such preparations. Toxic effects may include supraphysiological elevation of liver enzymes and vascular leak syndrome, and other undesired effects. In addition, immunotoxins
15 are themselves highly immunogenic upon administration to a host, and host antibodies generated against the immunotoxin limit potential usefulness for repeated therapeutic treatments of an individual.

Fusion proteins in which immunoglobulin constant region polypeptide sequences are present and nonimmunoglobulin sequences are substituted for the
20 antibody variable regions have also been investigated. For example, CD4, the T cell surface protein recognized by HIV, was recombinantly fused to an immunoglobulin Fc effector domain, and an IL-2-IgG1 fusion protein reportedly effected complement-mediated lysis of IL-2 receptor-bearing cells. *See Sensel et al.*, *Chem. Immunol.* **65**: 129-158 (1997).

25 An extensive introduction as well as detailed information about all aspects of recombinant antibody technology can be found in the textbook "Recombinant Antibodies" (John Wiley & Sons, NY, 1999). A comprehensive collection of detailed antibody engineering lab Protocols can be found in R. Kontermann and S. Dubel (eds.), "The Antibody Engineering Lab Manual" (Springer
30 Verlag, Heidelberg/NY, 2000). Diseases and disorders thought to be amenable to some type of immunoglobulin therapy include cancer and immune system disorders. Cancer includes a broad range of diseases, affecting approximately one in four individuals worldwide. Rapid and unregulated proliferation of malignant cells is a

hallmark of many types of cancer, including hematological malignancies. Although patients with a hematologic malignant condition have benefited from advances in cancer therapy in the past two decades, Multani *et al.*, 1998 *J. Clin. Oncology* 16: 3691-3710, and remission times have increased, most patients still relapse and succumb to their disease. Barriers to cure with cytotoxic drugs include, for example, tumor cell resistance and the high toxicity of chemotherapy, which prevents optimal dosing in many patients.

Nevertheless, patients have been treated with immunotherapeutics that target malignant cells, *i.e.*, to antigens expressed on tumor cells. With regard to the selection of tumor cell surface antigens suitable for use as immunotherapy targets, it is preferable that such a target antigen is not expressed by normal tissues, particularly where the preservation of such tissue is important to host survival. In the case of hematologic malignancy, malignant cells express many antigens that are not expressed on the surfaces of stem cells or other essential cells. Treatment of a hematologic malignant condition using a therapeutic regimen that depletes both normal and malignant cells of hematological origin has been acceptable where regeneration of normal cells from progenitors can occur after therapy has ended. Additionally, the target antigen is desirably expressed on all or virtually all clonogenic populations of tumor cells, and it is best that expression persists despite selective pressure from immunoglobulin therapy. Strategies that employ selection of a cell surface idiotype (*e.g.*, a particular idiotope) as a target for therapy of B cell malignancy have been limited by the outgrowth of tumor cell variants with altered surface idiotype expression, even where the antigen exhibits a high degree of tumor selectivity. Meeker *et al.*, 1985 *N. Engl. J. Med.* 312: 1658-65. The selected antigen should also traffic properly after an immunoglobulin binds to it. Shedding or internalization of a cell surface target antigen after an immunoglobulin binds to the antigen may allow tumor cells to escape destruction, thus limiting the effectiveness of serotherapy. Finally, binding of an immunoglobulin to cell surface target antigens that transmit or transduce cellular activation signals may result in improved functional responses to immunotherapy in tumor cells, and can lead to growth arrest and/or apoptosis. While all of these properties are important, the triggering of apoptosis after an immunoglobulin binds to the target antigen may also be a critical factor in achieving successful serotherapy.

Antigens that have been tested as targets for serotherapy of B and T cell malignancies include Ig idiotype (Brown *et al.*, 1989 *Blood* 73: 651-61), CD19 (Hekman *et al.*, 1991 *Cancer Immunol. Immunother.* 32: 364-72), Vlasveld *et al.*, 1995 *Cancer Immunol. Immunother.* 40: 37-47), CD20 (Press *et al.*, 1987 *Blood* 69: 584-91), Maloney *et al.*, 1997 *J. Clin. Oncol.* 15: 3266-74), CD21 (Scheinberg *et al.*, 1990 *J. Clin. Oncol.* 8: 792-803), CD5 (Dillman *et al.*, 1986 *J. Biol. Respn. Mod.* 5: 394-410), and CD52 (CAMPATH) (Pawson *et al.*, 1997 *J. Clin. Oncol.* 15: 2667-72). Of these, greater benefit for therapy of B cell lymphomas has been obtained using molecules that target CD20. Other targets have been limited by biological properties of the antigen. For example, surface idiotype can be altered through somatic mutation, allowing tumor cell escape. CD5, CD21, and CD19 are rapidly internalized after monoclonal antibody binding, allowing tumor cells to escape destruction unless monoclonal antibodies are conjugated with toxin molecules. CD22 is expressed on only a subset of B cell lymphomas, thereby limiting its usefulness, while CD52 is expressed on both T cells and B cells and may therefore generate counterproductive immunosuppression by depletion.

Treatment of patients with low grade or follicular B cell lymphoma using a chimeric CD20 monoclonal antibody has been reported to induce partial or complete responses in patients. McLaughlin *et al.*, 1996 *Blood* 88: 90a (abstract, suppl. 1); Maloney *et al.*, 1997 *Blood* 90: 2188-95. However, as noted above, tumor relapse commonly occurs within six months to one year. Further improvements in serotherapy are needed to induce more durable responses, for example, in low grade B cell lymphoma, and to allow effective treatment of high grade lymphoma and other B cell diseases.

Another approach has been to target radioisotopes to B cell lymphomas using monoclonal antibodies specific for CD20. While the effectiveness of therapy is reportedly increased, associated toxicity from the long *in vivo* half-life of the radioactive antibody increases also, sometimes requiring that the patient undergo stem cell rescue. Press *et al.*, 1993 *N. Eng. J. Med.* 329: 1219-1224; Kaminski *et al.*, 1993 *N. Eng. J. Med.* 329: 459-65. Monoclonal antibodies to CD20 have also been cleaved with proteases to yield F(ab')₂ or Fab fragments prior to attachment of radioisotope. This has been reported to improve penetration of the radioisotope conjugate into the tumor and to shorten the *in vivo* half-life, thus reducing the toxicity to normal tissues.

However, these molecules lack effector functions, including complement fixation and/or ADCC.

CD20 was the first human B cell lineage-specific surface molecule identified by a monoclonal antibody. It is a non-glycosylated, hydrophobic 35 kDa B cell transmembrane phosphoprotein that has both amino and carboxy ends situated in the cytoplasm. Einfeld *et al.*, 1988 *EMBO J.* **7**: 711-17. CD20 is expressed by all normal mature B cells, but is not expressed by precursor B cells. Natural ligands for CD20 have not been identified, and the function of CD20 in B cell biology is still incompletely understood.

Anti-CD20 monoclonal antibodies affect the viability and growth and growth of B cells. Clark *et al.*, 1986 *Proc. Natl. Acad. Sci. USA* **83**: 4494-98. Extensive cross-linking of CD20 can induce apoptosis in B lymphoma cell lines, Shan *et al.*, 1998 *Blood* **91**: 1644-52, and cross-linking of CD20 on the cell surface has been reported to increase the magnitude and enhance the kinetics of signal transduction, for example, as detected by measuring tyrosine phosphorylation of cellular substrates. Deans *et al.*, 1993 *J. Immunol.* **146**: 846-53. Therefore, in addition to cellular depletion by complement and ADCC mechanisms, Fc-receptor binding by CD20 monoclonal antibodies *in vivo* may promote apoptosis of malignant B cells by CD20 cross-linking, consistent with the theory that effectiveness of CD20 therapy of human lymphoma in a SCID mouse model may be dependent upon Fc-receptor binding by the CD20 monoclonal antibody. Funakoshi *et al.*, 1996 *J. Immunotherapy* **19**: 93-101. The presence of multiple membrane spanning domains in the CD20 polypeptide (Einfeld *et al.*, 1988 *EMBO J.* **7**: 711-17; Stamenkovic *et al.*, 1988 *J. Exp. Med.* **167**: 1975-80; Tedder *et al.*, 1988 *J. Immunol.* **141**: 4388-4394), prevent CD20 internalization after antibody binding, and this was recognized as an important feature for therapy of B cell malignancies when a murine CD20 monoclonal antibody, IF5, was injected into patients with B cell lymphoma, resulting in significant depletion of malignant cells and partial clinical responses. Press *et al.*, 1987 *Blood* **69**: 584-91.

Because normal mature B cells also express CD20, normal B cells are depleted by anti-CD20 antibody therapy. Reff, M. E. *et al.*, 1994 *Blood* **83**: 435-445. After treatment is completed, however, normal B cells can be regenerated from CD20 negative B cell precursors; therefore, patients treated with anti-CD20 therapy do not experience significant immunosuppression. Depletion of normal B cells may also be

beneficial in diseases that involve inappropriate production of autoantibodies or other diseases where B cells may play a role. A chimeric monoclonal antibody specific for CD20, consisting of heavy and light chain variable regions of mouse origin fused to human IgG1 heavy chain and human kappa light chain constant regions, reportedly retained binding to CD20 and the ability to mediate ADCC and to fix complement. Liu *et al.*, 1987 *J. Immunol.* **139**: 3521-26. The mechanism of anti-tumor activity of rituximab, discussed above, is thought to be a combination of several activities, including ADCC, complement fixation, and triggering of signals that promote apoptosis in malignant B cells, although the large size of rituximab prevents optimal diffusion of the molecule into lymphoid tissues that contain malignant B cells, thereby limiting these anti-tumor activities. Autoimmune diseases include autoimmune thyroid diseases, which include Graves' disease and Hashimoto's thyroiditis. In the United States alone, there are about 20 million people who have some form of autoimmune thyroid disease. Autoimmune thyroid disease results from the production of autoantibodies that either stimulate the thyroid to cause hyperthyroidism (Graves' disease) or destroy the thyroid to cause hypothyroidism (Hashimoto's thyroiditis). Stimulation of the thyroid is caused by autoantibodies that bind and activate the thyroid stimulating hormone (TSH) receptor. Destruction of the thyroid is caused by autoantibodies that react with other thyroid antigens. Current therapy for Graves' disease includes surgery, radioactive iodine, or antithyroid drug therapy. Radioactive iodine is widely used, since antithyroid medications have significant side effects and disease recurrence is high. Surgery is reserved for patients with large goiters or where there is a need for very rapid normalization of thyroid function. There are no therapies that target the production of autoantibodies responsible for stimulating the TSH receptor. Current therapy for Hashimoto's thyroiditis is levothyroxine sodium, and therapy is usually lifelong because of the low likelihood of remission. Suppressive therapy has been shown to shrink goiters in Hashimoto's thyroiditis, but no therapies that reduce autoantibody production to target the disease mechanism are known.

Rheumatoid arthritis (RA) is a chronic disease characterized by inflammation of the joints, leading to swelling, pain, and loss of function. RA affects an estimated 2.5 million people in the United States. RA is caused by a combination of events including an initial infection or injury, an abnormal immune response, and genetic factors. While autoreactive T cells and B cells are present in RA, the detection

of high levels of antibodies that collect in the joints, called rheumatoid factor, is used in the diagnosis of RA. Current therapy for RA includes many medications for managing pain and slowing the progression of the disease. No therapy has been found that can cure the disease. Medications include nonsteroidal antiinflammatory drugs (NSAIDS), and disease modifying antirheumatic drugs (DMARDS). NSAIDS are useful in benign disease, but fail to prevent the progression to joint destruction and debility in severe RA. Both NSAIDS and DMARDS are associated with significant side effects. Only one new DMARD, Leflunomide, has been approved in over 10 years. Leflunomide blocks production of autoantibodies, reduces inflammation, and slows progression of RA. However, this drug also causes severe side effects including nausea, diarrhea, hair loss, rash, and liver injury.

Systemic Lupus Erythematosus (SLE) is an autoimmune disease caused by recurrent injuries to blood vessels in multiple organs, including the kidney, skin, and joints. SLE is estimated to affect over 500,000 people in the United States. In patients with SLE, a faulty interaction between T cells and B cells results in the production of autoantibodies that attack the cell nucleus. These include anti-double stranded DNA and anti-Sm antibodies. Autoantibodies that bind phospholipids are also found in about half of SLE patients, and are responsible for blood vessel damage and low blood counts. Immune complexes accumulate the kidneys, blood vessels, and joints of SLE patients, where they cause inflammation and tissue damage. No treatment for SLE has been found to cure the disease. NSAIDS and DMARDS are used for therapy depending upon the severity of the disease. Plasmapheresis with plasma exchange to remove autoantibodies can cause temporary improvement in SLE patients. There is general agreement that autoantibodies are responsible for SLE, so new therapies that deplete the B cell lineage, allowing the immune system to reset as new B cells are generated from precursors, would offer hope for long lasting benefit in SLE patients.

Sjogren's syndrome is an autoimmune disease characterized by destruction of the body's moisture-producing glands. Sjogren's syndrome is one of the most prevalent autoimmune disorders, striking up to an estimated 4 million people in the United States. About half of people stricken with Sjogren's syndrome also have a connective tissue disease, such as RA, while the other half have primary Sjogren's syndrome with no other concurrent autoimmune disease. Autoantibodies, including

anti-nuclear antibodies, rheumatoid factor, anti-fodrin, and anti-muscarinic receptor are often present in patients with Sjogren's syndrome. Conventional therapy includes corticosteroids, and additional more effective therapies would be of benefit.

Immune thrombocytopenic purpura (ITP) is caused by autoantibodies that bind to blood platelets and cause their destruction. Some cases of ITP are caused by drugs, and others are associated with infection, pregnancy, or autoimmune disease such as SLE. About half of all cases are classified as "idiopathic"; meaning the cause is unknown. The treatment of ITP is determined by the severity of the symptoms. In some cases, no therapy is needed although in most cases immunosuppressive drugs, including corticosteroids or intravenous infusions of immune globulin to deplete T cells, are provided. Another treatment that usually results in an increased number of platelets is removal of the spleen, the organ that destroys antibody-coated platelets. More potent immunosuppressive drugs, including cyclosporine, cyclophosphamide, or azathioprine are used for patients with severe cases. Removal of autoantibodies by passage of patients' plasma over a Protein A column is used as a second line treatment in patients with severe disease. Additional more effective therapies are desired.

Multiple sclerosis (MS) is also an autoimmune disease. It is characterized by inflammation of the central nervous system and destruction of myelin, which insulates nerve cell fibers in the brain, spinal cord, and body. Although the cause of MS is unknown, it is widely believed that autoimmune T cells are primary contributors to the pathogenesis of the disease. However, high levels of antibodies are present in the cerebral spinal fluid of patients with MS, and some theories predict that the B cell response leading to antibody production is important for mediating the disease. No B cell depletion therapies have been studies in patients with MS, and there is no cure for MS. Current therapy is corticosteroids, which can reduce the duration and severity of attacks, but do not affect the course of MS over time. New biotechnology interferon (IFN) therapies for MS have recently been approved but additional more effectiver therapies are desired.

Myasthenia Gravis (MG) is a chronic autoimmune neuromuscular disorder that is characterized by weakness of the voluntary muscle groups. MG effects about 40,000 people in the united states. MG is caused by autoantibodies that bind to acetylcholine receptors expressed at neuromuscular junctions. The autoantibodies reduce or block acetylcholine receptors, preventing the transmission of signals from

nerves to muscles. There is no known cure for mg. Common treatments include immunosuppression with corticosteroids, cyclosporine, cyclophosphamide, or azathioprine. Surgical removal of the thymus is often used to blunt the autoimmune response. Plasmapheresis, used to reduce autoantibody levels in the blood, is effective
5 in mg, but is short-lived because the production of autoantibodies continues. Plasmapheresis is usually reserved for severe muscle weakness prior to surgery. New and effective therapies would be of benefit.

Psoriasis effects approximately five million people, and is characterized by autoimmune inflammation in the skin. Psoriasis is also associated
10 with arthritis in 30% (psoriatic arthritis). Many treatments, including steroids, uv light retenoids, vitamin d derivatives, cyclosporine, methotrexate have been used but it is also plain that psoriasis would benefit from new and effective therapies. Scieroderma is a chronic autoimmune disease of the connective tissue that is also known as systemic sclerosis. Scleroderma is characterized by an overproduction of collagen,
15 resulting in a thickening of the skin, and approxiamtely 300,000 people in the united states have scleroderma, which would also benefit from new and effective therapies.

There is a clear need for improved compositions and methods to treat malignacies, including B cell malignancies and disorders including autoimmune diseases, disorders, and conditions, as well as other diseases, disorders, and
20 conditions. The compositions and methods of the present invention described and claimed herein provide such improved compositions and methods as well as other advantages.

SUMMARY OF THE INVENTION

It is an aspect of the present invention to provide a binding domain-
25 immunoglobulin fusion protein, comprising a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge or hinge-acting region polypeptide, which in turn is fused or otherwise connected to a region comprising one or more native or engineered constant regions from an immunoglobulin heavy chain, other than CH1, for example, the CH2 and CH3 regions of IgG and IgA, or the CH3
30 and CH4 regions of IgE. The binding domain-immunoglobulin fusion protein further comprises a region that comprises, consists essentially of, or consists of, a native or engineered immunoglobulin heavy chain CH2 constant region polypeptide (or CH3 in

the case of a construct derived in whole or in part from IgE) that is fused or otherwise connected to the hinge region polypeptide and a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide (or CH4 in the case of a construct derived in whole or in part from IgE) that is fused or otherwise connected to the CH2 constant region polypeptide (or CH3 in the case of a construct derived in whole or in part from IgE). Such binding domain-immunoglobulin fusion proteins are capable of at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity and complement fixation. Such binding domain polypeptides are also capable of binding or specifically binding to a target, for example, a target antigen.

In certain embodiments, for example, the binding domain polypeptide comprises at least one immunoglobulin variable region polypeptide that is selected from a native or engineered immunoglobulin light chain variable region polypeptide and/or a native or engineered immunoglobulin heavy chain variable region polypeptide. In certain further embodiments the binding domain-immunoglobulin fusion protein comprises a native or engineered immunoglobulin heavy chain variable region polypeptide, wherein the heavy chain variable region polypeptide is an engineered human immunoglobulin heavy chain variable region polypeptide (or an engineered immunoglobulin heavy chain variable region polypeptide from a non-human species) comprising a mutation, substitution, or deletion of an amino acid(s) at a location corresponding to any one or more of amino acid positions 9, 10, 11, 12, 108, 110, and/or 112. Mutations, substitutions, or deletions of an amino acid(s) at a location corresponding to any one or more of amino acid positions 9, 10, 11, 12, 108, 110, and/or 112 in a heavy chain variable region may be included within a construct such as the construct corresponding to, for example, SEQ ID NO: 212. In certain embodiments the immunoglobulin variable region polypeptide is derived from, for example, a human immunoglobulin, and in certain other embodiments the immunoglobulin variable region polypeptide comprises a humanized immunoglobulin polypeptide sequence. In certain embodiments the immunoglobulin variable region polypeptide, whether or not humanized, is derived from a murine immunoglobulin, or is derived from an immunoglobulin from another species, including, for example a rat, a pig, a monkey, or a camelid.

According to certain embodiments of the present invention, the binding domain polypeptide comprises, consists essentially of, or consists of, (a) at least one native or engineered immunoglobulin light chain variable region polypeptide; (b) at least one native or engineered immunoglobulin heavy chain variable region polypeptide; and (c) at least one linker polypeptide that is fused or otherwise connected to the polypeptide of (a) and to the polypeptide of (b). In certain further embodiments the native or engineered immunoglobulin light chain variable region and heavy chain variable region polypeptides are derived or constructed from human immunoglobulins, and in certain other further embodiments the linker polypeptide comprises at least one polypeptide including or having as an amino acid sequence Gly-Gly-Gly-Gly-Ser [SEQ ID NO: 516]. In other embodiments the linker polypeptide comprises at least two or three repeats of a polypeptide having as an amino acid sequence Gly-Gly-Gly-Gly-Ser [SEQ ID NO: 516]. In other embodiments the linker comprises a glycosylation site, which in certain further embodiments is an asparagine-linked glycosylation site, an O-linked glycosylation site, a C-mannosylation site, a glypiation site or a phosphoglycation site. In another embodiment at least one of a native or engineered immunoglobulin heavy chain CH2 constant region polypeptide and a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide is derived from an IgG or IgA human immunoglobulin heavy chain. In another embodiment at least one of a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide and a native or engineered immunoglobulin heavy chain CH4 constant region polypeptide is derived from an IgE human immunoglobulin heavy chain.

An immunoglobulin hinge region polypeptide may comprise, consist essentially or, or consist of, for example, any of (1) any hinge or hinge-acting peptide or polypeptide that occurs naturally for example, a human immunoglobulin hinge region polypeptide including, for example, a wild-type human IgG hinge or a portion thereof, a wild-type human IgA hinge or a portion thereof, a wild-type human IgD hinge or a portion thereof, or a wild-type human IgE hinge-acting region, *i.e.*, IgE CH2, or a portion thereof, a wild-type camelid hinge region or a portion thereof (including a IgG1 llama hinge region or portion thereof, a IgG2 llama hinge region or portion thereof, and a IgG3 llama hinge region or portion thereof), a nurse shark hinge region or portion thereof, and/or a spotted ratfish hinge region or a portion thereof; (2)

a mutated or otherwise altered or engineered hinge region polypeptide that contains no cysteine residues and that is derived or constructed from a wild-type immunoglobulin hinge region polypeptide having one or more cysteine residues; (3) a mutated or otherwise altered or engineered hinge region polypeptide that contains one cysteine residue and that is derived from a wild-type immunoglobulin hinge region polypeptide having one or more cysteine residues; (4) a hinge region polypeptide that has been mutated or otherwise altered or engineered to contain or add one or more glycosylation sites, for example, an asparagine-linked glycosylation site, an O-linked glycosylation site, a C-mannosylation site, a glypiation site or a phosphoglycation site; (5) a mutated or otherwise altered or engineered hinge region polypeptide in which the number of cysteine residues is reduced by amino acid substitution or deletion, for example, a mutated or otherwise altered or engineered IgG1 or IgG4 hinge region containing for example zero, one, or two cysteine residues, a mutated or otherwise altered or engineered IgG2 hinge region containing for example zero, one, two or three cysteine residues, a mutated or otherwise altered or engineered IgG3 hinge region containing for example zero, one, two, three, or from four to ten cysteine residues, or a mutated or otherwise altered or engineered human IgA1 or IgA2 hinge region polypeptide that contains zero or only one or two cysteine residues (*e.g.*, an "SCC" hinge), a mutated or otherwise altered or engineered IgD hinge region containing no cysteine residues, or a mutated or otherwise altered or engineered human IgE hinge-acting region, *i.e.*, IgE CH2 region polypeptide that contains zero or only one, two, three or four cysteine residues; or (6) any other connecting region molecule described or referenced herein or otherwise known or later discovered as useful for connecting adjoining immunoglobulin domains such as, for example, a CH1 domain and a CH2 domain. For example, a hinge region polypeptide may be selected from the group consisting of (i) a wild-type human IgG1 immunoglobulin hinge region polypeptide, for example, (ii) a mutated or otherwise altered or engineered human IgG 1 or other immunoglobulin hinge region polypeptide that is derived or constructed from a wild-type immunoglobulin hinge region polypeptide having three or more cysteine residues, wherein said mutated human IgG1 or other immunoglobulin hinge region polypeptide contains two cysteine residues and wherein a first cysteine of the wild-type hinge region is not mutated, (iii) a mutated or otherwise altered or engineered human IgG1 or other immunoglobulin hinge region polypeptide that is derived from a wild-type immunoglobulin hinge region

polypeptide having three or more cysteine residues, wherein said mutated human IgG1 or other immunoglobulin hinge region polypeptide contains no more than one cysteine residue, and (iv) a mutated or otherwise altered or engineered human IgG1 or other immunoglobulin hinge region polypeptide that is derived from a wild-type immunoglobulin hinge region polypeptide having three or more cysteine residues, wherein said mutated or otherwise altered or engineered human IgG1 or other immunoglobulin hinge region polypeptide contains no cysteine residues. In certain embodiments, for example, the immunoglobulin hinge region polypeptide is a mutated or otherwise altered or engineered hinge region polypeptide and exhibits a reduced ability to dimerize, relative to a wild-type human immunoglobulin G or other wild type hinge region or hinge-acting polypeptide.

The immunoglobulin heavy chain constant region polypeptides may be, for example, native or engineered CH2 and CH3 domains of an isotype that is human IgG or human IgA. The immunoglobulin heavy chain constant region polypeptides may also be, for example, native or engineered immunoglobulin heavy chain constant region CH3 and CH4 polypeptides of an isotype that is human IgE.

In certain other embodiments the target or target antigen may be, for example, CD19 (B-lymphocyte antigen CD19, also referred to as B-lymphocyte surface antigen B4, or Leu-12), CD20 (B-lymphocyte antigen 20, also referred to as B-lymphocyte surface antigen B1, Leu-16, or Bp35), CD22 (B-cell receptor CD22, also referred to as Leu-14, B-lymphocyte cell adhesion molecule, or BL-CAM), CD37 (leukocyte antigen CD37), CD40 (B-cell surface antigen CD40, also referred to as Tumor Necrosis Factor receptor superfamily member 5, CD40L receptor, or Bp50), CD80 (T lymphocyte activation antigen CD80, also referred to as Activation B7-1 antigen, B7, B7-1, or BB1), CD86 (T lymphocyte activation antigen CD86, also referred to as Activation B7-2 antigen, B70, FUN-1, or BU63), CD137 (also referred to as Tumor Necrosis Factor receptor superfamily member 9), CD152 (also referred to as cytotoxic T-lymphocyte protein 4 or CTLA-4), CD45 (Leukocyte common antigen, also referred to as L-CA, T200, and EC 3.1.3.48), CD45RA (an isoform of CD45, and an antigen expressed on naive or immature lymphocytes), CD45RB (an isoform of CD45); CD45RO (an isoform of CD45, and a common leukocyte antigen expressed on memory B and T cells), L6 (Tumor-associated antigen L6, also referred to as Transmembrane 4 superfamily member 1, Membrane component surface marker 1, or

M3S1), CD2 (T-cell surface antigen CD2, also referred to as T-cell surface antigen T11/Leu-5, LFA-2, LFA-3 receptor, Erythrocyte receptor, or Rosette receptor), CD28 (T-cell-specific homodimer surface protein CD28, also referred to as Tp44), CD30 (lymphocyte activation antigen CD30, also referred to as Tumor Necrosis Factor receptor superfamily member 8, CD30L receptor, or Ki-1), CD50 (also referred to as Intercellular adhesion molecule-3 (ICAM3), or ICAM-R), CD54 (also referred to as Intercellular adhesion molecule-1 (ICAM1), or Major group rhinovirus receptor), B7-H1 (ligand for an immunoinhibitory receptor expressed by activated T cells, B cells, and myeloid cells, also referred to as PD-L1; see Dong, *et al.*, "B7-H1, a third member of the B7 family, co-stimulates T-cell proliferation and interleukin-10 secretion," 1999 *Nat. Med.* **5**: 1365-1369), CD134 (also referred to as Tumor Necrosis Factor receptor superfamily member 4, OX40, OX40L receptor, ACT35 antigen, or TAX-transcriptionally activated glycoprotein 1 receptor), 41BB (4-1BB ligand receptor, T-cell antigen 4-1BB, or T-cell antigen ILA), CD153 (also referred to as Tumor Necrosis Factor ligand superfamily member 8, CD30 ligand, or CD30-L), CD154 (also referred to as Tumor Necrosis Factor ligand superfamily member 5, CD40 ligand, CD40-L, TNF-related activation protein, TRAP, or T cell antigen Gp39), ICOS (Inducible Costimulator), CD3 (one or more of the delta, epsilon, gamma, eta and/or zeta chains, alone or in combination), CD4 (T-cell surface glycoprotein CD4, also referred to as T-cell surface antigen T4/Leu-3), CD25 (also referred to as Interleukin-2 receptor alpha chain, IL-2 receptor alpha subunit, p55, or Tac antigen), CD8 α (T-cell surface glycoprotein CD8 alpha chain, also referred to as T-lymphocyte differentiation antigen, T8/Leu-2, and Lyl-2), CD11b (also referred to as Integrin alpha-M, Cell surface glycoprotein MAC-1 alpha subunit, CR-3 alpha chain, Leukocyte adhesion receptor Mol, or Neutrophil adherence receptor), CD14 (Monocyte differentiation antigen CD14, also referred to as Myeloid cell-specific leucine-rich glycoprotein or LPS receptor), CD56 (also referred to as Neural cell adhesion molecule 1), or CD69 (also referred to as Early T-cell activation antigen p60, Gp32/28, Leu-23, MLR-3, Activation inducer molecule, or AIM). The above list of construct targets and/or target antigens is exemplary only and is not exhaustive.

In another aspect, the invention includes a binding construct (or a polynucleotide encoding such a construct) that comprises a CD154 extracellular domain, or desired functional portion thereof. In one embodiment of this aspect of the

invention, for example, the binding construct comprises a CD 154 extracellular domain fused or otherwise connected to a second binding domain. The second binding domain, for example, may comprise, consist essentially of, or consist of at least one immunoglobulin variable region polypeptide. The at least one

5 immunoglobulin variable region polypeptide may be a native or engineered scFv. The native or engineered scFv may be a native or engineered scFv disclosed or described herein. The second binding domain, including a native or engineered scFv, may be one that binds, for example, to any of the targets, including target antigens, disclosed or described herein, including but not limited to, for example, any of B7-H1, ICOS,

10 L6, CD2, CD3, CD8, CD4, CD11b, CD14, CD19, CD20, CD22, CD25, CD28, CD30, CD37, CD40, CD45, CD50, CD54, CD56, CD69, CD80, CD86, CD134, CD137, CD152, CD153, or CD154.

In another embodiment the binding domain polypeptide comprises a CTLA-4 extracellular domain, or desired functional portion thereof, and in further

15 embodiments at least one of the immunoglobulin heavy chain constant region polypeptides selected from a CH2 constant region polypeptide and a CH3 constant region polypeptide is a human IgG1 constant region polypeptide, either native or engineered.

In another further embodiment at least one of the immunoglobulin

20 heavy chain constant region polypeptides selected from a CH2 constant region polypeptide and a CH3 constant region polypeptide is a human IgA constant region polypeptide, either native or engineered.

In another further embodiment at least one of the immunoglobulin heavy chain constant region polypeptides selected from a CH3 constant region

25 polypeptide and a CH4 constant region polypeptide is a human IgE constant region polypeptide, either native or engineered.

Turning to another embodiment, the present invention provides a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of, (a) a binding domain polypeptide that is fused or otherwise

30 connected to an immunoglobulin hinge region polypeptide; (b) a native or engineered immunoglobulin heavy chain CH2 (or IgE Ch3) constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide; and (c) a native or

engineered immunoglobulin heavy chain CH3 (or IgE CH4) constant region polypeptide that is fused or otherwise connected to the CH2 (or IgE CH3) constant region polypeptide, wherein (1) the binding domain polypeptide comprises a CTLA-4 extracellular domain, or a portion thereof, that is capable of binding or specifically binding to at least one CTLA-4 ligand selected from the group consisting of CD80 and CD86, (2) the immunoglobulin hinge region polypeptide may be as described above or herein, and may comprise, consist essentially of, or consist of, for example, a polypeptide that is selected from the group consisting of a native or engineered human IgA hinge region polypeptide, a native or engineered human IgG1 hinge region polypeptide, and a native or engineered human IgE CH2 region polypeptide (3) a immunoglobulin heavy chain constant region polypeptide that comprises, consists essentially of, or consists of, a polypeptide that is selected from the group consisting of a native or engineered human IgA heavy chain CH2 constant region polypeptide, a native or engineered human IgG1 heavy chain CH2 constant region polypeptide, and a native or engineered human IgE heavy chain CH3 constant region polypeptide (4) a immunoglobulin heavy chain constant region polypeptide that comprises, consists essentially of, or consists of, a polypeptide that is selected from the group consisting of a native or engineered human IgA heavy chain CH3 constant region polypeptide, a native or engineered human IgG1 heavy chain CH3 constant region polypeptide, and a native or engineered human IgE heavy chain CH4 constant region polypeptide, and (5) the binding domain-immunoglobulin fusion protein is capable of inducing at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity, CDC, and complement fixation.

In another embodiment the present invention provides a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of (a) a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge region polypeptide, wherein said hinge region polypeptide may be as described above or herein, and may comprise, consist essentially of, or consist of, for example, a native or engineered human IgE hinge-acting region, *i.e.*, a IgE CH2 region polypeptide; (b) a first native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide, wherein said native or engineered constant region polypeptide comprises, consists essentially of, or consists of, a native or engineered human IgE

CH3 constant region polypeptide; and (c) a second native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the first native or engineered constant region polypeptide, wherein said native or engineered second constant region polypeptide comprises, consists essentially of, or consists of, a native or engineered human IgE CH4 constant region polypeptide and wherein (1) the binding domain-immunoglobulin fusion protein is capable of inducing at least one immunological activity selected from antibody dependent cell-mediated cytotoxicity and induction of an allergic response mechanism, and (2) the binding domain polypeptide is capable of binding or specifically binding to an antigen. In a further embodiment the antigen is a tumor antigen.

In certain other embodiments the present invention provides a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of, (a) a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge region polypeptide, wherein the binding domain polypeptide is capable of binding or specifically binding to at least one antigen that is present on an immune effector cell and wherein the hinge region polypeptide may be as described above or herein, and may comprise, consist essentially of, or consist of, for example, a polypeptide selected from the group consisting of a native or engineered human IgA hinge region polypeptide, a native or engineered human IgG hinge region polypeptide, and a native or engineered human IgE hinge-acting region, *i.e.*, IgE CH2 region polypeptide; (b) a first native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide, wherein said first native or engineered constant region polypeptide comprises, consists essentially of, or consists of, a polypeptide selected from the group consisting of a native or engineered human IgA CH2 constant region polypeptide, a native or engineered human IgG CH2 constant region polypeptide, and a native or engineered human IgE CH3 constant region polypeptide; (c) a second native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the first constant region polypeptide, wherein said second constant region polypeptide comprises, consists essentially of, or consists of, a polypeptide selected from the group consisting of a native or engineered human IgA CH3 constant region polypeptide, a native or engineered human IgG CH3 constant

region polypeptide, and a native or engineered human IgE CH4 constant region polypeptide; and (d) a native or engineered plasma membrane anchor polypeptide. In one example of this embodiment, the plasma membrane anchor domain polypeptide links to a membrane via a native or engineered glycosyl-
 5 phosphatidylinositol-linkage- . In a further embodiment the plasma membrane anchor domain polypeptide comprises, consists essentially of, or consists of, a native or engineered transmembrane domain polypeptide. In another further embodiment the membrane anchor domain polypeptide comprises, consists essentially of, or consists of, a native or engineered transmembrane domain polypeptide and a native or
 10 engineered cytoplasmic tail polypeptide. In a still further embodiment the cytoplasmic tail polypeptide comprises, consists essentially of, or consists of, a native or engineered apoptosis signaling polypeptide sequence, which in a still further embodiment is derived or constructed from a native or engineered receptor death domain polypeptide, a death domain, or a functional portion of either. In a further
 15 embodiment the native or engineered death domain polypeptide comprises, consists essentially of, or consists of, for example, a native or engineered polypeptide selected from an ITIM domain (immunoreceptor Tyr-based inhibition motif), an ITAM domain (immunoreceptor Tyr-based activation motif), TRAF, RIP, CRADD, FADD (Fas-associated death domain), TRADD (Tumor Necrosis Factor receptor type 1 associated DEATH domain protein), RAIDD (also referred to as RAID), CD95 (Tumor Necrosis Factor receptor superfamily member 6, also referred to as FASL receptor, Apoptosis-mediating surface antigen FAS, FAS and Apo-1 antigen),
 20 TNFR1, and/or DR5 (death receptor-5). In another embodiment the native or engineered apoptosis signaling polypeptide sequence comprises, consists essentially of, or consists of, for example, a polypeptide sequence derived from a native or engineered caspase polypeptide that is caspase-3 or caspase-8 or caspase-10, including caspase 8/FLICE/MACH/Mch5 and caspase 10/Flice2/Mch4. In another embodiment the plasma membrane anchor domain polypeptide comprises, consists essentially of, or consists of, for example, a native or engineered glycosyl-
 25 phosphatidylinositol-linkage polypeptide sequence. In another embodiment the antigen that is present on an immune effector cell is, for example, CD2, CD16, CD28, CD30, CD32, CD40, CD50, CD54, CD64, CD80, CD86, B7-H1, CD134, CD137, CD152, CD153, CD154, ICOS, CD19, CD20, CD22, CD37, L6, CD3, CD4, CD25, CD8, CD11b, CD14, CD56, or CD69. In another embodiment the human IgG is a

native or engineered human IgG1. These binding domain-immunoglobulin fusion proteins may be capable of inducing, for example, at least one immunological activity selected from antibody dependent cell-mediated cytotoxicity and/or complement fixation and/or CDC, and are capable of binding or specifically binding to a target, including, for example, a target antigen. Immune effector cells include, for example, granulocytes, mast cells, monocytes, macrophages, dendritic cells, neutrophils, eosinophils, basophils, NK cells, T cells (including Th1 cells, Th2 cells, Tc cells, memory T cells, null cells, and large granular lymphocytes, *etc.*), and B cells. This embodiment of the invention further includes the use of such proteins for therapy, and, for example, the use of such vectors for *in vivo* and *ex vivo* gene therapy. The above lists of construct components and targets are not exhaustive and may include any desired target or component that may function as, or be useful for the purposes, described herein.

In another embodiment, the invention provides a protein having a first protein motif that comprises, consists essentially of, or consists of, (1) a native or engineered immunoglobulin hinge region or hinge-acting region (*e.g.*, IgE CH2) polypeptide that is fused or otherwise connected to (2) a native or engineered CH2 constant region polypeptide (or native or engineered IgE CH3 constant region polypeptide). Said first protein motif may be fused or otherwise connected to one or more other such first protein motifs to form a second protein motif, the second protein motif being fused or otherwise connected to (3) a native or engineered CH3 constant region (or a native or engineered IgE CH4 constant region) to form a third protein motif. Said first, second or third protein motifs may be fused or otherwise connected to one or more of the herein-described native or engineered plasma membrane anchor domain polypeptides, including, for example, a native or engineered transmembrane domain polypeptide, and a native or engineered transmembrane domain polypeptide and a native or engineered cytoplasmic tail polypeptide, such as for example, a native or engineered apoptosis signaling polypeptide sequence, which may be derived or constructed from a native or engineered receptor death domain polypeptide, a death domain, or a functional portion of either. Thus, a protein or polynucleotide within this aspect of the invention may be, for example, a Hinge-CH2-CH3-TransmembraneDomain-DeathDomain construct. It may also be, for example, a (Hinge-CH2)_x-CH3- TransmembraneDomain-DeathDomain construct, where X is

from 2 to about 5, or such other number as may be needed to achieve a desired length or Fc receptor binding and/or complement fixation function(s). This embodiment of the invention also includes polynucleotides encoding such proteins, vectors including such polynucleotides, and host cells containing such polynucleotides and vectors. This
5 embodiment of the invention further includes the use of such proteins for therapy, and, for example, the use of such polynucleotides and/or vectors for *in vivo* and *ex vivo* gene therapy. The invention provides, in another embodiment, a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of, (a) a binding domain polypeptide that is fused or otherwise connected to an
10 immunoglobulin hinge region polypeptide, wherein the binding domain polypeptide is capable of binding or specifically binding to at least one antigen that is present on a cancer cell surface and wherein the hinge region polypeptide may be as described above or herein, and may comprise, consist essentially of, or consist of, for example, a polypeptide selected from the group consisting of a native or engineered human IgA
15 hinge region polypeptide, a native or engineered human IgG hinge region polypeptide, and a native or engineered human IgE hinge-acting region, *i.e.*, IgE CH2, region polypeptide; (b) a first native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide, wherein the first constant region polypeptide comprises, consists
20 essentially of, or consists of, a polypeptide that is a native or engineered human IgA CH2 constant region polypeptide, a native or engineered human IgG CH2 constant region polypeptide, or a native or engineered human IgE CH3 constant region polypeptide; and (c) a second native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the first constant
25 region polypeptide, wherein the second constant region polypeptide comprises, consists essentially of, or consists of, a polypeptide that is a native or engineered human IgA CH3 constant region polypeptide, a native or engineered human IgG CH3 constant region polypeptide, or a native or engineered human IgE CH4 constant region polypeptide. In a further embodiment the human IgG polypeptides are native
30 or engineered human IgG1 polypeptides.

In another embodiment the present invention provides a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of, (a) a binding domain polypeptide that is fused or otherwise connected to

an immunoglobulin hinge region polypeptide, wherein said hinge region polypeptide may be as described above or herein, and may comprises, consist essentially of, or consist of, for example, a wild-type or engineered human IgA hinge region polypeptide; (b) a native or engineered immunoglobulin heavy chain CH2 constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide, wherein said native or engineered CH2 constant region polypeptide comprises, consists essentially of, or consists of, a native or engineered human IgA CH2 constant region polypeptide; and (c) a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide that is fused or otherwise connected to the native or engineered CH2 constant region polypeptide, wherein the native or engineered CH3 constant region polypeptide comprises, consists essentially of, or consists of, a polypeptide that is (i) a wild-type human IgA CH3 constant region polypeptide or other IgA region, preferably human or humanized, that is capable of associating with J Chain, (ii) a mutated, altered or otherwise engineered human IgA CH3 constant region polypeptide that is, for example, incapable of associating with a J chain, wherein (1) the binding domain-immunoglobulin fusion protein is capable of at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity, CDC, and complement fixation, and (2) the binding domain polypeptide is capable of binding or specifically binding to a target such as, for example, an antigen. In certain further embodiments the mutated human IgA CH3 constant region polypeptide that is incapable of associating with a J chain is (i) a polypeptide comprising, consisting essentially of, or consisting of, an amino acid sequence as set forth in SEQ ID NO: 69 or (ii) a polypeptide comprising, consisting essentially of, or consisting of, an amino acid sequence as set forth in SEQ ID NO: 74. In other embodiments, the IgA hinge region polypeptide is a native or engineered IgA1 hinge region polypeptide or a native or engineered IgA2 hinge region polypeptide. In still other embodiments, the IgA hinge region polypeptide is different from a wild-type IgA1 or IgA2 hinge region polypeptide by, for example, the alteration, substitution, or deletion of one or more of the cysteine residues within said wild-type hinge region.

In certain other embodiments the present invention provides a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of (a) a binding domain polypeptide that is fused or otherwise connected to

an immunoglobulin hinge region polypeptide; (b) a native or engineered immunoglobulin heavy chain CH2 constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide, wherein the native or engineered CH2 constant region polypeptide comprises, consists essentially of, or consists of, a native or engineered llama CH2 constant region polypeptide that is a native or engineered llama IgG1 CH2 constant region polypeptide, a native or engineered llama IgG2 CH2 constant region polypeptide, or a native or engineered llama IgG3 CH2 constant region polypeptide; and (c) a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide that is fused or otherwise connected to the native or engineered CH2 constant region polypeptide, wherein said native or engineered CH3 constant region polypeptide comprises, consists essentially of, or consists of, a native or engineered llama CH3 constant region polypeptide that is selected from the group consisting of a native or engineered llama IgG1 CH3 constant region polypeptide, a native or engineered llama IgG2 CH3 constant region polypeptide and a native or engineered llama IgG3 CH3 constant region polypeptide wherein (1) the binding domain-immunoglobulin fusion protein is capable of at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity, fixation of complement and CDC, and (2) the binding domain polypeptide is capable of binding or specifically binding to a target, for example a target antigen. In a further embodiment the immunoglobulin hinge region polypeptide, the native or engineered llama CH2 constant region polypeptide and the native or engineered llama CH3 constant region polypeptide comprise sequences derived from a native or engineered llama IgG1 polypeptide and the fusion protein does not include a native or engineered llama IgG1 CH1 domain. In certain embodiments the invention provides any of the above described binding domain-immunoglobulin fusion proteins wherein the hinge region polypeptide is mutated, engineered, or otherwise altered to contain a glycosylation site, which in certain further embodiments is an asparagine-linked glycosylation site, an O-linked glycosylation site, a C-mannosylation site, a glypiation site or a phosphoglycation site.

In certain embodiments the invention, there are provided any of the above or herein described binding constructs, including binding domain-immunoglobulin fusion proteins, wherein a binding region or binding domain

polypeptide comprises two or more binding domain polypeptide sequences wherein each of the binding domain polypeptide sequences is capable of binding or specifically binding to a target(s) such as an antigen(s), which target(s) or antigen(s) may be the same or may be different. A native, for more preferably an engineered, IgD hinge is a desired connecting region between binding domains of a bispecific molecule of the invention, *i.e.*, one with two or more binding domains, preferably two. The wild type human IgD hinge has one cysteine that forms a disulfide bond with the light chain in the native IgD structure. It is desirable to mutate or delete this cysteine in the human IgD hinge for use as a connecting region between binding domains of, for example, a bispecific molecule. Other amino acid changes or deletions or alterations in an IgD hinge that do not result in undesired hinge inflexibility are within the scope of the invention native or engineered IgD hinge regions from other species are also within the scope of the invention, as are humanized native or engineered IgD hinges from non-human species. The present invention also provides, in certain embodiments, a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of (a) a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge region polypeptide, wherein the hinge region polypeptide may be as described above or herein, and may comprise, consist essentially of, or consist of, for example, an alternative hinge region polypeptide sequence; (b) a first native or engineered immunoglobulin heavy chain constant region, such as an IgG or IgA CH2 constant region polypeptide (or an IgE CH3 constant region polypeptide) that is fused or otherwise connected to the hinge region polypeptide; and (c) a second native or engineered immunoglobulin heavy chain constant region, such as an IgG or IgA CH3 constant region polypeptide (or an IgE CH4 constant region polypeptide) that is fused or otherwise connected to the first constant region polypeptide, wherein: (1) the binding domain-immunoglobulin fusion protein is capable of at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity, CDC, and complement fixation, and (2) the binding domain polypeptide is capable of binding or specifically binding to a target, such as an antigen.

Turning to another embodiment there is provided a binding domain-immunoglobulin fusion protein, comprising, consisting essentially of, or consisting of

(a) a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge region polypeptide, wherein the binding domain polypeptide is capable of binding or specifically binding to at least one target, such as an antigen, that is present on a cancer cell surface and wherein the hinge region polypeptide may be as described above or herein, and may comprise, consist essentially of, or consist of, for example, an alternative hinge region polypeptide sequence; (b) a first native or engineered immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide, wherein said native or engineered constant region polypeptide comprises, consists essentially of, or consists of, a polypeptide selected from the group consisting of a native or engineered human IgA CH2 constant region polypeptide, a native or engineered human IgG CH2 constant region polypeptide, and a native or engineered human IgE CH3 constant region polypeptide; and (c) a second immunoglobulin heavy chain constant region polypeptide that is fused or otherwise connected to the first constant region polypeptide, wherein the second constant region polypeptide comprises, consists essentially of, or consists of, a polypeptide that is a native or engineered human IgA CH3 constant region polypeptide, a native or engineered human IgG CH3 constant region polypeptide, or a native or engineered human IgE CH4 constant region polypeptide. In certain further embodiments the alternative hinge region polypeptide sequence comprises, consists essentially of, or consists of, a polypeptide sequence of at least ten continuous amino acids of an Ig hinge region in any one of the constructs set out herein.

In certain embodiments the present invention provides polynucleotides or vectors (including cloning vectors and expression vectors) or transformed or transfected cells, including isolated or purified or pure polynucleotides, vectors, and isolated transformed or transfected cells, encoding or containing any one of the above or herein described polypeptide or protein constructs of the invention, for example, including binding domain-immunoglobulin fusion proteins. Thus, in various embodiments the invention provides a recombinant cloning or expression construct comprising any such polynucleotide that is operably linked to a promoter.

In other embodiments there is provided a host cell transformed or transfected with, or otherwise containing, any such recombinant cloning or expression

construct. Host cells include the cells of a subject undergoing *ex vivo* cell therapy including, for example, *ex vivo* gene therapy.

5 In a related embodiment there is provided a method of producing a polypeptide or protein or other construct of the invention, for example, including a binding domain-immunoglobulin fusion protein, comprising the steps of (a) culturing a host cell as described or provided for herein under conditions that permit expression of the construct, for example, a binding domain-immunoglobulin fusion protein; and (b) isolating the construct, for example, the binding domain-immunoglobulin fusion protein from the host cell or host cell culture.

10 In another embodiment there is provided a pharmaceutical composition comprising any one of the above or herein described polypeptide or protein or other constructs of the invention, for example (including, for example, binding domain-immunoglobulin fusion proteins), in combination with a physiologically acceptable carrier.

15 In another embodiment the invention provides a pharmaceutical composition comprising, for example, an isolated, purified, or pure polynucleotide encoding any one of the polypeptide or protein constructs of the invention, for example (including, for example, binding domain-immunoglobulin fusion proteins), in combination with a physiologically acceptable carrier, or for example, in combination with, or in, a gene therapy delivery vehicle or vector.

In another embodiment the invention provides a method of treating a subject having or suspected of having a malignant condition or a B cell disorder, comprising administering to a patient a therapeutically effective amount of any of the pharmaceutical compositions described or claimed herein.

25 In certain further embodiments the malignant condition or B cell disorder is a B cell lymphoma or B cell leukemia, or a disease characterized by autoantibody production, and in certain other further embodiments the B cell disorder is, for example, rheumatoid arthritis, myasthenia gravis, Grave's disease, type I diabetes mellitus, multiple sclerosis or an autoimmune disease. In certain other
30 embodiments the malignant condition is, for example, melanoma, myeloma, glioma, astrocytoma, lymphoma, leukemia, carcinoma, or sarcoma, and so on.

It is another aspect of the present invention to provide a binding domain-immunoglobulin fusion protein, comprising, consisting essentially or, or consisting of, (a) a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge region polypeptide, wherein said hinge region polypeptide is as described herein, and may be selected from the group consisting of (i) a mutated, engineered or otherwise altered hinge region polypeptide that contains no cysteine residues and that is derived from a wild-type immunoglobulin hinge region polypeptide having one or more cysteine residues, (ii) a mutated, engineered or otherwise altered hinge region polypeptide that contains one cysteine residue and that is derived from a wild-type immunoglobulin hinge region polypeptide having two or more cysteine residues, (iii) a wild-type human IgA hinge region polypeptide, (iv) a mutated, engineered or otherwise altered human IgA hinge region polypeptide that contains no cysteine residues, (v) a mutated, engineered or otherwise altered human IgA hinge region polypeptide that contains one cysteine residue and (vi) a mutated, engineered or otherwise altered human IgA hinge region polypeptide that contains two cysteine residues; (b) a native or engineered immunoglobulin heavy chain CH2 constant region polypeptide that is fused or otherwise connected to the hinge region polypeptide; and (c) a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide that is fused or otherwise connected to the CH2 constant region polypeptide, wherein: (1) the binding domain-immunoglobulin fusion protein is capable of at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity and complement fixation, and (2) the binding domain polypeptide is capable of binding or specifically binding to an antigen. In one embodiment the immunoglobulin hinge region polypeptide is a mutated hinge region polypeptide, for example, and the resulting construct exhibits a reduced ability to dimerize, relative to a construct containing a wild-type human immunoglobulin G hinge region polypeptide. In another embodiment the binding domain polypeptide comprises, consists essentially of, or consists of, at least one native or engineered immunoglobulin variable region polypeptide that is a native or engineered immunoglobulin light chain variable region polypeptide and/or a native or engineered immunoglobulin heavy chain variable region polypeptide. In a further embodiment the native or engineered immunoglobulin variable region polypeptide is derived from a human immunoglobulin and, for example, may be humanized.

In another embodiment, the invention provides a binding domain-immunoglobulin fusion protein includes a binding domain polypeptide that comprises, consists essentially of, or consists of, (a) at least one native or engineered immunoglobulin light chain variable region polypeptide; (b) at least one native or engineered immunoglobulin heavy chain variable region polypeptide; and (c) at least one linker peptide that is fused or otherwise connected to the polypeptide of (a) and to the polypeptide of (b). In a further embodiment the native or engineered immunoglobulin light chain variable region and the native or engineered heavy chain variable region polypeptides are derived from human immunoglobulins and may, for example, be humanized. In another embodiment at least one of the native or engineered immunoglobulin heavy chain CH2 (or IgE CH3) constant region polypeptide and the native or engineered immunoglobulin heavy chain CH3 (or IgE CH4) constant region polypeptide is derived or constructed from a human immunoglobulin heavy chain. In another embodiment the native or engineered immunoglobulin heavy chain constant region CH2 and CH3 polypeptides are of, or are derived or otherwise prepared or constructed from, an isotype selected from human IgG and human IgA. In another embodiment the target, for example, the target antigen is selected from the group consisting of CD16, CD19, CD20, CD37, CD40, CD45RO, CD80, CD86, CD137, CD152, and L6. In certain further embodiments of the above described fusion protein construct, the binding domain comprises, consists essentially of, or consists of, an scFv and the scFv contains a linker polypeptide that comprises, consists essentially of, or consists of, at least one polypeptide comprising or having as an amino acid sequence Gly-Gly-Gly-Gly-Ser [SEQ ID NO: 516], and in certain other embodiments the linker polypeptide comprises, consists essentially of, or consists of, at least three repeats of a polypeptide having as an amino acid sequence Gly-Gly-Gly-Gly-Ser [SEQ ID NO: 516]. In certain embodiments the immunoglobulin hinge region polypeptide comprises, consists essentially of, or consists of, a native or engineered human IgG, IgA, IgD hinge region polypeptide, or a native or engineered IgE CH2 region polypeptide. In certain embodiments the binding domain polypeptide comprises, consists essentially of, or consists of, a native or engineered CD154 extracellular domain. In certain embodiments the binding domain polypeptide comprises, consists essentially of, or consists of, a native or engineered CD154 extracellular domain and at least one a native or engineered immunoglobulin variable region polypeptide.

In other embodiments the invention provides an isolated polynucleotide encoding any of the constructs of the invention, for example, protein or polypeptide constructs of the invention including binding domain-immunoglobulin fusion proteins, and in related embodiments the invention provides a recombinant expression construct comprising such a polynucleotide, and in certain further
5 embodiments the invention provides a host cell transformed or transfected with, or otherwise containing, such a recombinant expression construct. In another embodiment the invention provides a method of producing a construct of the invention, for example, a protein or polypeptide construct of the invention such as a
10 binding domain-immunoglobulin fusion protein, comprising the steps of (a) culturing a host cell that has been transformed or transfected with, or otherwise made to contain, a polynucleotide construct of the invention under conditions that permit expression of the construct, for example, a construct encoding a binding domain-immunoglobulin fusion protein; and (b) isolating the construct, for example, the
15 binding domain-immunoglobulin fusion protein, from the host cell culture.

The inventions described and claimed herein include novel molecules useful, for example, as therapeutics and other purposes including diagnostic and research purposes. Such molecules have, for example, antigen-binding or other binding function(s) and one or more effector functions. DNA constructs of the
20 invention are useful in, for example, gene therapies, including *in vivo* and *ex vivo* gene therapies.

In one aspect, various constructs of the molecules of the invention include molecules comprising a "binding region", a "tail" region, and a "connecting" region that joins a binding region and a tail region.

25 Binding regions within the molecules of the invention may comprise, for example, binding domains for desired targets, including antigen-binding targets. Binding domains for antigen binding targets may comprise, for example, single chain Fvs and scFv domains. In certain embodiments, molecules of the invention may comprise a binding region having at least one immunoglobulin variable region
30 polypeptide, which may be a light chain or a heavy chain variable region polypeptide. In certain embodiments, molecules of the invention may comprise at least one such light chain V-region and one such heavy chain V-region and at least one linker peptide that connects the V-regions. ScFvs useful in the invention also include those

with chimeric binding or other domains or sequences. Other ScFvs useful in the invention also include those with humanized binding or other domains or sequences. In such embodiments, all or a portion of an immunoglobulin binding or other sequence that is derived from a non-human source may be "humanized" according to
5 recognized procedures for generating humanized antibodies, *i.e.*, immunoglobulin sequences into which human Ig sequences are introduced to reduce the degree to which a human immune system would perceive such proteins as foreign.

Example of scFvs useful in the invention, whether included as murine or other scFvs (including human scFvs), chimeric scFvs, or humanized scFvs, in
10 whole or in part, include anti-human CD20 scFvs (for example, "2H7" scFvs), anti-human CD37 scFvs (for example, "G28-1" scFvs), anti-human CD40 scFvs (for example, "G28-5" scFvs and "40.2.220" scFvs), anti-carcinoma-associated antigen scFvs (for example, "L6" scFvs), anti-CTLA-4 (CD152) scFvs (for example, "10A8" scFvs), anti-human CD28 scFvs (for example, "2E12" scFvs), anti-murine CD3 scFvs
15 (for example, "500A2" scFvs), anti-human CD3 scFvs (for example, G19-4 scFvs), anti-murine 4-1BB scFvs (for example, "1D8" scFvs), anti-human 4-1BB scFvs (for example, "5B9" scFvs), anti-human CD45RO (for example, "UCHL-1" scFvs), and anti-human CD16 (for example, "Fc2" scFvs) scFvs useful in the invention also include scFvs, including chimeric and humanized scFvs, having one or more amino
20 acid substitutions. A preferred amino acid substitution is at amino acid position 11 in the variable heavy chain (the V_H). Such a substitution may be referred to herein as " $X_{xx}V_{H11}Z_{xx}$ ". Thus, for example, where the normally occurring amino acid at position V_{H11} is a Leucine, and a Serine amino acid residue is substituted therefor, the substitution is identified as " $L V_{H11}S$ " or "Leu V_{H11} Ser." Other preferred
25 embodiments of the invention include molecules containing scFvs wherein the amino acid residue normally found at position V_{H11} is deleted. Still other preferred embodiments of the invention include molecules containing scFvs wherein the amino acid residues normally found at positions V_{H10} and/or V_{H11} and/or V_{H12} are substituted or deleted.

30 Other binding regions within the molecules of the invention may include domains that comprise sites for glycosylation, for example, covalent attachment of carbohydrate moieties such as monosaccharides or oligosaccharides.

Still other binding regions within molecules of the invention include polypeptides that may comprise proteins or portions thereof that retain the ability to specifically bind another molecule, including an antigen. Thus, binding regions may comprise or be derived from hormones, cytokines, chemokines, and the like; cell
5 surface or soluble receptors for such polypeptide ligands; lectins; intercellular adhesion receptors such as specific leukocyte integrins, selectins, immunoglobulin gene superfamily members, intercellular adhesion molecules (ICAM-1, -2, -3) and the like; histocompatibility antigens; and so on. Binding regions derived from such molecules generally will include those portions of the molecules necessary or desired
10 for binding to a target.

Certain constructs include binding regions that comprise receptor or receptor-binding domains. Receptor domains useful for binding to a target include, for example, a CD154 extracellular domain, or a CTLA-4 extracellular domain. In another example, the binding domain may include a first portion comprising,
15 consisting essentially or, or consisting of, a CD154 extracellular domain and a second portion comprising, consisting essentially or, or consisting of, at least one immunoglobulin variable region polypeptide, said second portion including, for example, an scFv or a V_H. Examples of other cell surface receptors that may comprise, consist essentially or, or consist of, or a portion of which may provide, a
20 binding region or binding domain polypeptide, include, for example, HER1, HER2, HER3, HER4, epidermal growth factor receptor (EGFR), vascular endothelial cell growth factor, vascular endothelial cell growth factor receptor, insulin-like growth factor-I, insulin-like growth factor-II, transferrin receptor, estrogen receptor, progesterone receptor, follicle stimulating hormone receptor (FSH-R), retinoic acid
25 receptor, MUC-1, NY-ESO-1, Melan-A/MART-1, tyrosinase, Gp-100, MAGE, BAGE, GAGE, any of the CTA class of receptors including in particular HOM-MEL-40 antigen encoded by the SSX2 gene, carcinoembryonic antigen (CEA), and PyLT. Additional cell surface receptors that may be sources of binding regions or binding domain polypeptides include, for example, CD2, 4-1BB, 4-1BB ligand, CD5, CD10,
30 CD27, CD28, CD152/CTLA-4, CD40, interferon- γ (IFN- $\gamma\gamma$), interleukin-4 (IL-4), interleukin-17 (IL-17) and interleukin-17 receptor (IL-17R). Still other cell surface receptors that may be sources of binding regions and/or binding domain polypeptides include, for example, CD59, CD48, CD58/LFA-3, CD72, CD70, CD80/B7.1,

CD86/B7.2, B7-H1/B7-DC, IL-17, CD43, ICOS, CD3 (*e.g.*, gamma subunit, epsilon subunit, delta subunit), CD4, CD25, CD8, CD11b, CD14, CD56, CD69 and VLA-4 ($\alpha_4\beta_7$). The following cell surface receptors are typically associated with B cells: CD19, CD20, CD22, CD30, CD153 (CD30 ligand), CD37, CD50 (ICAM-3), CD106 (VCAM-1), CD54 (ICAM-1), interleukin-12, CD134 (OX40), CD137 (41BB), CD83, and DEC-205. These lists are not exhaustive. Binding regions such as those set forth above may be connected, for example, by a native or engineered IgD hinge region polypeptide, preferably a human or humanized native or engineered IgD hinge region polypeptide. The invention thus further provides constructs that comprise, consist essentially of, or consist of, two binding regions, for example, an scFv and a cell surface receptor (or portion thereof), connected by a third molecule, for example, an IgD hinge region polypeptide as described herein.

Various molecules of the invention described and claimed herein include a connecting region joining one end of the molecule to another end. Such connecting regions may comprise, for example, immunoglobulin hinge region polypeptides, including any hinge peptide or polypeptide that occurs naturally. A connecting region may also include, for example, any artificial peptide or other molecule (including, for example, non-peptide molecules, partial peptide molecules, and peptidomimetics, *etc.*) useful for joining the tail region and the binding region. These may include, for example, alterations of molecules situated in an immunoglobulin heavy chain polypeptide between the amino acid residues responsible for forming intrachain immunoglobulin-domain disulfide bonds in CH1 and CH2 regions. Naturally occurring hinge regions include those located between the constant region domains, CH1 and CH2, of an immunoglobulin. Useful immunoglobulin hinge region polypeptides include, for example, human immunoglobulin hinge region polypeptides and llama or other camelid immunoglobulin hinge region polypeptides. Other useful immunoglobulin hinge region polypeptides include, for example, nurse shark and spotted ratfish immunoglobulin hinge region polypeptides. Human immunoglobulin hinge region polypeptides include, for example, wild type IgG hinges including wild-type human IgG1 hinges, human IgG-derived immunoglobulin hinge region polypeptides, a portion of a human IgG hinge or IgG-derived immunoglobulin hinge region, wild-type human IgA hinge region polypeptides, human IgA-derived immunoglobulin

hinge region polypeptides, a portion of a human IgA hinge region polypeptide or IgA-derived immunoglobulin hinge region polypeptide, wild-type human IgD hinge region polypeptides, human Ig-D derived immunoglobulin hinge region polypeptides, a portion of a human IgD hinge region polypeptide or IgD-derived immunoglobulin hinge region polypeptide, wild-type human IgE hinge-acting region, *i.e.*, IgE CH2 region polypeptides (which generally have 5 cysteine residues), human IgE-derived immunoglobulin hinge region polypeptides, a portion of a human IgE hinge-acting region, *i.e.*, IgE CH2 region polypeptide or IgE-derived immunoglobulin hinge region polypeptide, and so on. A polypeptide "derived from" or that is "a portion or fragment of" an immunoglobulin polypeptide chain region regarded as having hinge function has one or more amino acids in peptide linkage, for example 15-115 amino acids, preferably 95-110, 80-94, 60-80, or 5-65 amino acids, preferably 10-50, more preferably 15-35, still more preferably 18-32, still more preferably 20-30, still more preferably 21, 22, 23, 24, 25, 26, 27, 28 or 29 amino acids. Llama immunoglobulin hinge region polypeptides include, for example, an IgG1 llama hinge. The connecting region may comprise a stretch of consecutive amino acids from an immunoglobulin hinge region. For example, the connecting region can comprise at least five consecutive hinge region amino acids, at least ten consecutive hinge region amino acids, at least fifteen consecutive hinge region amino acids, at least 20 consecutive hinge region amino acids, and at least twenty five or more consecutive hinge region amino acids from human IgG hinge, human IgA hinge, human IgE hinge, camelid hinge region, IgG1 llama hinge region, nurse shark hinge region, and spotted ratfish hinge region, including for example an IgG₁ hinge region, a IgG₂ hinge region, a IgG₃ hinge region, an IgG₃ hinge region, and an IgG₄ hinge region.

Such connecting regions also include, for example, mutated or otherwise altered or engineered immunoglobulin hinge region polypeptides. A mutated or otherwise altered or engineered immunoglobulin hinge region polypeptide may comprise, consist essentially of, or consist of, a hinge region that has its origin in an immunoglobulin of a species, of an immunoglobulin isotype or class, or of an immunoglobulin subclass that is the same or different from that of any included native or engineered CH2 and CH3 domains. Mutated or otherwise altered or engineered immunoglobulin hinge region polypeptides include those derived or constructed from, for example, a wild-type immunoglobulin hinge region that contains one or more

cysteine residues, for example, a wild-type human IgG or IgA hinge region that naturally comprises three cysteines. In such polypeptides the number of cysteine residues may be reduced by amino acid substitution or deletion or truncation, for example. These polypeptides include, for example, mutated human or other IgG1 or IgG4 hinge region polypeptides containing zero, one, or two cysteine residues, and mutated human or other IgA1 or IgA2 hinge region polypeptides that contain zero, one, or two cysteine residues. Mutated or otherwise altered or engineered immunoglobulin hinge region polypeptides include those derived or constructed from, for example, a wild-type immunoglobulin hinge region that contains three or more cysteine residues, for example, a wild-type human IgG2 hinge region (which has 4 cysteines) or IgG4 hinge region (which has 11 cysteines). Mutated or otherwise altered or engineered immunoglobulin hinge region polypeptides include those derived or constructed from, for example, an IgE CH2 wild-type immunoglobulin region that generally contains five cysteine residues. In such polypeptides the number of cysteine residues may be reduced by one or more cysteine residues by amino acid substitution or deletion or truncation, for example. Also included are an altered hinge region polypeptides in which cysteine residues in the hinge region are substituted with serine or one or more other amino acids that are less polar, less hydrophobic, more hydrophilic, and/or neutral. Such mutated immunoglobulin hinge region polypeptides include, for example, mutated hinge region polypeptides that contain one cysteine residue and that are derived from a wild-type immunoglobulin hinge region polypeptide having two or more cysteine residues, such as a mutated human IgG or IgA hinge region polypeptide that contains one cysteine residue and that is derived from a wild-type human IgG or IgA region polypeptide. Connecting region polypeptides include immunoglobulin hinge region polypeptides that are compromised in their ability to form interchain, homodimeric disulfide bonds.

Mutated immunoglobulin hinge region polypeptides also include mutated hinge region polypeptides that exhibit a reduced ability to dimerize, relative to a wild-type human immunoglobulin G hinge region polypeptide, and mutated hinge region polypeptides that allow expression of a mixture of monomeric and dimeric molecules. Mutated immunoglobulin hinge region polypeptides also include hinge region polypeptides engineered to contain a glycosylation site. Glycosylation sites include, for example, an asparagine-linked glycosylation site, an O-linked

glycosylation site, a C-mannosylation site, a glypiation site, and a phosphoglycation site.

Specific connecting regions useful in molecules of the invention described and claimed herein include, for example, the following 18 amino acid sequences, DQEPKSCDKTHTCPPCPA, DQEPKSSDKTHTSPPPSPA, and DLEPKSCDKTHTCPPCPA. Other specific connecting regions include, for example, the mutant hinges within the sequences referred to herein as "2H7 scFv (SSS-S)H WCH2 WCH3" and "2H7 scFv (CSS)H WCH2 WCH3", and the human IgA-derived hinge referred to herein as "2H7 scFv IgAH WCH2 WCH3".

Tail regions within the molecules of the invention may include heavy chain constant region immunoglobulin sequences. Tail regions may thus include, for example, a polypeptide having at least one of an immunoglobulin heavy chain CH2 constant region polypeptide and an immunoglobulin heavy chain CH3 constant region polypeptide. At least one of the immunoglobulin heavy chain CH2 constant region polypeptide and the immunoglobulin heavy chain CH3 constant region polypeptide may be derived from a human immunoglobulin heavy chain. Thus, for example, CH2 and/or CH3 polypeptides may be derived from human IgG, human IgA, or human IgD molecules. Tail regions may also include, for example, a polypeptide having at least one of an immunoglobulin heavy chain CH3 constant region polypeptide and an immunoglobulin heavy chain CH4 constant region polypeptide. At least one of the immunoglobulin heavy chain CH3 constant region polypeptide and the immunoglobulin heavy chain CH4 constant region polypeptide may be derived from a human immunoglobulin heavy chain. Thus, for example, CH3 and/or CH4 polypeptides may be derived from human IgE. An immunoglobulin heavy chain CH2 region polypeptide included within a molecule of the invention may, for example, be from the IgG1, IgG2, IgG3 and/or IgG4 subclasses. An immunoglobulin heavy chain CH3 region polypeptide included within a molecule of the invention may also, for example, be from the IgG1, IgG2, IgG3 and/or IgG4 subclasses. Additionally, both the immunoglobulin heavy chain CH2 region polypeptide and the immunoglobulin heavy chain CH3 region polypeptide included within a molecule of the invention may, for example, be from the IgG1, IgG2, IgG3 and/or IgG4 subclasses. In other molecules of the invention at least one of the immunoglobulin heavy chain constant region polypeptides selected from a CH2 constant region polypeptide and a CH3

constant region polypeptide is a human IgA constant region polypeptide. An immunoglobulin heavy chain CH2 region polypeptide included within a molecule of the invention may, for example, be from the IgA1 and/or IgA2 subclasses. An immunoglobulin heavy chain CH3 region polypeptide included within a molecule of the invention may also, for example, be from the IgA1 and/or IgA2 subclasses. Additionally, both the immunoglobulin heavy chain CH2 region polypeptide and the immunoglobulin heavy chain CH2 region polypeptide included within a molecule of the invention may, for example, be from the IgA1 and/or IgA2 subclasses. In still other molecules of the invention, the tail region may comprise or consist essentially of a CH2 and/or CH3 constant region polypeptide comprising a polypeptide from human IgA and/or human IgE. In other embodiments, for example, the tail region within a molecule of the invention may include an immunoglobulin heavy chain CH2 and/or CH3 constant region polypeptide that is a mutated (for example, a mutated IgA CH3 constant region polypeptide that is incapable of associating with a J chain in which, for example, the IgA CH3 constant region polypeptide is of human origin). The tail region may also comprise, consist essentially of, or consist of an extracellular portion of a protein from the TNF superfamily, for example, CD154.

For molecules of the invention intended for use in humans, these regions will typically be substantially or completely human to minimize a potential human immune responses against the molecules and to provide appropriate effector functions. In certain embodiments of the invention, for example, the tail region includes a human IgG1 CH3 region sequence, a wild-type IgA heavy chain constant region polypeptide sequence that is capable or incapable of associating with J chain.

In preferred embodiments of the invention, a CH1 domain is not included in the tail region of the molecule, and the carboxyl end of the binding region is joined to the amino terminus of a CH2 portion of a tail region either directly or indirectly. A binding region may be indirectly joined to a tail region, for example via a connecting region polypeptide or other connecting molecule.

The invention also includes molecules that have mutated CH2 and/or CH3 sequences within a tail region. For example, a molecule of the invention may include a mutated Fc domain that has one or more mutations introduced into the CH2, CH3 and/or CH4 domains. In certain embodiments of the invention, molecules may include an IgA CH3 constant region polypeptide such as a human IgA CH3 constant

region polypeptide in which two or more residues from the C-terminus have been deleted to yield a truncated CH3 constant region polypeptide. In other embodiments of the invention, molecules include a mutated human IgA CH3 constant region polypeptide that is incapable of associating with a J chain that comprises a C-terminal deletion of either four or 18 amino acids. However, the invention need not be so limited, such that molecules containing the mutated IgA CH3 constant region polypeptide may comprise a deletion of 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-25, 26-30 or more amino acids, so long as the fusion protein is capable of specifically binding an antigen and capable of at least one immunological activity such as ADCC, CDC or complement fixation. The invention also includes molecules containing a tail region that comprises a mutated IgA CH3 constant region polypeptide that is incapable of associating with a J chain by virtue of replacement of the penultimate cysteine, or by chemical modification of that amino acid residue, in a manner that prevents interchain disulfide bond formation.

Various molecules of the invention include, for example, a binding domain scFv- fusion protein having a binding domain polypeptide comprising, consisting essentially of, or consisting of, (a) at least one immunoglobulin light chain variable region polypeptide, (b) at least one immunoglobulin heavy chain variable region polypeptide, and at least one linker peptide that joins the polypeptide of (a) and the polypeptide of (b). Such polypeptides may, for example, be derived from human immunoglobulins or non-human immunoglobulins.

Thus, in one aspect, the invention includes a non-naturally occurring single chain protein and/or V_H protein and/or V_L protein, or a desired portion of any of the above, including a first polypeptide comprising a binding domain polypeptide capable of binding to a target molecule, a second polypeptide comprising a flexible or other desired linker attached to said first polypeptide, a third polypeptide comprising a tail region, for example, an N-terminally truncated immunoglobulin heavy chain constant region polypeptide (or desired portion thereof) attached to the second polypeptide. The flexible linker may comprise, consist essentially of, or consist of, an immunoglobulin hinge region or portion thereof that has been mutated or otherwise altered or engineered, for example, one that contains a number of cysteine residues that is less than the number of cysteine residues present in the wild type immunoglobulin hinge region or portion (for example, zero, one, or two cysteines in

the case of IgG1 or IgG4), and wherein said non-naturally occurring single-chain protein is capable of at least one immunological activity, for example, ADCC, CDC, and/or complement fixation. This protein may include a binding domain polypeptide which is a single chain Fv. Additionally, this protein may include a binding domain
5 polypeptide which is a single chain Fv wherein the heavy chain variable region of the single chain Fv has an amino acid deletion or substitution at one or more of amino acid positions 9, 10, 11, 12, 108, 110, and 112. The protein may also include a binding domain polypeptide which is a single chain Fv wherein the light chain variable region of the single chain Fv has an amino acid deletion or substitution at one
10 or more of amino acid positions 12, 80, 81, 83, 105, 106, and 107.

In another aspect, the invention includes a non-naturally occurring V_H protein, or a desired portion thereof, that comprises, consists essentially of, or consists of, alone or in combination with any other molecule or construct, a V_H region or portion thereof that has an amino acid deletion or substitution at one or more of amino
15 acid positions 9, 10, 11, 12, 108, 110, and 112 of said V_H region. Amino acids may be substituted with either naturally-occurring or non-naturally occurring amino acids, or any other desired useful molecule.

Also described and claimed are uses of V_H proteins, or desired portions thereof, that comprise, consist essentially of, or consist of, alone or in combination
20 with any other molecule or construct, a V_H region or portion thereof that has an amino acid deletion or substitution at one or more of amino acid positions 9, 10, 11, 12, 108, 110, and 112 of said V_H region. Such uses include uses in phage display, yeast display, and ribosome display systems and methods.

In yet another aspect, the invention includes a non-naturally occurring
25 V_L protein, or a desired portion thereof, that comprises, consists essentially of, or consists of, alone or in combination with any other molecule, a V_L region or portion thereof that has an amino acid deletion or substitution at one or more of amino acid positions 12, 80, 81, 83, 105, 106, and 107 of said V_L region. Amino acids may be substituted with either naturally-occurring or non-naturally occurring amino acids, or
30 any other desired useful molecule.

Also described and claimed are uses of V_L proteins, or desired portions thereof, that comprises, consists essentially of, or consists of, alone or in combination

with any other molecule, a V_L region or portion thereof that has an amino acid deletion or substitution at one or more of amino acid positions 12, 80, 81, 83, 105, 106, and 107 of said V_L region. Such uses include uses in phage display, yeast display, and ribosome display systems and methods.

5 In yet another aspect, the invention includes a molecule comprising, consisting essentially of, or consisting of, (1) a V_H protein, or a desired portion thereof, wherein the V_H protein or portion thereof has an amino acid deletion or substitution at one or more of amino acid positions 9, 10, 11, 12, 108, 110, and 112, and (2) a non-naturally occurring V_L protein, or a desired portion thereof, alone or in
10 combination with any other molecule, wherein the V_L protein or portion thereof has an amino acid deletion or substitution at one or more of amino acid positions 12, 80, 81, 83, 105, 106, and 107. Amino acids may be substituted with either naturally-occurring or non-naturally occurring amino acids, or any other desired useful molecule.

 Also described and claimed are uses of a molecule comprising,
15 consisting essentially of, or consisting of, (1) a V_H protein, or a desired portion thereof, wherein the V_H protein or portion thereof has an amino acid deletion or substitution at one or more of amino acid positions 9, 10, 11, 12, 108, 110, and 112, and (2) a non-naturally occurring V_L protein, or a desired portion thereof, alone or in
20 combination with any other molecule, wherein the V_L protein or portion thereof has an amino acid deletion or substitution at one or more of amino acid positions 12, 80, 81, 83, 105, 106, and 107. Such uses include uses in phage display, yeast display, and ribosome display systems and methods.

 The invention also includes molecular constructs wherein the binding domain is a single chain Fv and the heavy chain variable region of said single chain
25 Fv has an amino acid substitution at amino acid position 11. The amino acid substituted for the amino acid at position of 11 of the single chain Fv heavy chain variable region may be selected from the group consisting of serine, threonine, tyrosine, asparagine, glutamine, aspartic acid, glutamic acid, lysine, arginine, and histidine. The invention thus includes, for example, a construct wherein the binding
30 domain is a single chain Fv and the heavy chain variable region of said single chain Fv has a serine amino acid substitution at amino acid position 11. Other amino acid position changes, substitutions, and deletions, are noted herein.

The invention also includes, for example, a construct wherein the binding domain is a single chain Fv and the amino acid at position 10 and/or 11 of the heavy chain variable region of said single chain Fv has been deleted.

5 In another aspect, the invention includes constructs wherein the binding region binds to a tumor or tumor-associated antigen. The binding region of a construct of the invention may bind, for example, to a cancer cell antigen. Cancer cell antigens to which constructs of the invention bind include cancer cell surface antigens and intracellular cancer cell antigens.

10 In yet another aspect, the invention includes a construct wherein the binding region binds to an antigen on an immune effector cell.

In another aspect, the invention includes a construct wherein the binding region binds to a B cell antigen including, for example, a B cell antigen selected from the group consisting of CD19, CD20, CD22, CD37, CD40, CD80, and CD86. Constructs of the invention that bind to such B cell antigens include, for
15 example, binding regions comprising an single chain Fv. Examples of such single chain Fv binding regions include molecules comprising or consisting essentially of single chain Fvs selected from the group consisting of HD37 single chain Fv, 2H7 single chain Fv, G28-1 single chain Fv, and 4.4.220 single chain Fv. Other examples include a binding region comprising, consisting essentially of, or consisting of, an
20 extracellular domain of CTLA-4.

In another aspect, the invention includes a construct wherein the binding region binds to a B cell differentiation antigen. B cell differentiation antigens include, for example, CD19, CD20, CD21, CD22, CD23, CD37, CD40, CD45RO, CD80, CD86, and HLA class II.

25 In another aspect, the invention includes a construct wherein the binding region binds to a target selected from the group consisting of CD2, CD3, CD4, CD5, CD6, CD8, CD10, CD11b, CD14, CD19, CD20, CD21, CD22, CD23, CD24, CD25, CD28, CD30, CD37, CD40, CD43, CD50 (ICAM3), CD54 (ICAM1), CD56, CD69, CD80, CD86, CD134 (OX40), CD137 (41BB), CD152 (CTLA-4),
30 CD153 (CD30 ligand), CD154 (CD40 ligand), ICOS, L6, B7-H1, and HLA class II.

The invention also includes protein constructs having a binding region, a tail region, and a connecting region, wherein the protein construct is capable of existing in solution as a monomer or in substantially monomeric form.

5 The invention also includes protein constructs having a binding region, a tail region, and a connecting region, wherein the protein construct is capable of forming a complex comprising two or more of said protein constructs including, for example, wherein said complex is a dimer.

10 In another aspect, constructs of the invention are capable of participating in or inducing or eliciting or helping to induce or elicit, directly or indirectly, at least one immunological activity selected from the group consisting of antibody dependent cell-mediated cytotoxicity, complement-dependent cytotoxicity (or complement-mediated lysis), complement fixation, induction of apoptosis, induction of one or more biologically active signals, induction of one or more immune effector cells, activation of cellular differentiation, cellular activation, release of one
15 or more biologically active molecules, and neutralization of an infectious agent or toxin.

In another aspect, binding constructs of the invention are capable of induction of biologically active signals by activation or inhibition of one or more molecules selected from the group consisting of protein kinases, protein phosphatases, G-proteins, cyclic nucleotides or other second messengers, ion channels, and
20 secretory pathway components. Such biologically active molecules are, for example, proteases. Other biologically active molecules are, for example, cytokines, including by way of example monokines, lymphokines, chemokines, growth factors, colony stimulating factors, interferons, and interleukins.

25 In another aspect, constructs of the invention are capable of induction, or participation in the induction, of one or more immune effector cells selected from the group consisting of NK cells, monocytes, macrophages, B cells, T cells, mast cells, neutrophils, eosinophils, and basophils.

30 In another aspect, constructs of the invention are capable of induction, or participation in the induction, of one or more immune effector cells that results in antibody dependent cell-mediated cytotoxicity or the release of one or more biologically active molecules.

In another aspect, constructs of the invention are capable of participating in and/or initiating apoptosis within target cells, for example, by activating one or more signalling mechanisms or molecules.

5 In another aspect, constructs of the invention are capable of induction, or participation in the induction, of cellular activation, wherein said activation leads to changes in cellular transcriptional activity. In one embodiment, cellular transcriptional activity is increased. In another embodiment, cellular transcriptional activity is decreased.

10 In another aspect, constructs of the invention having tail regions comprising, consisting essentially of, or consisting of, constant regions from IgA or IgE molecules, are capable of induction, or participation in the induction, of degranulation of neutrophils and/or mast cells.

In another aspect, constructs of the invention are capable of promotion, or participation in the promotion, of neutralization of an infectious agent, wherein said
15 infectious agent is, for example, a bacterium, a virus, a parasite, or a fungus.

In another aspect, constructs of the invention are capable of promoting, or participating in the promotion of, neutralization of a toxin, wherein said toxin is selected from the group consisting of endotoxins and exotoxins. Such toxins include, for example, exotoxins selected from the group consisting of anthrax toxin, cholera
20 toxin, diphtheria toxin, pertussis toxin, *E. coli* heat-labile toxin LT, *E. coli* heat stable toxin ST, shiga toxin *Pseudomonas* Exotoxin A, botulinum toxin, tetanus toxin, *Bordetella pertussis* AC toxin, and *Bacillus anthracis* EF toxin. Other toxins include, for example, saxitoxins, tetrodotoxin, mushroom toxins (amatoxins, gyromitrin, orellanine, *etc.*), aflatoxins, pyrrolizidine alkaloids, phytohemagglutinins, and
25 grayanotoxins.

In another aspect, constructs of the invention are capable of binding to an intracellular target to, for example, effect (or participate in effecting) a cellular function. Such constructs include, for example, constructs that include a tail region comprising, consisting essentially of, or consisting of, a native or engineered IgA
30 CH2 domain region and a native or engineered IgA CH3 domain region, said tail region being capable of binding J chain. Such a tail region is found, for example, in the 2H7 scFv IgAH WlgACH2 WCH3+JChain construct. Thus, the invention includes

constructs having, for example, an "Anti-Intracellular Target" binding domain (for example, and "Anti-Intracellular Target" scFv), a connecting region, and a native or engineered IgA constant region capable of binding J chain (for example, WIgACH2 WCH3).

5 In still another aspect, constructs of the invention include a molecule wherein an N-terminally immunoglobulin heavy chain constant region polypeptide comprises an IgG CH2 constant region polypeptide attached to an immunoglobulin heavy chain IgG CH3 constant region polypeptide.

10 In yet another aspect, the invention includes a method of reducing a target cell population in a subject comprising administering to said subject a therapeutically effective amount of a protein molecule that is less than about 120 kK, or less than about 150 kD, as measured, for example, by HPLC and non-reducing gels and consists essentially of (a) a first protein or peptide molecule that is capable of binding to cells within said target cell population, and (b) a second protein or peptide molecule that is capable of (i) binding to an Fc receptor and/or (ii) inducing target cell apoptosis, and/or (iii) fixing complement, wherein said first protein or peptide molecule is directly connected to said second protein or peptide molecule, or, optionally, said first protein or peptide molecule and said second protein or peptide molecule are linked by a third protein or peptide molecule, wherein said protein molecule is not an antibody, a member of the TNF family or the TNF receptor family, and is not conjugated with a bacterial toxin, a cytotoxic drug, or a radioisotope.

Various specific constructs of the invention include, by way of example only, the following:

1. 2H7 scFv VH L11S (CSC-S)H WCH2 WCH3
- 25 2. 2H7 scFv VH L11 S IgE CH2 CH3 CH4
3. 2H7 scFv VH L11S mIgE CH2 CH3 CH4
4. 2H7 scFv VH L11S mIgAH WIgACH2 T4CH3
5. 2H7 scFv VH L11S(SSS-S)H K322S CH2 WCH3
6. 2H7 scFv VH L11S (CSS-S)H K322S CH2 WCH3
- 30 7. 2H7 scFv VH L11S (SSS-S)H P331S CH2 WCH3

	8. 2HU scFv VH L11S(CSS-S)H P331S CH2 WCH3
	9. 2H7 scFv VH L11S (SSS-S)H T256N CH2 WCH3
	10. 2H7 scFv VH L11S (SSS-S)H RTPE/QNAK (255-258) CH2
	WCH3
5	11. 2H7 scFv VH L11S (SSS-S)H K290Q CH2 WCH3
	12. 2H7 scFv VH L11S (SSS-S)H A339P CH2 WCH3
	13. G28-1 scFv (SSS-S)H WCH2 WCH3
	14. G28-1 scFv IgAH WCH2 WCH3
	15. G28-1 scFv VH L11S (SSS-S)H WCH2 WCH3
10	16. G28-1 scFv VH L11S (CSS-S)H WCH2 WCH3
	17. G28-1 scFv VH L11S (CSC-S)H WCH2 WCH3
	18. G28-1 scFv VH L11S (SSC-P)H WCH2 WCH3
	19. CTLA4 (SSS-S)H P238SCH2 WCH32
	20. CTLA4 (CCC-P) WH WCH2 WCH3
15	21. FC2-2 scFv (SSS-S)H WCH2 WCH3
	22. FC2-2 scFv VHL11S (SSS-S)H WCH2 WCH3
	23. UCHL-1 scFv (SSS-S)H WCH2 WCH3
	24. UCHL-1 scFv L VH 11S (SSS-S)H WCH2 WCH3
	25. 5B9 scFv (SSS-S)H WCH2 WCH3
20	26. 5B9 scFv VHL11S (SSS-S)H WCH2 WCH3
	27. 2H7 scFv (SSS-S)H WCH2 WCH3
	28. 2H7 scFv (SSS-S)H P238SCH2 WCH3
	29. 2H7 scFv IgAH WCH2 WCH3
	30. 2H7 scFv IgAH WIgACH2 T4CH3
25	31. 2H7 scFv IgAH WIgACH2 WCH3+JChain
	32. 2H7 scFv (CCC-P) WH WCH2 WCH3

33. 2H7 scFv (SSS-S)H WCH2 F405YCH3
34. 2H7 scFv (SSS-S)H WCH2 F405ACH3
35. 2H7 scFv (SSS-S)H WCH2 Y407ACH3
36. 2H4 scFv (SSS-S) HWCH2 F405A, Y407ACH3
- 5 37. 2H7 scFv (CSS-S)H WCH2 WCH3
38. 2H7 scFv (SCS-S)H WCH2 WCH3
39. 2H7 scFv (SSC-P)H WCH2 WCH3
40. 2H7 scFv (CSC-S)H WCH2 WCH3
41. 2H7 scFv (CCS-P)H WCH2 WCH3
- 10 42. 2H7 scFv (SCC-P)H WCH2 WCH3
43. 2H7 scFv VH L11S (SSS-S)H WCH2 WCH3
44. 2H7 scFv VH L11S (CSS-S)H WCH2 WCH3
45. G28-1 scFv VH L11S (SCS-S)H WCH2 WCH3
46. G28-1 scFv VH L11S (CCS-P)H WCH2 WCH3
- 15 47. G28-1 scFv VH L11S (SCC-P)H WCH2 WCH3
48. G28-1 scFv VH L11S migE CH2 CH3 CH4
49. G28-1 scFv VH L11S mIgAH WlgACH2 T4CH3
50. G28-1 scFv VH L11S hIgE CH2 CH3 CH4
51. G28-1 scFv VH L11S hIgAH WlgACH2 T4CH3
- 20 52. HD37 scFv IgAH WCH2 WCH3
53. HD37 scFv (SSS-S)H WCH2 WCH3
54. HD37 scFv VH L11S(SSS-S)H WCH2 WCH3
55. L6 scFv IgAH WCH2 WCH3
56. L6 scFv L VH 11S (SSS-S)H WCH2 WCH3
- 25 57. 2H7 scFv-llama IgG1
58. 2H7 scFv-llama IgG2

	59. 2H7 scFv-Illama IgG3
	60. CD16-6 low (ED)(SSS-S)H P238SCH2 WCH3
	61. CD16-9 high (ED)(SSS-S)H P238SCH2 WCH3
	62. 2e12 scFv (SSS-s)H P238SCH2 WCH3--hCD80TM/CT
5	63. 10A8 scFv (SSS-s)H P238SCH2 WCH3--hCD80TM/CT
	64. 40.2.36 scFv (SSS-s)H P238SCH2 WCH3--hCD80TM/CT
	65. 2H7 scFv (SSS-s)H P238SCH2 WCH3--hCD80TM/CT
	66. G19-4 scFv (SSS-s)H P238SCH2 WCH3--hCD80TM/CT
	67. 2e12 scFv (SSS-s)H WCH2 WCH3--hCD80TM/CT
10	68. 2e12 scFv IgAH IgACH2 T4CH3--hCD80TM/CT
	69. 2e12 scFv IgE CH2CH3CH4--hCD80TM/CT
	70. 2e12 scFv (SSS-s)H P238SCH2 WCH3--mFADD-TM/CT
	71. 2e12 scFv (SSS-s)H WCH2 WCH3--mFADD-TM/CT
	72. 2e12 scFv (SSS-s)H WCH2 WCH3--mCasp3-TM/CT
15	73. 2e12 scFv (SSS-s)H P238SCH2 WCH3--mCasp3-TM/CT
	74. 2e12 scFv (SSS-s)H WCH2 WCH3--mCasp8-TM/CT
	75. 2e12 scFv (SSS-s)H P238SCH2 WCH3--mCasp8-TM/CT
	76. 2e12 scFv (SSS-s)H WCH2 WCH3--hCasp3-TM/CT
	77. 2e12 scFv (SSS-s)H P238SCH2 WCH3--hCasp3-TM/CT
20	78. 2e12 scFv (SSS-s)H WCH2 WCH3--hCasp8-TM/CT
	79. 2e12 scFv (SSS-s)H P238SCH2 WCH3--hCasp8-TM/CT
	80. 1D8 scFv-hIgG1 (SSS-s)H P238SCH2 WCH3--hCD80TM/CT
	81. 1D8 scFv-hIgG1 (SSS-s)H WCH2 WCH3--hCD80TM/CT
	82. 1D8 scFv-mIgAT4--hCD80TM/CT
25	83. 1D8 scFv-hIgE--hCD80TM/CT
	84. 1D8 scFv-hIgG1 (SSS-s)H P238SCH2 WCH3--mFADD-TM/CT

85. 1D8 scFv-hIgG1 (SSS-s)H WCH2 WCH3--mFADD-TM/CT
86. 1D8 scFv-hIgG1 (SSS-s)H WCH2 WCH3--mcasep3-TM/CT
87. 1D8 scFv-hIgG1 (SSS-s)H P238SCH2 WCH3--mcasep3-TM/CT
88. 1D8 scFv-hIgG1 (SSS-s)H WCH2 WCH3--mcasep8-TM/CT
- 5 89. 1D8 scFv-hIgG1 (SSS-s)H P238SCH2 WCH3--mcasep8-TM/CT
90. 1D8 scFv-hIgG1 (SSS-s)H WCH2 WCH3--hcasep3-TM/CT
91. 1D8 scFv-hIgG1 (SSS-s)H P238SCH2 WCH3--hcasep3-TM/CT
92. 1D8 scFv-hIgG1 (SSS-s)H WCH2 WCH3--hcasep8-TM/CT
93. 1D8 scFv-hIgG1 (SSS-s)H P238SCH2 WCH3--hcasep8-TM/CT L6
- 10 scFv (SSS-S)H WCH2 WCH3
94. 2H7 scFv CD1 54 (L2)
95. 2H7 scFv CD154 (S4)
96. CTLA4 IgAH IGACH2CH3
97. CTLA4 IgAH IgACH2 T4CH3
- 15 98. 2H7 scFv IgAH IgACH2CH3
99. 2H7 scFv IgAH IgAHCH2 TI 8CH3
100. 2H&-40.2.220 scFv (SSS-S)H WCH2 WCH3 (bispecific anti-ccd20-anti-cd40)
101. 2H7 scFv IgAH IgACH2 T4CH3-hCD89 TM/CT
- 20 102. G19-4 scFv (CCC-P) WH WCH2 WCH3-hCD89 TM/CT
103. 2e12 scFv (CCC-P) WH WCH2 WCH3-hCD89 TM/CT

These and other aspects of the present invention will become further apparent upon reference to the following detailed description and attached drawings. As noted herein, all referenced patents, articles, documents, and other materials

25 disclosed or identified herein are hereby incorporated by reference in their entireties as if each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows DNA and deduced amino acid sequences [SEQ ID NOs: 27 and 28] of 2H7scFv-Ig, a binding domain-immunoglobulin fusion protein capable of specifically binding CD20.

5 **Figure 2** shows production levels of 2H7 scFv-Ig by transfected, stable CHO lines and generation of a standard curve by binding of purified 2H7 scFv-Ig to CHO cells expressing CD20.

Figure 3 shows SDS-PAGE analysis of multiple preparations of isolated 2H7scFv-Ig protein.

10 **Figure 4** shows complement fixation (Figure 4A) and mediation of antibody-dependent cellular cytotoxicity (Figure 4B) by 2H7scFv-Ig.

Figure 5 shows the effect of simultaneous ligation of CD20 and CD40 on growth of normal B cells.

15 **Figure 6** shows the effect of simultaneous ligation of CD20 and CD40 on CD95 expression (Figure 6A) and induction of apoptosis (Figure 6B) in a B lymphoblastoid cell line.

Figure 7 shows DNA and deduced amino acid sequences of 2H7scFv-CD154 L2 (Figure 7A-7B, SEQ ID NOS: 690 and 691) and 2H7scFv-CD154 S4 (FIG. 7C-7D, SEQ ID NOS: 692 and 693) binding domain-immunoglobulin fusion proteins capable of specifically binding CD20 and CD40.

20 **Figure 8** shows binding of 2H7scFv-CD154 binding domain-immunoglobulin fusion proteins to CD20+CHO cells by flow immunocytofluorimetry.

25 **Figure 9** shows binding of Annexin V to B cell lines Ramos, BJAB, and T51 after binding of 2H7scFv-CD 154 binding domain-immunoglobulin fusion protein to cells.

Figure 10 shows effects on proliferation of B cell line T51 following binding of 2H7scFv-CD 154 binding domain-immunoglobulin fusion protein.

30 **Figure 11** depicts schematic representations of the structures of 2H7ScFv-Ig fusion proteins referred to as CytosB or CytosB derivatives: CytosB-

MHWTG1C (2H7 ScFv, mutant hinge, wild-type human IgG1 Fc domain), CytoxB-MHMG1C (2H7 ScFv, mutant hinge, mutated human IgG1 Fc domain) and CytoxB-IgAHWTHG1C (2H7 ScFv, human IgA-derived hinge (SEQ ID NO: 41), wild-type human IgG1 Fc domain). Arrows indicate position numbers of amino acid residues
5 believed to contribute to FcR binding and ADCC activity (heavy arrows), and to complement fixation (light arrows). Note absence of interchain disulfide bonds.

Figure 12 shows SDS-PAGE analysis of isolated CytoxB and 2H7scFv-CD 154 binding domain-immunoglobulin fusion proteins.

Figure 13 shows antibody dependent cell-mediated cytotoxicity
10 activity of CytoxB derivatives.

Figure 14 shows complement dependent cytotoxicity of CytoxB derivatives.

Figure 15 shows serum half-life determinations of CytoxB-MHWTG1C in macaque blood samples.

Figure 16 shows effects of CytoxB-MHWTG1C on levels of
15 circulating CD40+B cells in macaque blood samples.

Figure 17 shows production levels of HD37 (CD19-specific) ScFv-Ig by transfected mammalian cell lines and generation of a standard curve by binding of purified HD37 ScFv-Ig to cells expressing CD19.

Figure 18 shows production levels of L6 (carcinoma antigen) ScFv-Ig
20 by transfected, stable CHO lines and generation of a standard curve by binding of purified L6 ScFv-Ig to cells expressing L6 antigen.

Figure 19 shows antibody dependent cell-mediated cytotoxicity activity of binding domain-immunoglobulin fusion proteins 2H7 ScFv-Ig (Figure
25 19A), HD37 ScFv-Ig (Figure 19C) and G28-1 (CD37-specific) ScFv-Ig (Figure 19B).

Figure 20 shows antibody dependent cell-mediated cytotoxicity activity of L6 ScFv-Ig fusion proteins.

Figure 21 shows SDS-PAGE analysis of L6 ScFv-Ig and 2H7 ScFv-Ig fusion proteins.

Figure 22 shows SDS-PAGE analysis of G28-1 ScFv-Ig and HD37 ScFv-Ig fusion proteins.

Figure 23 presents a sequence alignment of immunoglobulin hinge and CH2 domains of human IgG1 (SEQ ID NO: 10 and 13) with the hinge and CH2 domains of llama IgG1 (SEQ ID NO: 4), IgG2 (SEQ ID NO: 6), and IgG3 (SEQ ID NO: 8).

Figure 24 illustrates migration of purified 2H7 scFv llama IgG fusion proteins in a 10% SDS polyacrylamide gel. Purified fusion proteins (5 µg per sample) were prepared in non-reducing sample buffer (lanes 2-5) and in reducing sample buffer (lanes 6-9). Lane 1: molecular weight markers (non-reduced); lanes 2 and 6: 2H7 scFv-llama IgG1 (DNA sequence is set forth in SEQ ID NO: 21; amino acid sequence is set forth in SEQ ID NO: 22); Lanes 3 and 7: 2H7 scFv-llama IgG2 (DNA sequence is set forth in SEQ ID NO: 23; amino acid sequence is set forth in SEQ ID NO: 24); lanes 4 and 8: 2H7 scFv-llama IgG3 (DNA sequence is set forth in SEQ ID NO: 25; amino acid sequence is set forth in SEQ ID NO: 26; and Lanes 5 and 9: Rituximab (chimeric anti-CD20 antibody (human IgG1 constant region)).

Figure 25 shows binding of 2H7 scFv-llama IgG1 (DNA sequence is set forth in SEQ ID NO: 21; amino acid sequence is set forth in SEQ ID NO: 22), 2H7 scFv-llama IgG2 (DNA sequence is set forth in SEQ ID NO: 23; amino acid sequence is set forth in SEQ ID NO: 24), and 2H7 scFv-llama IgG3 (DNA sequence is set forth in SEQ ID NO: 25; amino acid sequence is set forth in SEQ ID NO: 26) to CD20+CHO cells detected by flow immunocytometry.

Figure. 26 depicts CDC activity of 2H7 scFv llama IgG fusion proteins, 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, and 2H7 scFv-llama IgG3, and 2H7 scFv human IgG1 (2H7 scFv IgG WTH WTCH2CH3) (DNA sequence is set forth in SEQ ID NO: 27; amino acid sequence is set forth in SEQ ID NO: 28) against BJAB cells in the presence of rabbit complement. Rituximab was included as a control.

Figure 27 shows antibody dependent cell-mediated cytotoxicity activity of 2H7 scFv llama IgG fusion proteins, 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, and 2H7 scFv-llama IgG3. Effector cells (human PBMC) were combined with target cells (BJAB cells) at three different ratios, 1:25, 1:50, and 1:100.

Rituximab was included as a control. Each data point represents three separate measurements.

Figure 28 shows antibody dependent cell-mediated cytotoxicity activity of 2H7 scFv llama IgG fusion proteins, 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, and 2H7 scFv-llama IgG3. Effector cells (llama PBMC) were combined with target cells (BJAB cells) at three different ratios, 1:25, 1:50, and 1:100. Rituximab was included as a control. Each data point represents three separate measurements.

Figure 29 depicts complement dependent cytotoxicity activity of Reh cells (acute lymphocytic leukemia) expressing scFv-Ig fusion proteins on the cell surface. Reh cells were transfected with constructs encoding scFv antibodies specific for human costimulatory molecules, CD152, CD28, CD40, and CD20, fused to human IgG1 wild-type hinge-CH2-CH3, which was fused to human CD80 transmembrane and cytoplasmic tail domains. Complement dependent cytotoxicity activity was measured in the presence and absence of rabbit complement (plus C' and no C', respectively). The data represent the average of duplicate samples. Reh anti-hCD 152 scFvIg: 10A8 scFv (SEQ ID NO: 53); Reh anti-hCD28scFvIg: 2E12 scFv (SEQ ID NO: 51); Reh anti-hCD40scFvIg: 4.2.220 scFv (SEQ ID NO: 49); and Reh anti-hCD20scFvIg: 2H7 scFv IgG (SEQ ID NO: 57 and 29).

Figure 30 presents antibody dependent cell-mediated cytotoxicity activity of Reh cells that were transfected with constructs encoding scFv antibodies specific for human costimulatory molecules, CD152, CD28, CD40, and CD20, as described for Figure 29, and for murine CD3, fused to human mutant IgG1 hinge and mutant CH2 and wild type CH3 (Reh anti-mCD3scFv designating Reh cells transfected with polynucleotide 500A2 scFv IgG MTH (SSS) MTCH2WTCH3 (SEQ ID NO: 55), which was fused to human CD80 transmembrane and cytoplasmic tail domains. The data represent the average of quadruplicate samples.

Figure 31 lists immunoglobulin constant region constructs that were used in experiments illustrated in subsequent figures.

Figure 32 depicts complement dependent cytotoxicity activity of CTLA-4 Ig fusion proteins, CTLA-4 IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 78) (2 µg/ml) and CTLA-4 IgG MTH MTCH2WTCH3 (SEQ ID NOs: 530 and 86) (2

µg/ml), in the presence and absence of rabbit complement (plus C' and no C', respectively). The target cells were Reh cells and Reh cells transfected with CD80 (Reh CD80.10).

Figure 33 shows antibody dependent cell-mediated cytotoxicity activity of CTLA-4 Ig fusion proteins, CTLA-4 IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 78) (2 µg/ml) and CTLA-4 IgG MTH MTCH2WTCH3 (SEQ ID NO: 530 or 86) (2 µg/ml). Effector cells, human PBMC, were added to target cells, Reh or Reh CD80.1, at the ratios indicated. Figure 33A presents the level of natural killing in Reh CD80.1 cells in the absence of any Ig fusion protein. Figure 33B presents antibody dependent cell-mediated cytotoxicity mediated by CTLA-4 IgG MTH MTCH2WTCH3, and FIGURE 33C presents antibody dependent cell-mediated cytotoxicity mediated by CTLA-4 IgG WTH (CCC) WTCH2CH3. Each data point represents the average percent specific killing measured in four sample wells.

Figure 34 illustrates binding of 2H7 (anti-CD20) scFv Ig fusion proteins to (CD20+) CHO cells by flow immunocytofluorimetry.

Figure 35 presents an immunoblot of 2H7 scFv IgG and IgA fusion proteins. COS cells were transiently transfected with various 2H7 scFv Ig fusion protein constructs. The expressed polypeptides were immune precipitated with protein A, separated in a non-reducing SDS polyacrylamide gel, and then transferred to a polyvinyl fluoride membrane. Proteins were detected using an anti-human IgG (Fc specific) horseradish peroxidase conjugate. Lane 1: vector only; lane 2: 2H7 scFv IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 28); lane 3: 2H7 scFv IgG MTH (CSS) WTCH2CH3 (SEQ ID NO: 135); lane 4: 2H7 scFv IgG MTH (SCS) WTCH2CH3 (SEQ ID NO: 137); lane 5: 2H7 scFv IgAH IgG WTCH2CH3 (SEQ ID NO: 60); and lane 6: 2H7 scFv IgG MTH (SSS) WTCH2CH3 (SEQ ID NO: 50).

Figure 36 illustrates binding of 2H7 scFv IgAH IgACH2CH3 polypeptide (SEQ ID NO: 62) and 2H7 scFv IgAH IgAT4 (SEQ ID NO: 71) to (CD20+) CHO cells by flow immunocytofluorimetry. The source of the polypeptides was culture supernatants from transiently transfected COS cells. COS cells transfected with a plasmid comprising a sequence encoding 2H7 scFv IgAH IgACH2CH3 were co-transfected with a plasmid containing nucleotide sequence encoding human J chain.

Figure 37 illustrates antibody dependent cell-mediated cytotoxicity activity of anti-CD20 (2H7) scFv Ig fusion proteins against BJAB target cells using whole blood as the source of effector cells. Purified 2H7 scFv Ig fusion proteins were titrated and combined with ^{51}Cr -labeled BJAB cells (5×10^4) and whole blood (1:4
5 final dilution). Each data point represents the average percent specific killing measured in four sample wells.

Figure 38 demonstrates antibody dependent cell-mediated cytotoxicity activity of 2H7 scFv Ig fusion proteins (5 $\mu\text{g/ml}$) against ^{51}Cr -labeled BJAB cells at 0.25, 0.125, and 0.625 dilutions of whole blood. Each data point represents the
10 average percent specific killing measured in four sample wells.

Figure 39 shows a comparison of antibody dependent cell-mediated cytotoxicity activity of 2H7 scFv IgG MTH (SSS) WTCH2CH3 (5 $\mu\text{g/ml}$) and 2H7 scFv IgAH IgACH2CH3 (5 $\mu\text{g/ml}$) when human PBMC are the source of effector cells (Figure 39A) and when human whole blood is the source of effector cells (Figure
15 39B).

Figure 40 presents an immunoblot of 2H7 scFv IgG fusion proteins. COS cells were transiently transfected with various 2H7 scFv Ig fusion protein constructs. Culture supernatants containing the expressed polypeptides were separated in a non-reducing SDS polyacrylamide gel, and then were transferred to a polyvinyl
20 fluoride membrane. Proteins were detected using an anti-human IgG (Fc specific) horseradish peroxidase conjugate. Lanes 1-5: purified 2H7 scFv IgG MTH (SSS) WTCH2CH3 at 40 ng, 20 ng, 10 ng/5 ng, and 2.5 ng per lane, respectively. Culture supernatants were separated in lanes 6-9. Lane 6: 2H7 scFv IgG WTH (CCC) WTCH2CH3; lane 7: 2H7 scFv IgG MTH (CSS) WTCH2CH3; lane 8: 2H7 scFv IgG
25 MTH (SCS) WTCH2CH3; and lane 9: 2H7 scFv VHSE11 IgG MTH (SSS) WTCH2CH3. The molecular weight (kDa) of marker proteins is indicated on the left side of the immunoblot.

Figure 41 illustrates cell surface expression of 1D8 (anti-murine 4-1BB) scFv IgG WTH WTCH2CH3-CD80 fusion protein on K1735 melanoma cells
30 by flow immunofluorimetry (Fig. 41A). The scFv fusion protein was detected with phycoerythrin-conjugated F(ab')₂ goat anti-human IgG. Fig. 41B depicts growth of tumors in naive C3H mice transplanted by subcutaneous injection with wild type

K1735 melanoma cells (K1735-WT) or with K1735 cells transfected with 1D8 scFv IgG WTH WTCH2CH3-CD80 (K1735-1D8). Tumor growth was monitored by measuring the size of the tumor. Fig. 41C demonstrates the kinetics of tumor growth in naive C3H mice injected intraperitoneally with monoclonal antibodies to remove
5 CD8⁺, CD4⁺, or both CD4⁺ and CD8⁺ T cells prior to transplantation of the animals with K1735-1D8 cells.

Figure 42 demonstrates therapy of established K1735-WT tumors using K1735-1D8 as an immunogen. Six days after mice were transplanted with K1735-WT tumors, one group (five animals) was injected subcutaneously with
10 K1735-1D8 cells (open circles) or irradiated K1735-WT cells (solid squares) on the contralateral side. A control group of mice received PBS (open squares). Treatments were repeated on the days indicated by the arrows.

Figure 43 shows the growth of tumors in animals that were injected subcutaneously with 2×10^6 K1735-WT cells (solid squares) and the growth of
15 tumors in animals that were injected subcutaneously with 2×10^6 K1735-WT cells plus 2×10^5 K1735-1D8 cells (open triangles).

Figure 44 presents a flow cytometry analysis of antigen 104 murine sarcoma tumor cells transfected with 1D8 scFv IgG WTH WTCH2CH3-CD80 isolated after repeated rounds of panning against anti-human IgG. Transfected cells
20 expressing 1D8 scFv IgG WTH WTCH2CH3-CD80 were detected with fluorescein isothiocyanate (FITC)-conjugated goat anti-human IgG (depicted in black). Untransfected cells are shown in gray.

Figure 45 illustrates migration of various 2H7 scFv Ig fusion proteins in a 10% SDS-PAGE gel. 2H7 was the anti-CD20 scFv and 40.2.220 was the anti-
25 CD40 scFv. Lane 1: Bio-Rad prestained molecular weight standards; lane 2: anti-CD20 scFv IgG MTH (SSS) MTCH2WTCH3; lane 3: anti-CD20 scFv IgG MTH (SSS) WTCH2CH3; lane 4: 2H7 scFv IgAH IgG WTCH2CH3; lane 5: anti-CD20-anti-CD40 scFv IgG MTH (SSS) MTCH2WTCH3; lane 6: Rituximab; lane 7: Novex MULTIMARK® molecular weight standards.

Figure 46 illustrates effector function as measured in an antibody dependent cell-mediated cytotoxicity assay of 2H7 Ig fusion proteins that contain a mutant CH2 domain or wild type CH2 domain. The percent specific killing of BJAB
30

target cells in the presence of human PBMC effector cells by 2H7 scFv IgG MTH (SSS) MTCH2WCH3 (diamonds) was compared to 2H7 scFv IgG MTH (SSS) WCH2CH3 (squares) and 2H7 scFv IgAH IgG WCH2CH3 (triangles) and Rituximab (circles).

5 **Figure 47** shows cell surface expression of an anti-human CD3 scFv IgG WTH WCH2CH3-CD80 (SEQ ID NO: 443) fusion protein on Reh cells (Fig. 47A) and T51 lymphoblastoid cells (Fig. 47B) by measuring the linear fluorescent equivalent (LFE) using flow immunocytometry.

10 **Figure 48** presents the percent specific killing of untransfected Reh and T51 cells and the percent specific killing of Reh cells (Reh anti-hCD3) (Fig. 48A) and T51 cells (T51 anti-hCD3) (Fig. 48B) that were transfected with a construct encoding scFv antibodies specific for human CD3, fused to human IgG1 wild-type hinge-CH2-CH3, which was fused to human CD80 transmembrane and cytoplasmic tail domains. Human PBMC (effector cells) were combined with BJAB target cells at
15 the ratios indicated.

Figure 49 illustrates binding of 5B9, an anti-murine CD137 (4-1 BB) monoclonal antibody, and a 5B9 scFv IgG fusion protein (5B9 scFv IgG MTH (SSS) WCH2CH3 (SEQ ID NO: 133) to stimulated human PBMC. Binding of the 5B9 scFv IgG fusion protein was detected by flow immunocytometry using FITC
20 conjugated goat anti-human IgG. Binding of the 5B9 monoclonal antibody was detected with FITC conjugated goat anti-mouse IgG.

Figure 50 illustrates the effect of the LV_H11S mutation on the expression of 2H7 LV_H11S scFv WCH2 WCH3 ("CytosB scFv Ig"; SEQ ID NO: 369) in CHO cell lines.

25 **Figure 51** shows a semi-quantitative SDS-PAGE analysis examining the expression of 2H7 LV_H11S scFv WCH2 WCH3 (SEQ ID NO: 369) when transiently transfected in CHO cells. Lanes 2-5 are various amounts of 2H7 LV_H11S scFv WCH2 WCH3. Lanes 6-10 are 10 µl samples from five different clones expressing 2H7 LV_H11S scFv WCH2 WCH3.

30 **Figure 52** shows differences in binding capacity between a G28-1 LV_H11S scFv Ig construct (SEQ ID NO: 324) and a G28-1 wild type scFv Ig binding domain fusion protein construct (SEQ ID NO: 320), both obtained from transiently

transfected COS cells. Binding to Ramos cells was determined using flow cytometry. The data illustrates a significant increase in binding of the LV_H11S protein to CD37+Ramos cells.

Figure 53 illustrates increased levels of expression of a G28-1 LV_H11S scFv Ig construct (SEQ ID NO: 324) compared to a G28-1 wild type scFv Ig construct in COS. Protein levels were compared using immunoblot analysis. Both immunoblot gels have quantitated amounts of purified a G28-1 scFv Ig (SSS-S)H WCH2 WCH3 construct of the invention in lanes 1-4. Lanes 5-9 of the first immunoblot represent five different clones each transfected with G28-1 scFv (SSS-S)H WCH2 WCH3, while lanes 5-9 of the second immunoblot represent five different clones transfected with G28-1 LV_H11S scFv (SSS-S)H WCH2 WCH3. The immunoblots illustrate that the LV_H11S form causes the G28-1 scFv Ig construct to express at very high levels.

Figure 54 illustrates the binding of 2H7 scFv Ig derivatives with altered hinges (SEQ ID NOs: 370, 58, 28, 166, 137, and 135) to CHO cells expressing CD20 (CD20+CHO) by flow cytometry, and indicates that these altered connecting region hinge constructs (including (SSS-S), (CSS-S), (SCS-S) and (CSC-S) hinge regions) retain binding function to CD20.

Figure 55 shows the ability to mediate antibody dependent cell-mediated cytotoxicity of various constructs against Bjab targets: (A) 2H7 scFv Ig constructs of the invention that contain connecting regions comprising (CSS-S), (SCS-S), (CSC-S), and (SSS-S) hinges (SEQ ID NOs: 58, 28, 166, 137, 135, 60) and (B) 2H7 scFv constructs of the invention with various connecting regions and tail regions. Percent specific killing is compared to total killing induced by a detergent. The controls are natural killing in target cells with effectors added and a 2H7 construct with an IgA hinge connecting region and IgA-derived tail region that does not bind PBMC effectors.

Figure 56 illustrates the ability of various 2H7 scFv Ig constructs of the invention (SEQ ID NOs: 370, 58, 28, 166, 137, 135) that include connecting regions having various hinge regions (*e.g.*, (CSC-S), (SSS-S), (SCS-S), and (CSS-S)) to mediate complement activity in Ramos cells. Percent specific killing is measured

against the control of complement only, and 100% killing was determined by exposure of cells to detergent.

Figure 57 illustrates the shows the binding of 2H7 scFv Ig constructs of the invention containing different tail regions to CD20+CHO using immunocytofluorimetry. The different proteins were detected using FITC conjugated to anti-IgG, anti-IgA, and anti-IgE.

Figure 58A shows the binding of 2H7 V_H L1 SscFv IgECH2CH3CH4, purified using Hydrophobic charge induction chromatography (HCIC) and eluted at different pHs 4.0 and 3.5, in CD20+CHO cells by flow cytometry, indicating that the proteins bound CD20 whether eluted at pH 4.0 or 3.5. **Figure 58B** is a data graph indicating the ability of these 2H7 V_H L1 S scFv IgE constructs of the invention to mediate, for example, ADCC in Bjab target cells.

Figure 59 shows the binding capacity of 2H7 V_H L1 S mIgECH2CH3CH4 tailed and IgA tailed to CD20+CHO cells by flow cytometry.

Figure 60 shows the High Performance Liquid Chromatography (HPLC) profiles of various protein constructs of the invention (A) 2H7 scFv (SSS-S) H (P238S)CH2 WCH3 (B) 2H7 scFv (CSS-S)H WCH2 WCH3, (C) 2H7 scFv (SCS-S)H WCH2 WCH3, and (D) 2H7 scFv (SSS-S)H WCH2 (Y407A)CH3, indicating that construct A has apparent molecular weight forms of 100 kD and 75 kD and that, by introducing certain changes a predominant 75 kD molecular weight form is obtained, as seen in constructs B, C, and D. See Example 40.

Figure 61 shows the HPLC profiles of various protein constructs of the invention (A) 2H7 scFv (SSS-S)H WCH2 WCH3, (B) 2H7 scFv (CSC--S)H WCH2 WCH3, (C) 2H7 scFv (CCC-P)H WCH2 WCH3, and (D) 2H7 scFv IgAH WCH2 WCH3, indicating that construct A has apparent molecular weight forms of 100 kD and 75 kD and that, by introducing certain changes a predominant 75 kD molecular weight form is obtained, as seen in constructs B and C, while construct D (which has an IgA tail regaion) has an apparent molecular weight of 150 kD. See Example 40.

Figure 62 shows the HPLC profiles of various protein constructs of the invention (A) 2H7 scFv (SSS-S)H WCH2 WCH3, (B) 2H7 scFv (SCS-S) H WCH2 WCH3, (C) 2H7 scFv IgA 3TCH2 WCH3, and (D) 2H7 scFv (SSS-S)H

WCH2 (F405A Y407A)CH3, indicating that construct A has two forms with apparent molecular weights at 100 kD and 75 kD, construct B has a predominant form with an apparent molecular weight of 75 kD, while construct C with a T4 mutation leads to three forms with apparent molecular weights near 600 kD and construct D with a double point mutation in the CH3 region leads to a predominant form having an apparent molecular weight less than 44 kD. A T4 mutation here refers to a truncation of four amino acids from a CH3 region. See Example 40.

Figure 63 compares the effect on binding CD20+CHO cells by 2H7. VH L11S scFv Ig constructs, with and without F405a and Y407A alterations in the CH3 region, by flow cytometry, indicating a loss of binding capability with this double amino acid change. See Example 41.

Figure 64 shows the binding capacity of FITC conjugated 2H7 V_H L11S scFv Ig derivatives, to CH₂O+CHO cells by flow cytometry, indicating that these constructs do not lose binding capacity when conjugated to a fluorescent marker. See Example 41.

Figure 65 shows a nonreducing SDS-PAGE analysis examining 10 fg (per lane) of various purified 2H7 V_H L11S scFv Ig constructs of the invention, indicating an apparent molecular weight for each construct in reference to a standard molecular weight marker in lane 1. See Example 41.

Figure 66 compares the CH2 domain sequences of four different human IgG regions, hIgG1, hIgG2, hIgG3, hIgG4, hIgG4, and one rat region, rIgG2b. Point mutations affecting ADCC and CDC are labeled with arrows. See Example 52.

Figure 67 demonstrates the ability of various 2H7 V_H L11S scFv Ig constructs to mediate ADCC in CHO and Lec13 CHO transiently transfected cells, indicating that constructs expressed in Lec 13 CHO cells had a 20% increase in specific killing over the same construct expressed in regular CHO cells. See Example 42.

Figure 68 shows SDS-PAGE analysis, both reduced and nonreduced, of high and low affinity alleles of soluble CD 16(ED) (SSS-S)H P283S CH2 WCH3 (SEQ ID NOs: 422 and 426). See Example 43.

Figure 69 demonstrates the different binding capabilities of the high and low affinity CD16 fusion proteins to 2H7 V_H L11S scFv (CSC-S) WCH2 WCH3

or 2H7 V_H L11S scFv (SSS-S) WCH2 WCH3, and indicates a loss of high and low affinity allele binding using P238S CH2 constructs. See Example 43.

Figure 70 shows a diagram of (A) an assay used to detect changes in Fc receptor binding using FITC conjugated CD16 extracellular domain Ig fusion protein with a mutated tail, which eliminates self-association and B) a mammalian display system using cell surface expression of SMIPs. See Example 44.

Figure 71 shows the induction of apoptosis in Bjab and Ramos cells by various mAbs and scFvlg constructs of the invention. See Example 45.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to novel molecules useful, for example, as therapeutics, as well as for other purposes including diagnostic and research purposes. Such molecules have, for example, antigen-binding or other binding function(s) and, for example, one or more effector functions. The invention includes molecular constructs, including binding domain-immunoglobulin fusion proteins, and related compositions and methods, which will be useful in immunotherapeutic and immunodiagnostic applications, and in research methods, and which offer certain advantages over antigen-specific compounds and polypeptides of the prior art. The constructs, including fusion proteins, of the present invention are preferably single polypeptide chains that comprise, in pertinent part, the following fused or otherwise connected domains or regions: a binding region construct, such as a binding domain or polypeptide, a connecting region construct including, for example, a native or engineered immunoglobulin hinge region polypeptide, and a tail region construct, including, for example, a construct that may comprise, consist essentially of, or consist of, a native or engineered immunoglobulin heavy chain CH2 constant region polypeptide and a native or engineered immunoglobulin heavy chain CH3 constant region polypeptide. According to certain embodiments that are particularly useful for gene therapy, the constructs, including fusion proteins, of the present invention may further comprise a native or engineered plasma membrane anchor domain. According to certain other preferred embodiments the constructs, including fusion proteins, of the present invention may further include a tail region having a native or engineered immunoglobulin heavy chain CH4 constant region polypeptide. In particularly preferred embodiments, the binding regions, such as

polypeptide domains, of which the constructs, including binding domain-immunoglobulin fusion proteins, are comprised are, or are derived from, polypeptides that are the products of human gene sequences, but the invention need not be so limited and may in fact relate to constructs, including binding domain-immunoglobulin fusion proteins, as provided herein that are derived from any natural or artificial source, including genetically engineered and/or mutated polypeptides.

The present invention relates in part to the surprising observation that the novel constructs, including binding domain-immunoglobulin fusion proteins, described herein are capable of immunological activity. More specifically, these proteins retain the ability to participate in well known immunological effector activities including, for example, antibody dependent cell mediated cytotoxicity (*e.g.*, subsequent to antigen binding on a cell surface, engagement and induction of cytotoxic effector cells bearing appropriate Fc receptors, such as Natural Killer cells bearing FcRγIII, under appropriate conditions) and/or complement fixation in complement dependent cytotoxicity (*e.g.*, subsequent to antigen binding on a cell surface, recruitment and activation of cytolytic proteins that are components of the blood complement cascade) despite having structures not be expected to be capable of promoting such effector activities or to promotion of such activities as described herein. For reviews of ADCC and CDC see, *e.g.*, Carter, 2001 *Nat. Rev. Canc.* **1**: 118; Sulica *et al.*, 2001 *Int. Rev. Immunol.* **20**: 371; Maloney *et al.*, 2002 *Semin. Oncol.* **29**: 2; Sondel *et al.*, 2001 *Hematol Oncol Clin North Am* **15**(4): 703-21; Maloney 2001 *Anticanc. Drugs* **12** Suppl.2: 1-4. IgA activation of complement by the alternative pathway is described, for example, in Schneiderman *et al.*, 1990 *J. Immunol.* **145**: 233. As described in greater detail below, ADCC, complement fixation, and CDC are unexpected functions for constructs, including fusion proteins, comprising for example immunoglobulin heavy chain regions and having the structures described herein, and in particular for immunoglobulin fusion proteins comprising, for example, immunoglobulin hinge region polypeptides that are compromised in their ability to form interchain, homodimeric disulfide bonds.

Another advantage afforded by the present invention is constructs, including binding domain-immunoglobulin fusion polypeptides, of the invention that can be produced in substantial quantities that are typically greater than those routinely attained with single-chain antibody constructs of the prior art, for example. In

preferred embodiments, constructs, including the binding domain-immunoglobulin fusion polypeptides, of the present invention are recombinantly expressed in mammalian or other desired and useful expression systems, which offer the advantage of providing polypeptides that are stable *in vivo* (e.g., under physiological conditions).

5 According to non-limiting theory, such stability may derive in part from posttranslational modifications, and specifically glycosylation. Production of the constructs, including binding domain-immunoglobulin fusion protein constructs, of the invention via recombinant mammalian expression has been attained in static cell cultures at a level of greater than 50 mg protein per liter culture supernatant and has
10 been routinely observed in such cultures at 10-50 mg/liter, such that preferably at least 10-50 mg/liter may be produced under static culture conditions; also contemplated are enhanced production, in whole or in part, of the protein constructs of the invention using art-accepted scale-up methodologies such as "fed batch" (*i.e.*, non-static) production, where yields of at least 5-500 mg/l, and in some instances at least 0.5-1
15 gm/l, depending on the particular protein product, are obtained.

A construct, including a binding domain polypeptide, according to the present invention may be, for example, any polypeptide that possesses the ability to specifically recognize and bind to a cognate biological molecule or complex of more than one molecule or assembly or aggregate, whether stable or transient, of such a
20 molecule. Such molecules include proteins, polypeptides, peptides, amino acids, or derivatives thereof, lipids, fatty acids or the like, or derivatives thereof; carbohydrates, saccharides or the like or derivatives thereof; nucleic acids, nucleotides, nucleosides, purines, pyrimidines or related molecules, or derivatives thereof, or the like; or any combination thereof such as, for example, glycoproteins, glycopeptides, glycolipids,
25 lipoproteins, proteolipids; or any other biological molecule that may be present in a biological sample. Biological samples may be provided, for example, by obtaining a blood sample, biopsy specimen, tissue explant, organ culture, biological fluid or any other tissue or cell or other preparation from a subject or a biological source. The subject or biological source may, for example, be a human or non-human animal, a
30 primary cell culture or culture adapted cell line including but not limited to genetically engineered cell lines that may contain chromosomally integrated or episomal recombinant nucleic acid sequences, immortalized or immortalizable cell lines, somatic cell hybrid cell lines, differentiated or differentiable cell lines,

transformed cell lines and the like, *etc.* In certain preferred embodiments of the invention, the subject or biological source may be suspected of having or being at risk for having a disease, disorder or condition, including a malignant disease, disorder or condition or a B cell disorder, which in certain further embodiments may be an autoimmune disease, and in certain other embodiments of the invention the subject or biological source may be known to be free of a risk or presence of such disease, disorder or condition.

A binding region, including a binding domain polypeptide, for example, may be any naturally occurring, synthetic, semi-synthetic, and/or recombinantly produced binding partner for a biological or other molecule that is a target structure of interest, herein sometimes referred to as an "antigen" but intended according to the present disclosure to encompass any target biological or other molecule to which it is desirable to have the subject invention, for example, a fusion protein, bind or specifically bind. Constructs of the invention, including binding domain-immunoglobulin fusion proteins, are defined to be "immunospecific" or capable of binding to a desired degree, including specifically binding, if they bind a desired target molecule such as an antigen as provided herein, at a desired level, for example, with a K_a of greater than or equal to about 10^4 M^{-1} , preferably of greater than or equal to about 10^5 M^{-1} , more preferably of greater than or equal to about 10^6 M^{-1} and still more preferably of greater than or equal to about 10^7 M^{-1} . Affinities of even greater than about 10^7 M^{-1} are still more preferred, such as affinities equal to or greater than about 10^7 M^{-1} , about 10^8 M^{-1} , and about 10^9 M^{-1} , and about 10^{10} M^{-1} . Affinities of binding domain-immunoglobulin fusion proteins according to the present invention can be readily determined using conventional techniques, for example those described by Scatchard *et al.*, 1949 *Ann. NY Acad. Sci.* **51**: 660. Such determination of fusion protein binding to target antigens of interest can also be performed using any of a number of known methods for identifying and obtaining proteins that specifically interact with other proteins or polypeptides, for example, a yeast two-hybrid screening system such as that described in U.S. Pat. No. 5,283,173 and U.S. Pat. No. 5,468,614, or the equivalent.

Preferred embodiments of the subject invention constructs, for example, binding domain-immunoglobulin fusion proteins, comprise binding regions or binding domains that may include, for example, at least one native or engineered

immunoglobulin variable region polypeptide, such as all or a portion or fragment of a native or engineered heavy chain and/or a native or engineered light chain V-region, provided it is capable of binding or specifically binding an antigen or other desired target structure of interest at a desired level of binding and selectivity. In other preferred embodiments the binding region or binding domain comprises, consists essentially of, or consists of, a single chain immunoglobulin-derived Fv product, for example, and scFv, which may include all or a portion of at least one native or engineered immunoglobulin light chain V-region and all or a portion of at least one native or engineered immunoglobulin heavy chain V-region, and a linker fused or otherwise connected to the V-regions; preparation and testing such constructs are described in greater detail herein. Other preparation and testing methods are well known in the art.

As described herein and known in the art, immunoglobulins comprise products of a gene family the members of which exhibit a high degree of sequence conservation. Amino acid sequences of two or more immunoglobulins or immunoglobulin domains or regions or portions thereof (*e.g.*, V_H domains, V_L domains, hinge regions, CH2 constant regions, CH3 constant regions) may be aligned and analyzed. Portions of sequences that correspond to one another may be identified, for instance, by sequence homology. Determination of sequence homology may be determined with any of a number of sequence alignment and analysis tools, including computer algorithms well known to those of ordinary skill in the art, such as Align or the BLAST algorithm (Altschul, 1991 *J. Mol. Biol.* **219**: 555-565; Henikoff and Henikoff, 1992 *Proc. Natl. Acad. Sci. USA* **89**: 10915-10919), which is available at the National Center for Biotechnology Information (NCBI) website. Default parameters may be used.

Portions, for example, of a particular immunoglobulin reference sequence and of any one or more additional immunoglobulin sequences of interest that may be compared to a reference sequence. "Corresponding" sequences, regions, fragments or the like, may be identified based on the convention for numbering immunoglobulin amino acid positions according to Kabat, *Sequences of Proteins of Immunological Interest*, (5th ed. Bethesda, MD: Public Health Service, National Institutes of Health (1991)). For example, according to this convention, the immunoglobulin family to which an immunoglobulin sequence of interest belongs is

determined based on conservation of variable region polypeptide sequence invariant amino acid residues, to identify a particular numbering system for the immunoglobulin family, and the sequence(s) of interest can then be aligned to assign sequence position numbers to the individual amino acids which comprise such sequence(s). Preferably at least about 70%, more preferably at least about 80%-85% or about 86%-89%, and still more preferably at least about 90%, about 92%, about 94%, about 96%, about 98% or about 99% of the amino acids in a given amino acid sequence of at least about 1000, more preferably about 700-950, more preferably about 350-700, still more preferably about 100-350, still more preferably about 80-100, about 70-80, about 60-70, about 50-60, about 40-50 or about 30-40 consecutive amino acids of a sequence, are identical to the amino acids located at corresponding positions in a reference sequence such as those disclosed by Kabat (1991) or in a similar compendium of related immunoglobulin sequences, such as may be generated from public databases (*e.g.*, Genbank, SwissProt, *etc.*) using sequence alignment tools such as, for example, those described above. In certain preferred embodiments, an immunoglobulin sequence of interest or a region, portion, derivative or fragment thereof is greater than about 95% identical to a corresponding reference sequence, and in certain preferred embodiments such a sequence of interest may differ from a corresponding reference at no more than about 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 amino acid positions.

For example, in certain embodiments the present invention is directed to a construct, including a binding domain-immunoglobulin fusion protein, comprising in pertinent part a human or other species immunoglobulin heavy chain variable region polypeptide comprising a mutation, alteration or deletion at an amino acid at a location or locations corresponding to one or more of amino acid positions 9, 10, 11, 12, 108, 110, 111, and 112 in, for example, SEQ ID NO: 212, which comprises, for example, a murine V_H-derived sequence. At a relatively limited number of immunoglobulin V_H sequence positions, for example, including position 11, amino acid conservation is observed in the overwhelming majority of V_H sequences analyzed across mammalian species lines (*e.g.*, Leu11, Val37, Gly44, Leu45, Trp47; Nguyen *et al.*, 1998 *J. Mol. Biol.* **275**: 413). Various such amino acid residues, and hence their side chains, are located at the surface of the variable domain (V_H). They may contact residues of the C_H1 (*e.g.*, Leu11) and the V_L domains (*e.g.*,

Val37, Gly44, Leu45, and Trp47) and may, in the absence of light chains, contribute to stability and solubility of the protein (*see, e.g., Chothia et al., 1985 J. Mol. Biol.* **186**: 651; Muyldermans *et al.*, 1994 *Prot. Engineer.* **7**: 1129; Desmyter *et al.*, 1996 *Nat. Struct. Biol.* **3**: 803; Davies *et al.*, 1994 *FEBS Lett.* **339**: 285). In certain

5 embodiments, for example, the present invention is also directed to a construct, including a binding domain-immunoglobulin fusion protein, comprising in pertinent part a human immunoglobulin light chain variable region polypeptide, or an immunoglobulin light chain variable region polypeptide from another species, comprising a mutation, alteration or deletion at an amino acid at a location or

10 locations corresponding to one or more of amino acid positions 12, 80, 81, 82, 83, 105, 106, 107 and 108. In still other certain embodiments, for example, the present invention is directed to a construct, including a binding domain-immunoglobulin fusion protein, comprising in pertinent part (1) a human immunoglobulin heavy chain variable region polypeptide, or an immunoglobulin light chain variable region

15 polypeptide from another species, comprising, consisting essentially of, or consisting of, said heavy chain sequence having a mutation, alteration or deletion at a location or locations corresponding to one or more of amino acid positions 9, 10, 11, 12, 108, 110, 111, and 112, and (2) a human immunoglobulin light chain variable region polypeptide, or an immunoglobulin light chain variable region polypeptide from

20 another species, comprising, consisting essentially of, or consisting of, said light chain sequence having a mutation, alteration or deletion at a location or locations corresponding to one or more of amino acid positions 12, 80, 81, 82, 83, 105, 106, 107 and 108.

As another example, by reference to immunoglobulin sequence

25 compendia and databases such as those cited above, for example, the relatedness of two or more immunoglobulin sequences to each other can readily and without undue experimentation be established in a manner that permits identification of the animal species of origin, the class and subclass (*e.g., isotype*) of a particular immunoglobulin or immunoglobulin region polypeptide sequence. Any immunoglobulin variable

30 region polypeptide sequence, including native or engineered V_H and/or V_L and/or single-chain variable region (sFv) sequences or other native or engineered V region-derived sequences or the like, may be used as a binding region or binding domain. Engineered sequences includes immunoglobulin sequences from any species,

preferably human or mouse, for example, that include, for example, a mutation, alteration or deletion at an amino acid at a location or locations corresponding to one or more of amino acid positions 9, 10, 11, 12, 108, 110, 111, and 112 in a heavy chain variable region sequence or an scFv, and/or a mutation, alteration or deletion at a
5 location or locations corresponding to one or more of amino acid positions 12, 80, 81, 82, 83, 105, 106, 107 and 108 in a light chain variable region sequence or an scFv.

Various preferred embodiments include, for example, native or engineered immunoglobulin V region polypeptide sequences derived, for example, from antibodies including monoclonal antibodies such as murine or other rodent
10 antibodies, or antibodies or monoclonal antibodies derived from other sources such as goat, rabbit, equine, bovine, camelid or other species, including transgenic animals, and also including human or humanized antibodies or monoclonal antibodies. Non-limiting examples include variable region polypeptide sequences derived from monoclonal antibodies such as those referenced herein and/or described in greater
15 detail in the Examples below, for instance, CD20-binding or specific murine monoclonal antibodies (*e.g.*, 2H7), monoclonal antibody L6 (specific for a carbohydrate-defined epitope and available from American Type Culture Collection, Manassas, VA, as hybridoma HB8677), and monoclonal antibodies that bind to or are
specific for CD28 (*e.g.*, monoclonal antibody 2E12), CD40, CD80, CD137 (*e.g.*,
20 monoclonal antibody 5B9 or monoclonal antibody 1D8 which recognizes the murine homologue of CD137, 41BB) and CD152 (CTLA-4).

Other binding regions, including binding domain polypeptides, may comprise any protein or portion thereof that retains the ability to bind or specifically bind to an antigen as provided herein, including non-immunoglobulins. Accordingly
25 the invention contemplates constructs, including fusion proteins, comprising binding region or binding domain polypeptides that are derived from polypeptide ligands such as hormones, cytokines, chemokines, and the like; cell surface or soluble receptors for such polypeptide ligands; lectins; intercellular adhesion receptors such as specific leukocyte integrins, selectins, immunoglobulin gene superfamily members,
30 intercellular adhesion molecules (ICAM-1, -2, -3) and the like; histocompatibility antigens; *etc.*

Examples of cell surface receptors useful in the preparation of, or as, binding regions, or that may provide a binding domain polypeptide, and that may also

be selected as a target molecule or antigen to which a construct, including for example, a binding domain-Ig fusion protein of the present invention desirably binds, include the following, or the like: HER1 (*e.g.*, GenBank Accession Nos. U48722, SEG_HEGFREXS, K03193), HER2 (Yoshino *et al.*, 1994 *J. Immunol.* 152: 2393; 5 Disis *et al.*, 1994 *Canc. Res.* **54**: 16; see also, *e.g.*, GenBank Acc. Nos. X03363, M17730, SEG_HUMHER20), HER3 (*e.g.*, GenBank Acc. Nos. U29339, M34309), HER4 (Plowman *et al.*, 1993 *Nature* **366**: 473; see also *e.g.*, GenBank Acc. Nos. L07868, T64105), epidermal growth factor receptor (EGFR) (*e.g.*, GenBank Acc. Nos. U48722, SEG_HEGFREXS, K03193), vascular endothelial cell growth factor 10 (*e.g.*, GenBank No. M32977), vascular endothelial cell growth factor receptor (*e.g.*, GenBank Acc. Nos. AF022375, 1680143, U48801, X62568), insulin-like growth factor-I (*e.g.*, GenBank Acc. Nos. X00173, X56774, X56773, X06043, *see also* European Patent No. GB 2241703), insulin-like growth factor-II (*e.g.*, GenBank Acc. Nos. X03562, X00910, SEG_HUMGFIA, SEG_HUMGFI2, M17863, M17862), 15 transferrin receptor (Trowbridge and Omary, 1981 *Proc. Nat. Acad. USA* **78**: 3039; see also *e.g.*, GenBank Acc. Nos. X01060, M11507), estrogen receptor (*e.g.*, GenBank Acc. Nos. M38651, X03635, X99101, U47678, M12674), progesterone receptor (*e.g.*, GenBank Acc. Nos. X51730, X69068, M15716), follicle stimulating hormone receptor (FSH-R) (*e.g.*, GenBank Acc. Nos. Z34260, M65085), retinoic acid 20 receptor (*e.g.*, GenBank Acc. Nos. L12060, M60909, X77664, X57280, X07282, X06538), MUC-1 (Barnes *et al.*, 1989 *Proc. Nat. Acad. Sci. USA* **86**: 7159; see also *e.g.*, GenBank Acc. Nos. SEG_MUSMUCI0, M65132, M64928) NY-ESO-1 (*e.g.*, GenBank Acc. Nos. AJ003149, U87459), NA 17-A (*e.g.*, European Patent No. WO 96/40039), Melan-A/MART-1 (Kawakami *et al.*, 1994 *Proc. Nat. Acad. Sci. USA* **91**: 3515; see also *e.g.*, GenBank Acc. Nos. U06654, U06452), tyrosinase (Topalian *et al.*, 25 1994 *Proc. Nat. Acad. Sci. USA* **91**: 9461; see also *e.g.*, GenBank Acc. Nos. M26729, SEG_HUMTYRO, *see also* Weber *et al.*, *J. Clin. Invest* (1998) **102**: 1258), Gp-100 (Kawakami *et al.*, 1994 *Proc. Nat. Acad. Sci. USA* **91**: 3515; see also *e.g.*, GenBank Acc. No. S73003, *see also* European Patent No. EP 668350; Adema *et al.*, 1994 *J. Biol. Chem.* **269**: 20126), MAGE (van den Bruggen *et al.*, 1991 *Science* **254**: 1643; 30 see also *e.g.*, GenBank Acc. Nos. U93163, AF064589, U66083, D32077, D32076, D32075, U10694, U10693, U10691, U10690, U10689, U10688, U10687, U10686, U10685, L18877, U10340, U10339, L18920, U03735, M77481), BAGE (*e.g.*, GenBank Acc. No. U19180; see also U.S. Pat. Nos. 5,683,886 and 5,571,711), GAGE

(e.g., GenBank Acc. Nos. AF055475, AF055474, AF055473, U19147, U19146, U19145, U19144, U19143, U19142), any of the CTA class of receptors including in particular HOM-MEL-40 antigen encoded by the SSX2 gene (e.g., GenBank Acc. Nos. X86175, U90842, U90841, X86174), carcinoembryonic antigen (CEA, Gold and
5 Freedman, 1985 *J. Exp. Med.* **121**: 439; *see also e.g.*, GenBank Acc. Nos. SEG_HUMCEA, M59710, M59255, M29540), and PyLT (e.g., GenBank Acc. Nos. J02289, J02038).

Additional cell surface receptors that may be sources of binding region or binding domain polypeptides or portions thereof, or that may be targets, including
10 target antigens, include the following, or the like: CD2 (e.g., GenBank Acc. Nos. Y00023, SEG_HUMCD2, M16336, M16445, SEG_MUSCD2, M14362), 4-1BB (CDw137, Kwon *et al.*, 1989 *Proc. Nat. Acad. Sci. USA* **86**: 1963, 4-1BB ligand (Goodwin *et al.*, 1993 *Eur. J. Immunol.* **23**: 2361; Melero *et al.*, 1998 *Eur. J. Immunol.* **3**: 116), CD5 (e.g., GenBank Acc. Nos. X78985, X89405), CD10 (e.g.,
15 GenBank Acc. Nos. M81591, X76732) CD27 (e.g., GenBank Acc. Nos. M63928, L24495, L08096), CD28 (June *et al.*, 1990 *Immunol. Today* **11**: 211; *see also, e.g.*, GenBank Acc. Nos. J02988, SEG_HUMCD28, M34563), CD152/CTLA-4 (e.g., GenBank Acc. Nos. L15006, X05719, SEG_HUMIGCTL), CD40 (e.g., GenBank Acc. Nos. M83312, SEG_MUSCO40A0, Y10507, X67878, X96710, U15637,
20 L07414), interferon- γ (IFN- γ ; *see, e.g.*, Farrar *et al.* 1993 *Ann. Rev. Immunol.* **11**: 571 and references cited therein, Gray *et al.* 1982 *Nature* **295**: 503, Rinderknecht *et al.* 1984 *J. Biol. Chem.* **259**: 6790, DeGrado *et al.* 1982 *Nature* **300**: 379), interleukin-4 (IL-4; *see, e.g.*, 53rd *Forum in Immunology*, 1993 *Research in Immunol.* **144**: 553-643; Banchereau *et al.*, 1994 in *The Cytokine Handbook*, 2nd ed., A. Thomson, ed.,
25 Academic Press, NY, p. 99; Keegan *et al.*, 1994 *J Leukocyt. Biol.* **55**: 272, and references cited therein), interleukin-17 (IL-17) (e.g., GenBank Acc. Nos. U32659, U43088) and interleukin-17 receptor (IL-17R) (e.g., GenBank Acc. Nos. U31993, U58917). Notwithstanding the foregoing, the present invention expressly does not encompass any immunoglobulin fusion protein that is disclosed in U.S. Pat. No.
30 5,807,734, or U.S. Pat. No. 5,795,572.

Additional cell surface receptors that may be sources of binding region or binding domain polypeptides or portions thereof, or that may serve as targets including target antigens or binding sites include the following, or the like: CD59

(e.g., GenBank Acc. Nos. SEG_HUMCD590, M95708, M34671), CD48 (e.g., GenBank Acc. Nos. M59904), CD58/LFA-3 (e.g., GenBank Acc. No. A25933, Y00636, E12817; see also JP 1997075090-A), CD72 (e.g., GenBank Acc. Nos. AA311036, S40777, L35772), CD70 (e.g., GenBank Acc. Nos. Y13636, S69339),
5 CD80/B7.1 (Freeman *et al.*, 1989 *J. Immunol.* **43**: 2714; Freeman *et al.*, 1991 *J. Exp. Med.* **174**: 625; see also e.g., GenBank Acc. Nos. U33208, 1683379), CD86/B7.2 (Freeman *et al.*, 1993 *J. Exp. Med.* **178**: 2185, Boriello *et al.*, 1995 *J. Immunol.* **155**: 5490; see also, e.g., GenBank Acc. Nos. AF099105, SEG_MMB72G, U39466, U04343, SEG_HSB725, L25606, L25259), B7-H1/B7-DC (e.g., Genbank Acc. Nos.
10 NM_014143, AF177937, AF317088; Dong *et al.*, 2002 *Nat. Med.* June 24 [epub ahead of print], PMID 12091876; Tseng *et al.*, 2001 *J. Exp. Med.* **193**: 839; Tamura *et al.*, 2001 *Blood* **97**: 1809; Dong *et al.*, 1999 *Nat. Med.* **5**: 1365), CD40 ligand (e.g., GenBank Acc. Nos. SEG_HUMCD40L, X67878, X65453, L07414), IL-17 (e.g., GenBank Acc. Nos. U32659, U43088), CD43 (e.g., GenBank Acc. Nos. X52075,
15 J04536), ICOS (e.g., Genbank Acc. No. AH011568), CD3 (e.g., Genbank Acc. Nos. NM_000073 (gamma subunit), NM_000733 (epsilon subunit), X73617 (delta subunit)), CD4 (e.g., Genbank Acc. No. NM-000616), CD25 (e.g., Genbank Acc. No. NM_000417), CD8 (e.g., Genbank Acc. No. M12828), CD11b (e.g., Genbank Acc. No. J03925), CD14 (e.g., Genbank Acc. No. XM 039364), CD56 (e.g., Genbank Acc.
20 No. U63041), CD69 (e.g., Genbank Acc. No. NM_001781) and VLA-4 ($\alpha_4\beta_7$) (e.g., GenBank Acc. Nos. L12002, X16983, L20788, U97031, L24913, M68892, M95632). The following cell surface receptors are typically associated with B cells: CD19 (e.g., GenBank Acc. Nos. SEG_HUMCD19W0, M84371, SEG_MUSCD19W, M62542), CD20 (e.g., GenBank Acc. Nos. SEG_HUMCD20, M62541), CD22 (e.g., GenBank
25 Acc. Nos. 1680629, Y10210, X59350, U62631, X52782, L16928), CD30 (e.g., Genbank Acc. Nos. M83554, D86042), CD153 (CD30 ligand, e.g., GenBank Acc. Nos. L09753, M83554), CD37 (e.g., GenBank Acc. Nos. SEG_MMCD37X, X14046, X53517), CD50 (ICAM-3, e.g., GenBank Acc. No. NM_002162), CD106 (VCAM-1) (e.g., GenBank Acc. Nos. X53051, X67783, SEG_MMVCAM1C, *see also* U.S. Pat.
30 No. 5,596,090), CD54 (ICAM-1) (e.g., GenBank Acc. Nos. X84737, S82847, X06990, J03132, SEG_MUSICAM0), interleukin-12 (see, e.g., Reiter *et al.*, 1993 *Crit. Rev. Immunol.* **13**: 1, and references cited therein), CD134 (OX40, e.g., GenBank Acc. No. AJ277151), CD137 (41BB, e.g., GenBank Acc. No. L12964,

NM_001561), CD83 (e.g., GenBank Acc. Nos. AF001036, AL021918), DEC-205 (e.g., GenBank Acc. Nos. AF011333, U19271).

Constructs, including binding domain-immunoglobulin fusion proteins, of the present invention comprise, for example, a binding domain, such as a binding domain polypeptide that, according to certain particularly preferred embodiments, is capable of binding or specifically binding at least one target, for example, a target antigen or other binding site that is present on an immune effector cell. According to non-limiting theory, such constructs, including for example binding domain-immunoglobulin fusion proteins, may advantageously recruit desired immune effector cell function(s) in a therapeutic context, where it is well known that immune effector cells having different specialized immune functions can be identified or distinguished from one another on the basis of their differential expression of a wide variety of cell surface antigens, including many of the antigens described herein to which constructs of the invention including binding domain polypeptides can specifically bind. As noted herein, immune effector cells include any cell that is capable of directly mediating an activity which is a component of immune system function, including cells having such capability naturally or as a result of genetic engineering.

In certain embodiments an immune effector cell comprises a cell surface receptor for an immunoglobulin or other peptide binding molecule, such as a receptor for an immunoglobulin constant region and including the class of receptors commonly referred to as "Fc receptors" ("FcR"s). A number of FcRs have been structurally and/or functionally characterized and are well known in the art, including FcR having specific abilities to interact with a restricted subset of immunoglobulin heavy chain isotypes, or that interact with Fc domains with varying affinities, and/or which may be expressed on restricted subsets of immune effector cells under certain conditions (e.g., Kijimoto-Ochichai *et al.*, 2002 *Cell Mol. Life Sci.* **59**: 648; Davis *et al.*, 2002 *Curr. Top. Microbiol. Immunol.* **266**: 85; Pawankar, 2001 *Curr. Opin. Allerg. Clin. Immunol.* **1**: 3; Radaev *et al.*, 2002 *Mol. Immunol.* **38**: 1073; Wurzburg *et al.*, 2002 *Mol. Immunol.* **38**: 1063; Sulica *et al.*, 2001 *Int. Rev. Immunol.* **20**: 371; Underhill *et al.*, 2002 *Ann. Rev. Immunol.* **20**: 825; Coggeshall, 2002 *Curr. Dir. Autoimm.* **5**: 1; Mimura *et al.*, 2001 *Adv. Exp. Med. Biol.* **495**: 49; Baumann *et al.*, 2001 *Adv. Exp. Med. Biol.* **495**: 219; Santoso *et al.*, 2001 *Ital. Heart J.* **2**: 811; Novak

et al., 2001 *Curr. Opin. Immunol.* **13**: 721; Fossati *et al.*, 2001 *Eur. J. Clin. Invest.* **31**: 821).

Cells that are capable of mediating ADCC are preferred examples of immune effector cells according to the present invention. Other preferred examples
5 include Natural Killer cells, tumor-infiltrating T lymphocytes (TILs), cytotoxic T lymphocytes, and granulocytic cells such as cells that comprise allergic response mechanisms. Immune effector cells thus include, but are not limited to, cells of hematopoietic origins including cells at various stages of differentiation within myeloid and lymphoid lineages and which may (but need not) express one or more
10 types of functional cell surface FcR, such as T lymphocytes, B lymphocytes, NK cells, monocytes, macrophages, dendritic cells, neutrophils, basophils, eosinophils, mast cells, platelets, erythrocytes, and precursors, progenitors (*e.g.*, hematopoietic stem cells), as well as quiescent, activated, and mature forms of such cells. Other immune effector cells may include cells of non-hematopoietic origin that are capable
15 of mediating immune functions, for example, endothelial cells, keratinocytes, fibroblasts, osteoclasts, epithelial cells, and other cells. Immune effector cells may also include cells that mediate cytotoxic or cytostatic events, or endocytic, phagocytic, or pinocytotic events, or that effect induction of apoptosis, or that effect microbial immunity or neutralization of microbial infection, or cells that mediate allergic,
20 inflammatory, hypersensitivity and/or autoimmune reactions.

Allergic response mechanisms are well known in the art and include an antigen (*e.g.*, allergen)-specific component such as an immunoglobulin (*e.g.*, IgE), as well as the cells and mediators which comprise sequelae to allergen-immunoglobulin (*e.g.*, IgE) encounters (*e.g.*, Ott *et al.*, 2000 *J. Allerg. Clin. Immunol.* **106**: 429;
25 Barnes, 2000 *J. Allerg. Clin. Immunol.* **106**: 5; Togias, 2000 *J. Allerg. Clin. Immunol.* **105**: S599; Akdis *et al.*, 2000 *Int. Arch. Allerg. Immunol.* **121**: 261; Beach, 2000 *Occup. Med.* **15**: 455). Particularly with regard to constructs, including binding domain-immunoglobulin fusion proteins, of the present invention that interact with FcR, certain embodiments of the present invention contemplate constructs including
30 fusion proteins that comprise one or more IgE-derived domains including, for example, those that are capable of inducing an allergic response mechanism that comprises IgE-specific FcR, or portions thereof, which IgE-specific FcRs include those noted above and described or identified in the cited articles. Without wishing to

be bound by particular theory or mechanism, and as disclosed herein, constructs, including fusion proteins, of the present invention may comprise portions of IgE heavy chain Fc domain polypeptides, for example, native or engineered IgE CH3 and CH4 domains, whether provided or expressed as cell surface proteins (*e.g.*, with a plasma membrane anchor domain) or as soluble or otherwise not cell-bound proteins (*e.g.*, without a plasma membrane anchor domain). Further according to non-limiting theory, recruitment and induction of an allergic response mechanism (*e.g.*, an FcR-epsilon expressing immune effector cell) may proceed as the result of either or both of the presence of an IgE Fc domain or portion thereof as described herein (*e.g.*, one that is capable of triggering an allergic mechanism by FcR crosslinking) and the presence of a target such as an antigen to which the binding region, for example a binding domain, binds or specifically binds. The present invention therefore exploits induction of allergic response mechanisms in heretofore unappreciated contexts, such as treatment of a malignant condition or a B cell disorder, including those described or referenced herein.

An immunoglobulin hinge region polypeptide includes any hinge peptide or polypeptide that occurs naturally, as an artificial peptide or as the result of genetic engineering and that is situated, for example, in an immunoglobulin heavy chain polypeptide between the amino acid residues responsible for forming intrachain immunoglobulin-domain disulfide bonds in CH1 and CH2 regions. Hinge region polypeptides for use in the present invention may also include a mutated or otherwise altered hinge region polypeptide. Accordingly, for example, an immunoglobulin hinge region polypeptide may be derived from, or may be a portion or fragment of (*i.e.*, one or more amino acids in peptide linkage, typically about 15-115 amino acids, preferably about 95-110, about 80-94, about 60-80, or about 5-65 amino acids, preferably about 10-50, more preferably about 15-35, still more preferably about 18-32, still more preferably about 20-30, still more preferably about 21, 22, 23, 24, 25, 26, 27, 28 or 29 amino acids) an immunoglobulin polypeptide chain region classically regarded as having hinge function, including those described herein, but a hinge region polypeptide for use in the instant invention need not be so restricted and may include one or more amino acids situated (according to structural criteria for assigning a particular residue to a particular domain that may vary, as known in the art) in an adjoining immunoglobulin domain such as a CH1 domain and/or a CH2 domain in the

cases of IgG, IgA and IgD (or in an adjoining immunoglobulin domain such as a CH1 domain and/or a CH3 domain in the case of IgE), or in the case of certain artificially engineered immunoglobulin constructs, an immunoglobulin variable region domain.

Wild-type immunoglobulin hinge region polypeptides include any
5 known or later-discovered naturally occurring hinge region that is located between the constant region domains, CH1 and CH2, of an immunoglobulin, for example, a human immunoglobulin (or between the CH1 and CH3 regions of certain types of immunoglobulins, such as IgE). For use in constructing one type of connecting region, the wild-type immunoglobulin hinge region polypeptide is preferably a human
10 immunoglobulin hinge region polypeptide, preferably comprising a hinge region from a human IgG, IgA, or IgD immunoglobulin (or the CH2 region of an IgE immunoglobulin), and more preferably, for example, a hinge region polypeptide from a wild-type or mutated human IgG1 isotype as described herein.

As is known to the art, despite the tremendous overall diversity in
15 immunoglobulin amino acid sequences, immunoglobulin primary structure exhibits a high degree of sequence conservation in particular portions of immunoglobulin polypeptide chains, notably with regard to the occurrence of cysteine residues which, by virtue of their sulfhydryl groups, offer the potential for disulfide bond formation with other available sulfhydryl groups. Accordingly, in the context of the present
20 invention wild-type immunoglobulin hinge region polypeptides for use as connecting regions include those that feature one or more highly conserved (*e.g.*, prevalent in a population in a statistically significant manner) cysteine residues, and in certain preferred embodiments a connecting region may comprise, or consist essentially of, or consist of, a mutated hinge region polypeptide may be selected that contains less than
25 the number of naturally-occurring cysteines, for example, zero or one or two cysteine residue(s) in the case of IgG1 and IgG4 hinge regions, and that is derived or constructed from (or using) such a wild-type hinge region sequence.

In certain preferred embodiments wherein the connecting region is a hinge region polypeptide and the hinge region polypeptide is a mutated, engineered or
30 otherwise altered human IgG1 immunoglobulin hinge region polypeptide that is derived or constructed from (or using) a wild-type hinge region sequence, it is noted that the wild-type human IgG1 hinge region polypeptide sequence comprises three non-adjacent cysteine residues, referred to as a first cysteine of the wild-type hinge

region, a second cysteine of the wild-type hinge region and a third cysteine of the wild-type hinge region, respectively, proceeding along the hinge region sequence from the polypeptide N-terminus toward the C-terminus. This can be referred to herein as a "CCC" hinge (or a "WTH", *i.e.*, a wild-type hinge). Examples of mutated or engineered hinge regions include those with no cysteines, which may be referred to herein as an "XXX" hinge (or, for example, as "MH-XXX," referring to a mutant or engineered hinge with three amino acids or other molecules in place of naturally occurring cysteines, such as, for example, "MH-SSS", which refers to a mutant hinge with three serine residues in place of the naturally occurring cysteine residues. It will be understood that the term "mutant" refers only to the fact that a different molecule or molecules is present, or no molecule, at the position of a naturally occurring residue and does not refer to any particular method by which such substitution, alteration, or deletion has been carried out. Accordingly, in certain embodiments of the present invention, the connecting region may be a hinge region polypeptide and the hinge region polypeptide is a mutated human IgG1 immunoglobulin hinge region polypeptide that contains two cysteine residues and in which the first cysteine of the wild-type hinge region has not changed or deleted, for example. This can be referred to as a "MH-CXX" hinge, for example, a "MH-CSC" hinge, in which case the cysteine residue has been replaced with a serine residue. In certain other embodiments of the present invention the mutated human IgG1 immunoglobulin hinge region polypeptide contains no more than one cysteine residue and include, for example, a "MH-CSS" hinge or a "MH-SSC" hinge or a "MH-CSC" hinge, and in certain other embodiments the mutated human IgG1 immunoglobulin hinge region polypeptide contains no cysteine residues such as, for example, a "MH-SSS" hinge.

The constructs, including binding domain-immunoglobulin fusion proteins, of the present invention expressly do not contemplate any fusion protein that is disclosed in U.S. Pat. No. 5,892,019. U.S. Pat. No. 5,892,019 refers to a human IgG1 hinge region in which the first IgG1 hinge region cysteine residue has been changed or deleted, but retains both of the second and third IgG1 hinge region cysteine residues that correspond to the second and third cysteines of the wild-type IgG1 hinge region sequence. The patent states that the first cysteine residue of the wild-type IgG1 hinge region is replaced to prevent interference by the first cysteine residue with proper assembly of the polypeptide described therein into a dimer. The

patent requires that the second and third cysteines of the IgG1 hinge region be retained to provide interchain disulfide linkage between two heavy chain constant regions to promote dimer formation so that the molecule contains has effector function such as the ability to mediate ADCC.

5 By contrast and as described herein, the constructs, including the binding domain-immunoglobulin fusion proteins, of the present invention, various of which are capable of ADCC, CDC and/or complement fixation, for example, are not so limited and may comprise, in pertinent part, for example, (i) a wild-type immunoglobulin hinge region polypeptide, such as a wild-type human
10 immunoglobulin hinge region polypeptide, for example, a human IgG1 immunoglobulin hinge region polypeptide, (ii) a mutated or otherwise altered immunoglobulin hinge region polypeptide, such as a mutated or otherwise altered human immunoglobulin hinge region polypeptide, for example, a mutated or otherwise altered human IgG1 immunoglobulin hinge region polypeptide that, for
15 example, is or has been derived or constructed from (or using) a wild-type immunoglobulin hinge region polypeptide or nucleic acid sequence having three or more cysteine residues, wherein the mutated or otherwise altered human IgG1 immunoglobulin hinge region polypeptide contains two cysteine residues and wherein a first cysteine of the wild-type hinge region is not mutated or deleted, (iii) a mutated
20 or otherwise altered immunoglobulin hinge region polypeptide, such as a mutated or otherwise altered human immunoglobulin hinge region polypeptide, for example, a mutated or otherwise altered human IgG1 immunoglobulin hinge region polypeptide that, for example, is or has been derived or constructed from (or using) a wild-type immunoglobulin hinge region polypeptide or nucleic acid sequence having three or
25 more cysteine residues, wherein the mutated or otherwise altered human IgG1 immunoglobulin hinge region polypeptide contains no more than one cysteine residue, or (iv) a mutated or otherwise altered immunoglobulin hinge region polypeptide, such as a mutated or otherwise altered human immunoglobulin hinge region polypeptide, for example, a mutated or otherwise altered human IgG1
30 immunoglobulin hinge region polypeptide that is or has been derived or constructed from (or using) a wild-type immunoglobulin hinge region polypeptide or nucleic acid sequence having three or more cysteine residues, wherein the mutated or otherwise altered (for example, by amino acid change or deletion) human IgG1 immunoglobulin

hinge region polypeptide contains no cysteine residues. The present invention offers unexpected advantages associated with retention by the constructs, including the fusion proteins, described herein of the ability to mediate ADCC and/or CDC and/or complement fixation notwithstanding that the ability to dimerize via IgG1 hinge
5 region interchain disulfide bonds is ablated or compromised by the removal or replacement of one, two or three hinge region cysteine residues, and even in constructs where the first cysteine of an IgG1 hinge region, for example, is not mutated or otherwise altered or deleted.

A connecting region may comprise a mutated or otherwise altered
10 immunoglobulin hinge region polypeptide, which itself may comprise a hinge region that has its origin in an immunoglobulin of a species, of an immunoglobulin isotype or class, or of an immunoglobulin subclass that is different from that of the tail region, for example, a tail region comprising, or consisting essentially of, or consisting of, CH2 and CH3 domains (or IgE CH3 and CH4 domains). For instance, in certain
15 embodiments of the invention, a construct, for example, a binding domain-immunoglobulin fusion protein, may comprise a binding region such as a binding domain polypeptide that is fused or otherwise connected to an immunoglobulin hinge region polypeptide comprising, or consisting essentially of, or consisting of, a wild-type human IgA hinge region polypeptide, or a mutated or otherwise altered human
20 IgA hinge region polypeptide that contains zero or only one or more cysteine residues (but less than the wild-type number of cysteines), as described herein, or a wild-type human IgG hinge, such as an IgG1 hinge, region polypeptide, or a wild-type human IgE hinge-acting region, *i.e.*, IgE CH2 region polypeptide, or a mutated or otherwise altered human IgG hinge, such as an IgG1 hinge, region polypeptide that is or has
25 been mutated or otherwise altered to contain zero, one or two cysteine residues wherein the first cysteine of the wild-type hinge region is not mutated or altered or deleted, as also described herein. Such a hinge region polypeptide may be fused or otherwise connected to, for example, a tail region comprising, or consisting essentially of, or consisting of, an immunoglobulin heavy chain CH2 region
30 polypeptide from a different Ig isotype or class, for example an IgA or an IgD or an IgG subclass (or a CH3 region from an IgE subclass), which in certain preferred embodiments will be the IgG1 or IgA or IgE subclass and in certain other preferred embodiments may be any one of the IgG2, IgG3 or IgG4 subclasses.

For example, and as described in greater detail herein, in certain embodiments of the present invention a connecting region may be selected to be an immunoglobulin hinge region polypeptide, which is or has been derived from a wild-type human IgA hinge region that naturally comprises three cysteines, where the
5 selected hinge region polypeptide is truncated or otherwise altered or substituted relative to the complete and/or naturally-occurring hinge region such that only one or two of the cysteine residues remain. Similarly, in certain other embodiments of the invention, the construct may be binding domain-immunoglobulin fusion protein comprising a binding domain polypeptide that is fused or otherwise connected to an
10 immunoglobulin hinge region polypeptide comprising a mutated or otherwise altered hinge region polypeptide in which the number of cysteine residues is reduced by amino acid substitution or deletion, for example a mutated or otherwise altered IgG1 hinge region containing zero, one or two cysteine residues as described herein, or an IgD hinge region containing zero cysteine residues.

15 A mutated or otherwise altered hinge region polypeptide may thus be derived or constructed from (or using) a wild-type immunoglobulin hinge region that contains one or more cysteine residues. In certain embodiments, a mutated or otherwise altered hinge region polypeptide may contain zero or only one cysteine residue, wherein the mutated or otherwise altered hinge region polypeptide is or has
20 been derived from a wild type immunoglobulin hinge region that contains, respectively, one or more or two or more cysteine residues. In the mutated or otherwise altered hinge region polypeptide, the cysteine residues of the wild-type immunoglobulin hinge region are preferably deleted or substituted with amino acids that are incapable of forming a disulfide bond. In one embodiment of the invention, a
25 mutated or otherwise altered hinge region polypeptide is or has been derived from a human IgG wild-type hinge region polypeptide, which may include any of the four human IgG isotype subclasses, IgG1, IgG2, IgG3 or IgG4. In certain preferred embodiments, the mutated or otherwise altered hinge region polypeptide is or has been derived from (or using) a human IgA or IgD wild-type hinge region polypeptide.
30 By way of example, a mutated or otherwise altered hinge region polypeptide that is or has been derived from a human IgG1 or IgA wild-type hinge region polypeptide may comprise mutations, alterations, or deletions at two of the three cysteine residues in

the wild-type immunoglobulin hinge region, or mutations, alterations, or deletions at all three cysteine residues.

The cysteine residues that are present in a wild-type immunoglobulin hinge region and that are removed or altered by mutagenesis or any other techniques according to particularly preferred embodiments of the present invention include cysteine residues that form, or that are capable of forming, interchain disulfide bonds. Without wishing to be bound by particular theory or mechanism of action, the present invention contemplates that mutation, deletion, or other alteration of such hinge region cysteine residues, which are believed to be involved in formation of interchain disulfide bridges, reduces the ability of the subject invention binding domain-immunoglobulin fusion protein to dimerize (or form higher oligomers) via interchain disulfide bond formation, while surprisingly not ablating or undesireably compromising the ability of a fusion protein or other construct to promote ADCC, and/or CDC and/or to fix complement. In particular, the Fc receptors that mediate ADCC (*e.g.*, FcRIII, CD16) exhibit low affinity for immunoglobulin Fc domains, supporting the idea that functional binding of Fc to FcR requires avidity stabilization of the Fc-FcR complex by virtue of the dimeric structure of heavy chains in a conventional antibody, and/or FcR aggregation and cross-linking by a conventional antibody Fc structure. Sonderman *et al.*, 2000 *Nature* **406**: 267; Radaev *et al.*, 2001 *J. Biol. Chem.* **276**: 16469; Radaev *et al.*, 2001 *J. Biol. Chem.* **276**: 16478; Koolwijk *et al.*, 1989 *J. Immunol.* **143**: 1656; Kato *et al.*, 2000 *Immunol. Today* **21**: 310. Hence, the constructs, including for example binding domain-immunoglobulin fusion proteins, of the present invention provide the advantages associated with single-chain constructs including single-chain immunoglobulin fusion proteins while also unexpectedly retaining one or more immunological activities. Similarly, the ability to fix complement is typically associated with immunoglobulins that are dimeric with respect to heavy chain constant regions such as those that comprise Fc, while various constructs, including binding domain-immunoglobulin fusion proteins, of the present invention, which may, due to the replacement or deletion of hinge region cysteine residues or due to other structural modifications as described herein, for example, have compromised or ablated abilities to form interchain disulfide bonds, exhibit the unexpected ability to fix complement. Additionally, according to certain embodiments of the present invention wherein a construct, including, for example, a binding

domain-immunoglobulin fusion protein, may comprise a connecting region and tail region comprising, or consisting essentially of, or consisting of, one or more of a human IgE hinge-acting region, *i.e.*, a IgE CH2 region polypeptide, a human IgE CH3 constant region polypeptide, and a human IgE CH4 constant region polypeptide, the
5 invention constructs including fusion proteins unexpectedly retain the immunological activity of mediating ADCC and/or of inducing an allergic response mechanism.

Selection of an immunoglobulin hinge region polypeptide as a connecting region according to certain embodiments of the subject invention constructs, such as binding domain-immunoglobulin fusion proteins, may relate to the
10 use of an "alternative hinge region" polypeptide sequence, which includes a polypeptide sequence that is not necessarily derived from any immunoglobulin hinge region sequence *per se*. Instead, an alternative hinge region refers to a hinge region polypeptide that comprises an amino acid sequence, or other molecular sequence, of at least about ten consecutive amino acids or molecules, and in certain embodiments
15 at least about 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-25, 26-30, 31-50, 51-60, 71-80, 81-90, or 91-110 amino acids or molecules that is present in a sequence disclosed herein, for example a polypeptide sequence that is or has been derived from a region located between intrachain disulfide-generated immunoglobulin-like loop domains of immunoglobulin gene superfamily members such as CD2 (*e.g.*, Genbank Acc. No.
20 NM_001767), CD4 (*e.g.*, Genbank Acc. No. NM_000616), CD5 (*e.g.*, Genbank Acc. No. BC027901), CD6 (*e.g.*, Genbank Acc. No. NM-006725), CD7 (*e.g.*, Genbank Acc. Nos. XM_046782, BC009293, NM_006137) or CD8 (*e.g.*, Genbank Acc. No. M12828), or other Ig superfamily members. By way of non-limiting example, an alternative hinge region used as a connecting region, for example, may provide a
25 glycosylation site as provided herein, or may provide a human gene-derived polypeptide sequence for purposes of enhancing the degree of "humanization" of a fusion protein, or may comprise, or consist essentially of, or consist of, an amino acid sequence that eliminates or reduces the ability of a construct of the invention, such as a fusion protein, to form multimers or oligomers or aggregates or the like. Certain
30 alternative hinge region polypeptide sequences, including those described herein, may be derived or constructed from (or using) the polypeptide sequences of immunoglobulin gene superfamily members that are not actual immunoglobulins *per se*. For instance and according to non-limiting theory, certain polypeptide sequences

that are situated between intrachain disulfide-generated immunoglobulin loop domain of immunoglobulin gene super-family member proteins may be used in whole or in part as alternative hinge region polypeptides as provided herein, or may be further modified for such use.

5 As noted above, the constructs of the invention, including binding domain-immunoglobulin fusion proteins, are believed, according to non-limiting theory, to be compromised in their ability to dimerize via interchain disulfide bond formation, and further according to theory, this property is a consequence, in whole or in part, of a reduction in the number of cysteine residues that are present in an
10 immunoglobulin hinge region polypeptide selected for inclusion in the construction of the construct, such as a fusion protein construct. Determination of the relative ability of a polypeptide to dimerize is well within the knowledge of the relevant art, where any of a number of established methodologies may be applied to detect protein dimerization (see, *e.g.*, Scopes, *Protein Purification: Principles and Practice*, 1987
15 Springer-Verlag, New York). For example, biochemical separation techniques for resolving proteins on the basis of molecular size (*e.g.*, gel electrophoresis, gel filtration chromatography, analytical ultracentrifugation, *etc.*), and/or comparison of protein physicochemical properties before and after introduction of sulfhydryl-active (*e.g.*, iodoacetamide, N-ethylmaleimide) or disulfide-reducing (*e.g.*, 2-
20 mercaptoethanol, dithiothreitol) agents, or other equivalent methodologies, may all be employed for determining a degree of polypeptide dimerization or oligomerization, and for determining possible contribution of disulfide bonds to such potential quarternary structure. In certain embodiments, the invention relates to a construct, for example a binding domain-immunoglobulin fusion protein, that exhibits a reduced
25 (*i.e.*, in a statistically significant manner relative to an appropriate IgG-derived control, for example) ability to dimerize, relative to a wild-type human immunoglobulin G hinge region polypeptide as provided herein. Those familiar with the art will be able readily to determine whether a particular fusion protein displays such reduced ability to dimerize.

30 Compositions and methods for preparation of immunoglobulin fusion proteins, for example, are well known in the art. See, *e.g.*, U.S. Pat. No. 5,892,019, which reports recombinant proteins that are the products of a single encoding

polynucleotide but which are not constructs, including binding domain-immunoglobulin fusion proteins, according to the present invention.

For a construct, for example, an immunoglobulin fusion protein of the invention which is intended for use in humans, any included Ig constant regions will typically be of human sequence origin, or humanized, to minimize a potential anti-human immune response and to provide appropriate and/or desired effector functions. Manipulation of sequences encoding antibody constant regions is referenced in the PCT publication of Morrison and Oi, WO 89/07142. In particularly preferred embodiments, a tail region is prepared from an immunoglobulin heavy chain constant region in which the CH1 domain is or has been deleted (the CH1 and CH2 regions in the case of IgE) and the carboxyl end of the binding domain, or where the binding domain comprises two immunoglobulin variable region polypeptides, the second (*i.e.*, more proximal to the C-terminus) variable region is joined to the amino terminus of CH2 through one or more connecting regions, such as a hinge or altered region. A schematic diagram depicting the structures of two exemplary binding domain-immunoglobulin fusion proteins is shown in Figure 11. In particularly preferred embodiments no interchain disulfide bonds are present, and in other embodiments a restricted number of interchain disulfide bonds may be present relative to the number of such bonds that would be present if wild-type hinge region polypeptides were instead present. In other embodiments a construct of the invention, such as for example, a fusion protein, comprises, or consists essentially of, or consists of, a mutated or otherwise altered hinge region polypeptide that exhibits a reduced ability to dimerize, relative to a wild-type human IgG hinge region polypeptide. Thus, an isolated polynucleotide molecule coding for such a single chain construct, such as an immunoglobulin fusion protein, has a binding region, for example, a domain that provides specific or otherwise desired binding affinity and selectivity for a target, such as a target antigen.

The invention also contemplates, for example, in certain embodiments, constructs including binding domain-immunoglobulin fusion proteins that comprise fused or otherwise connected polypeptide sequences or portions thereof derived or prepared from a plurality of genetic sources, for example, according to molecular "domain swapping" paradigms. Those having familiarity with the art will appreciate that selection of such polypeptide sequences for assembly into a construct, such as a

binding domain-immunoglobulin fusion protein, for example, may involve determination of appropriate portions of each component polypeptide sequence, for example, based on structural and/or functional properties of each such sequence (see, *e.g.*, Carayannopoulos *et al.*, 1996 *J. Exp. Med* **183**: 1579; Harlow *et al.*, Eds.,
5 *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor (1988)). The component polypeptide sequences of which the construct, such as a fusion protein, is comprised or prepared may therefore comprise intact or full-length binding domain, immunoglobulin, linker and/or plasma membrane anchor domain polypeptide sequences, or truncated versions or variants thereof such as those
10 provided herein. According to these and related embodiments of the invention, any two or more of the candidate component polypeptides of which the subject invention constructs, for example, fusion proteins, may be comprised will be derived or prepared from independent sources, such as from immunoglobulin sequences of differing allotype, isotype, subclass, class, or species of origin (*e.g.*, xenotype). Thus,
15 as a non-limiting example, a binding domain polypeptide (or its constituent polypeptides such as one or more variable region polypeptides and/or a linker polypeptide), a hinge region polypeptide, immunoglobulin heavy chain CH2 and CH3 constant region polypeptides and optionally an immunoglobulin heavy chain CH4 constant region polypeptide as may be obtained from an IgM or IgE heavy chain, and
20 a plasma membrane anchor domain polypeptide may all be separately obtained from distinct genetic sources and engineered into a chimeric or fusion protein using well known techniques and according to methodologies described herein, for example.

Accordingly, a construct of the invention, for example a binding domain-immunoglobulin fusion protein according to certain embodiments of the
25 present invention, may also therefore comprise in pertinent part an immunoglobulin heavy chain CH3 constant region polypeptide that is a wild-type IgA CH3 constant region polypeptide, or alternatively, that is a mutated or otherwise altered or substituted or truncated IgA CH3 constant region polypeptide that is incapable of associating with a J chain, or that will not associate to an undesired degree with a J
30 chain; preferably the IgA CH3 constant region polypeptides used in a tail region portion of a construct are of human origin or are humanized. By way of brief background, IgA molecules are known to be released into secretory fluids by a mechanism that involves association of IgA into disulfide-linked polymers (*e.g.*,

dimers) via a J chain polypeptide (*e.g.*, Genbank Acc. Nos. XM-059628, M12378, M12759; Johansen *et al.*, 1999 *Eur. J. Immunol.* **29**: 1701) and interaction of the complex so formed with another protein that acts as a receptor for polymeric immunoglobulin, and which is known as transmembrane secretory component (SC; Johansen *et al.*, 2000 *Sc. J. Immunol.* **52**: 240; see also, *e.g.*, Sorensen *et al.*, 2000 *Int. Immunol.* **12**: 19; Yoo *et al.*, 1999 *J. Biol. Chem.* **274**: 33771; Yoo *et al.*, 2002 *J. Immunol. Meth.* **261**: 1; Cortesy, 2002 *Trends Biotechnol.* **20**: 65; Symersky *et al.*, 2000 *Mol. Immunol.* **37**: 133; Crottet *et al.*, 1999 *Biochem. J.* **341**: 299). Interchain disulfide bond formation between IgA Fc domains and J chain is mediated through a penultimate cysteine residue in an 18-amino acid C-terminal extension that forms part of the IgA heavy chain constant region CH3 domain polypeptide (Yoo *et al.*, 1999; Sorensen *et al.*, 2000). Certain embodiments of the subject invention constructs, including for example, fusion proteins, therefore contemplate inclusion of the wild-type IgA heavy chain constant region polypeptide sequence, which is capable of associating with J chain. Certain other embodiments of the invention, however, contemplate fusion proteins that comprise a mutated or otherwise altered, substituted, or truncated IgA CH3 constant region polypeptide that is incapable of associating with a J chain. According to such embodiments, for example, two or more residues from the C-terminus of an IgA CH3 constant region polypeptide such as a human IgA CH3 constant region polypeptide may be deleted to yield a truncated CH3 constant region polypeptide as provided herein. In preferred embodiments and as described in greater detail herein, a mutated human IgA CH3 constant region polypeptide that is incapable of associating with a J chain comprises such a C-terminal deletion of either four or 18 amino acids. However, the invention need not be so limited, such that the mutated IgA CH3 constant region polypeptide may comprise a deletion of about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21-25, 26-30 or more amino acids, so long as the construct, for example, the fusion protein, is capable of specifically binding an antigen and of at least one immunological activity as provided herein. Alternatively, the invention also contemplates constructs, for example, fusion proteins, having a tail region that comprises a mutated IgA CH3 constant region polypeptide that is incapable of associating with a J chain by virtue of replacement of the penultimate cysteine, or by chemical modification of that amino acid residue, in a manner that prevents, or inhibits an undesired level of, interchain disulfide bond or multimer formation. Methods for determining whether a construct, for example a

fusion protein, can associate with a J chain will be known to those having familiarity with the art and are described or referenced herein.

As also described herein and according to procedures known in the art, the construct, for example a fusion protein, may further be tested routinely for immunological activity, for instance, in ADCC or CDC assays. As an illustrative example, a construct, for example a fusion protein, according to such an embodiment may comprise a binding domain polypeptide derived or constructed from (or using) a native or engineered human heavy chain variable region polypeptide sequence, a native or engineered human IgA-derived immunoglobulin hinge region polypeptide sequence, a native or engineered human IgG1 immunoglobulin heavy chain CH2 constant region polypeptide sequence, a native or engineered human IgG2 immunoglobulin heavy chain CH3 constant region polypeptide sequence, and optionally a native or engineered human IgE immunoglobulin heavy chain CH4 constant region polypeptide sequence and/or a native or engineered human TNF- α receptor type 1 (TNFR1) plasma membrane anchor domain polypeptide sequence that comprises a cytoplasmic tail polypeptide which is capable of apoptotic signaling or otherwise promoting apoptosis. The invention therefore contemplates these and other embodiments according to the present invention in which two or more polypeptide sequences that are present in a construct, for example a fusion protein, have independent genetic origins..

As noted above, in certain embodiments the construct, for example a binding protein-immunoglobulin fusion protein, comprises at least one native or engineered immunoglobulin variable region polypeptide, which may be a native or engineered light chain or a native or engineered heavy chain variable region polypeptide, and in certain embodiments the fusion protein comprises at least one such native or engineered light chain V-region and one such native or engineered heavy chain V-region and at least one linker peptide that is fused or otherwise connected to to each of the native or engineered V-regions. Construction of such binding domains, for example single chain Fv domains, is known in the art and is described in greater detail in the Examples below, and has been described, for example, in various documents cited herein; selection and assembly of single-chain variable regions and of linker polypeptides that may be fused or otherwise connected to each of a heavy chain-derived and a light chain-derived V region (*e.g.*, to generate

a binding region, such as a binding domain that comprises a single-chain Fv polypeptide) is also known to the art and described herein. See, e.g., U.S. Pat. Nos. 5,869,620, 4,704,692, and 4,946,778. In certain embodiments all or a portion or portions of an immunoglobulin sequence that is derived from a non-human source may be "humanized" according to recognized procedures for generating humanized antibodies, *i.e.*, immunoglobulin sequences into which human Ig sequences are introduced to reduce the degree to which a human immune system would perceive such proteins as foreign (see, *e.g.*, U.S. Pat. Nos. 5,693,762; 5,585,089; 4,816,567; 5,225,539; 5,530,101; and documents cited therein).

Constructs of the invention, including binding domain-immunoglobulin fusion proteins, as described herein may, according to certain embodiments, desirably comprise sites for glycosylation, *e.g.*, covalent attachment of carbohydrate moieties such as, for example, monosaccharides or oligosaccharides. Incorporation of amino acid sequences that provide substrates for polypeptide glycosylation is within the scope of the relevant art, including, for example, the use of genetic engineering or protein engineering methodologies to obtain a polypeptide sequence containing, for example, the classic Asn-X-Ser/Thr site for N-(asparagine)-linked glycosylation, or a sequence containing Ser or Thr residues that are suitable substrates for O-linked glycosylation, or sequences amenable to C-mannosylation, glypiation/glycosylphosphatidylinositol modification, or phosphoglycation, all of which can be identified according to art-established criteria (*e.g.*, Spiro, 2002 *Glybiology* 12: 43R). Without wishing to be bound by any particular theory or mechanism, glycosylated constructs such as fusion proteins having particular amino acid sequences may beneficially possess attributes associated with one or more of improved solubility, enhanced stability in solution, enhanced physiological stability, improved bioavailability including in vivo biodistribution, and superior resistance to proteases, all in a statistically significant manner, relative to constructs, including fusion proteins, having the same or highly similar amino acid sequences but lacking glycosyl moieties. In certain preferred embodiments the subject invention constructs, such as fusion protein constructs, may comprise a glycosylation site that is present in a linker as provided herein, and in certain other preferred embodiments the subject invention construct, for example, a fusion protein, comprises a glycosylation site that

is present in a connecting region, such as a hinge region polypeptide sequence as provided herein.

In certain preferred embodiments of the present invention, such as those useful for gene therapy applications or in display systems or assays, such as screening assays (including library display systems and library screening assays), the construct, for example, a binding domain-immunoglobulin fusion protein, is a protein or glycoprotein that is capable of being expressed by a host cell such that it localizes to the cell surface. Constructs, such as binding domain-immunoglobulin fusion proteins, that localize to the cell surface may do so by virtue of having naturally present or artificially introduced structural features that direct the fusion protein to the cell surface (*e.g.*, Nelson *et al.* 2001 *Trends Cell Biol.* **11**: 483; Ammon *et al.*, 2002 *Arch. Physiol. Biochem.* **110**: 137; Kasai *et al.*, 2001 *J. Cell Sci.* **114**: 3115; Watson *et al.*, 2001 *Am. J. Physiol. Cell Physiol.* **281**: C215; Chatterjee *et al.*, 200 *J. Biol. Chem.* **275**: 24013) including by way of illustration and not limitation, secretory signal sequences, leader sequences, plasma membrane anchor domain polypeptides and transmembrane domains such as hydrophobic transmembrane domains (*e.g.*, Heuck *et al.*, 2002 *Cell Biochem. Biophys.* **36**: 89; Sadlish *et al.*, 2002 *Biochem J.* **364**: 777; Phoenix *et al.*, 2002 *Mol. Membr. Biol.* **19**: 1; Minke *et al.*, 2002 *Physiol. Rev.* **82**: 429) or glycosylphosphatidylinositol attachment sites ("glypiation" sites, *e.g.*, Chatterjee *et al.*, 2001 *Cell Mol. Life Sci.* **58**: 1969; Hooper, 2001 *Proteomics* **1**: 748; Spiro, 2002 *Glycobiol.* **12**: 43R), cell surface receptor binding domains, extracellular matrix binding domains, or any other structural feature that causes at least a desired portion of the fusion protein population to localize, in whole or in part, to the cell surface. Particularly preferred are fusion protein constructs that comprise a plasma membrane anchor domain which includes a transmembrane polypeptide domain, typically comprising a membrane spanning domain which includes a hydrophobic region capable of energetically favorable interaction with the phospholipid fatty acyl tails that form the interior of the plasma membrane bilayer. Such features are known to those of ordinary skill in the art, who will further be familiar with methods for introducing nucleic acid sequences encoding these features into the subject expression constructs by genetic engineering, and with routine testing of such constructs to verify cell surface localization of the product.

According to certain further embodiments, a plasma membrane anchor domain polypeptide comprises such a transmembrane domain polypeptide and also comprises a cytoplasmic tail polypeptide, which refers to a region or portion of the polypeptide sequence that contacts the cytoplasmic face of the plasma membrane and/or is in contact with the cytosol or other cytoplasmic components. A large number of cytoplasmic tail polypeptides are known that comprise the intracellular portions of plasma membrane transmembrane proteins, and discrete functions have been identified for many such polypeptides, including biological signal transduction (*e.g.*, activation or inhibition of protein kinases, protein phosphatases, G-proteins, cyclic nucleotides and other second messengers, ion channels, secretory pathways), biologically active mediator release, stable or dynamic association with one or more cytoskeletal components, cellular differentiation, cellular activation, mitogenesis, cytostasis, apoptosis and the like (*e.g.*, Maher *et al.*, 2002 *Immunol. Cell Biol.* **80**: 131; El Far *et al.*, 2002 *Biochem J* **365**: 329; Teng *et al.*, 2002 *Genome Biol.* **2**REVIEWS: 3012; Simons *et al.*, 2001 *Cell Signal* **13**: 855; Furie *et al.*, 2001 *Thromb. Haemost.* **86**: 214; Gaffen, 2001 *Cytokine* **14**: 63; Dittel, 2000 *Arch. Immunol. Ther. Exp. (Warsz.)* **48**: 381; Parnes *et al.*, 2000 *Immunol. Rev.* **176**: 75; Moretta *et al.*, 2000 *Semin. Immunol.* **12**: 129; Ben Ze'ev, 1999 *Ann. NY Acad. Sci.* **886**: 37; Marsters *et al.*, *Recent Prog Horm. Res.* **54**: 225).

Figure 70 illustrates the binding, for example, of fluorocine-conjugated FcRIII (CD16) soluble fusion proteins to 2H7 scFv-binding domain constructs that are attached to CD20 expressed by cells, CHO cells in this example. CD16 binding to a construct of the invention, for example, scFv-binding domain construct, provides one example of a screening tool that may be used to detect and/or quantitate changes in CD16 binding to altered constructs of the invention, including scFv-binding domain constructs, that contains targeted or site-specific mutations, substitutions, deletions, or other alterations. Changes in CD16 binding properties may be reflected, for example, by changes in binding of either CD16 high affinity protein (158 V) or CD16 low affinity protein (158 F) or both.

A schematic representation of one example of such a screening process is diagrammed in the second drawing in Figure 70, in which scFv-binding domain constructs are displayed on the cell surface of mammalian cells scFv-binding domain molecules in this example are displayed on the cell surface through a molecule that

serves as a transmembrane domain anchor. These molecules may represent, for example, a single scFv-binding domain construct or may be introduced into a population of mammalian cells as a library of such molecules. Transfected cells with altered binding properties can then, for example, be panned, sorted, or otherwise
5 isolated from other cells by altering the stringency of the selection conditions and using CD16 fusion proteins as a binding probe. Cells that express scFv-Ig molecules with altered binding to either CD 16 high affinity allele (158 V) or CD16 low affinity allele (158F) or both, for example, can be isolated.

This display system can be used, for example, to create a library of
10 constructs of the invention with mutated or otherwise altered tail regions with short stretches of CH2 sequence replaced with randomized oligonucleotides or, for example, randomization of a single residue with all possible amino acid substitutions, natural or unnatural, including synthetic amino acids. Once such a library is constructed, it can be transfected into appropriate cells, for example, COS cells, by
15 methods known in the art. Transfectants can then be bound to, for example, labeled CD16 constructs, and panned or sorted based on their relative or desired binding properties to multiple allelotypes/isoforms. Desired cells may be harvested, and the DNA, for example, plasmid DNA, isolated and then transformed into, for example, bacteria. This process may be repeated iteratively multiple times until desired single
20 clones are isolated from the mammalian host cells. See Seed B and Aruffo A, *Pro. Nat'l Acad Sci USA* 1987 **84**: 3365-3369; Aruffo A and Seed B, *Pro. Nat'l Acad Sci USA* 1987 **84**: 8573-8577.

One such use of this type of screening system, for example, is for the identification and/or isolation of constructs of the invention having tail regions, or tail
25 regions, that bind equally well to both the high and low affinity alleles of CD16 with the goal of improving effector functions mediated by scFv-binding domain constructs in multiple subpopulations of patients. Constructs of the invention having tail regions, or tail regions with altered binding properties to other Fc receptors can also be selected using such a display system, for example, the display system described.
30 Other display systems that do not glycosylate proteins, for example, those that use bacteriophage or yeast, are not generally desired for selection of constructs of the invention having Ig-based tail regions, or Ig-based tail regions, with altered FcR

binding properties. Most non-mammalian systems, for example, do not glycosylate proteins.

Expression of constructs of the invention, for example, scFv-binding domain constructs, expressed on the surface of a mammalian cell by incorporation of an appropriate molecule into the construct, for example, by incorporation of a transmembrane domain or a GPI anchor signal, also have utility in other display systems that are useful, for example, for selection of constructs of the invention, for example, altered scFv-binding domain molecules that will be produced at higher or other desired levels. In one such an embodiment, cells that are useful in the production of glycosylated proteins, for example, mammalian cells such as COS cells, are transfected with a library of scFv-binding domain constructs in a plasmid that directs their expression to the cell surface. Cells, such as COS cells, that express the highest or other desired level of the scFv-binding domain molecules are selected by techniques known in the art (for example panning, sterile cell sorting, magnetic bead separation, *etc.*), and DNA, for example, plasmid DNA, is isolated for transformation into other cells, for example, bacteria. After one or more rounds of selection single clones are isolated that encode scFv-binding domain molecules capable of a high or other desired level of expression. The isolated clones may then be altered to remove the membrane anchor and expressed in an appropriate cells system, for example, a mammalian cell system, wherein the scFv-binding domain constructs will be produced, for example, by secretion into the culture fluid at desired levels. Without being bound by any particular mechanism or theory, this is believed to result from the common requirement of secreted glycoproteins and cell surface glycoproteins for a signal peptide and processing through the golgi for expression. Thus, selection for a molecule that shows an improvement expression levels on a cell surface will also result in the identification of a molecule having an improvement in levels of secreted protein.

These display systems utilizing a construct of the invention may also be used for screening and/or identifying and/or isolating affinity variants of the binding domain within a construct.

Particularly preferred are such display and/or screening systems, for example, that include or use constructs that include (1) an immunoglobulin variable region polypeptide sequence, including native or engineered V_H and/or V_L and/or

single-chain variable region (sFv) sequences, and which include, for example, a mutation, alteration or deletion at an amino acid at a location or locations corresponding to one or more of amino acid positions 9, 10, 11, 12, 108, 110, 111, and 112, in a V_H region sequence (including in a V_H region sequence within an scFv or other construct), and/or (2) an immunoglobulin variable region polypeptide sequence, including native or engineered V_H and/or V_L and/or single-chain variable region (sFv) sequences, and which include, for example, a a mutation, alteration or deletion at a location or locations corresponding to one or more of amino acid positions 12, 80, 81, 82, 83, 105, 106, 107 and 108 in a light chain variable region sequence (including in a V_L region sequence within an scFv or other construct). Especially preferred are such display and/or screening systems that include or use constructs that include an engineered V_H sequence (whether or not associated with one or more other sequences, including immunoglobulin-derived and other sequences contained, for example, within an sFv or scFv-containing construct), which includes a mutation, alteration or deletion at an amino acid at a location or locations corresponding to amino acid position 11. The V_H 11 amino acid, if substituted, may be substituted with another amino acid as described herein, or by another molecule as desired.

In the context of other methods of using constructs of the invention, including binding domain-immunoglobulin fusion proteins, for the treatment of a malignant condition or a B cell disorder(s) as provided herein, including, for example, by one or more of a number of gene therapy methods and related construct delivery techniques, the present invention also contemplates certain embodiments wherein a construct, for example, a binding domain-immunoglobulin fusion protein that comprises a plasma membrane anchor domain polypeptide is expressed (or capable of expression) at a cell surface and may further comprise a cytoplasmic tail polypeptide which comprises an apoptosis signaling polypeptide sequence. A number of apoptosis signaling polypeptide sequences are known to the art, as reviewed, for example, in *When Cells Die: A Comprehensive Evaluation of Apoptosis and Programmed Cell Death* (B.A. Lockshin *et al.*, Eds., 1998 John Wiley & Sons, New York; see also, *e.g.*, Green *et al.*, 1998 *Science* **281**: 1309 and references cited therein; Ferreira *et al.*, 2002 *Clin. Canc. Res.* **8**: 2024; Gurumurthy *et al.*, 2001 *Cancer Metastas. Rev.* **20**: 225; Kanduc *et al.*, 2002 *Int. J. Oncol.* **21**: 165). Typically an apoptosis signaling

polypeptide sequence comprises all or a portion of, or is derived or constructed from, a receptor death domain polypeptide, for instance, FADD (*e.g.*, Genbank Acc. Nos. U24231, U43184, AF009616, AF009617, NM_012115), TRADD (*e.g.*, Genbank Acc. No. NM_003789), RAIDD (*e.g.*, Genbank Acc. No. U87229), CD95 (FAS/Apo-1; *e.g.*, Genbank Acc. Nos. X89101, NM-003824, AF344850, AF344856), TNF- α -receptor-1 (TNFR1, *e.g.*, Genbank Acc. Nos. S63368, AF040257), DR5 (*e.g.*, Genbank Acc. No. AF020501, AF016268, AF012535), an ITIM domain (*e.g.*, Genbank Acc. Nos. AF081675, BC015731, NM_006840, NM_006844, NM_006847, XM_017977; see, *e.g.*, Billadeau *et al.*, 2002 *J. Clin. Invest.* **109**: 161), an ITAM domain (*e.g.*, Genbank Acc. Nos. NM_005843, NM_003473, BC030586; see, *e.g.*, Billadeau *et al.*, 2002), or other apoptosis-associated receptor death domain polypeptides known to the art, for example, TNFR2 (*e.g.*, Genbank Acc. No. L49431, L49432), caspase/procaspase-3 (*e.g.*, Genbank Acc. No. XM_54686), caspase/procaspase-8 (*e.g.*, AF380342, NM_004208, NM_001228, NM_033355, NM_033356, NM_033357, NM_033358), caspase/procaspase-2 (*e.g.*, Genbank Acc. No. AF314174, AF314175), *etc.*

Cells in a biological sample that are suspected of undergoing apoptosis may be examined for morphological, permeability or other changes that are indicative of an apoptotic state. For example by way of illustration and not limitation, apoptosis in many cell types may cause altered morphological appearance such as plasma membrane blebbing, cell shape change, loss of substrate adhesion properties or other morphological changes that can be readily detected by a person having ordinary skill in the art, for example by using light microscopy. As another example, cells undergoing apoptosis may exhibit fragmentation and disintegration of chromosomes, which may be apparent by microscopy and/or through the use of DNA-specific or chromatin-specific dyes that are known in the art, including fluorescent dyes. Such cells may also exhibit altered plasma membrane permeability properties as may be readily detected through the use of vital dyes (*e.g.*, propidium iodide, trypan blue) or by the detection of lactate dehydrogenase leakage into the extracellular milieu. These and other means for detecting apoptotic cells by morphologic criteria, altered plasma membrane permeability, and related changes will be apparent to those familiar with the art.

In another embodiment of the invention wherein a construct, such as a binding domain-immunoglobulin fusion protein, that is expressed at a cell surface comprises a plasma membrane anchor domain having a transmembrane domain and a cytoplasmic tail that comprises an apoptosis signaling polypeptide, cells in a biological sample may be assayed for translocation of cell membrane phosphatidylserine (PS) from the inner to the outer leaflet of the plasma membrane, which may be detected, for example, by measuring outer leaflet binding by the PS-specific protein annexin. Martin *et al.*, *J. Exp. Med.* **182**: 1545, 1995; Fadok *et al.*, *J. Immunol.* **148**: 2207, 1992. In still other related embodiments of the invention, including embodiments wherein a construct, such as a binding domain-immunoglobulin fusion protein, is expressed at the cell surface and comprises a plasma membrane anchor domain having an apoptosis signaling polypeptide and also including embodiments wherein the construct, such as a binding domain-immunoglobulin fusion protein, is a soluble protein that lacks a membrane anchor domain and that is capable of inducing apoptosis, a cellular response to an apoptogen is determined by an assay for induction of specific protease activity in any member of a family of apoptosis-activated proteases known as the caspases (see, *e.g.*, Green *et al.*, 1998 *Science* **281**: 1309). Those having ordinary skill in the art will be readily familiar with methods for determining caspase activity, for example by determination of caspase-mediated cleavage of specifically recognized protein substrates. These substrates may include, for example, poly-(ADP-ribose) polymerase (PARP) or other naturally occurring or synthetic peptides and proteins cleaved by caspases that are known in the art (see, *e.g.*, Ellerby *et al.*, 1997 *J. Neurosci.* **17**: 6165). The synthetic peptide Z-Tyr-Val-Ala-Asp-AFC (SEQ ID NO: 528), wherein "Z" indicates a benzoyl carbonyl moiety and AFC indicates 7-amino-4-trifluoromethylcoumarin (Kluck *et al.*, 1997 *Science* **275**: 1132; Nicholson *et al.*, 1995 *Nature* **376**: 37), is one such substrate. Other non-limiting examples of substrates include nuclear proteins such as U1-70 kDa and DNA-PKcs (Rosen and Casciola-Rosen, 1997 *J. Cell. Biochem.* **64**: 50; Cohen, 1997 *Biochem. J.* **326**: 1). Cellular apoptosis may also be detected by determination of cytochrome c that has escaped from mitochondria in apoptotic cells (*e.g.*, Liu *et al.*, *Cell* **86**: 147, 1996). Such detection of cytochrome c may be performed spectrophotometrically, immunochemically or by other well established methods for determining the presence of a specific protein. Persons having ordinary skill in the art

will readily appreciate that there may be other suitable techniques for quantifying apoptosis.

Particularly preferred embodiments of constructs useful for gene therapy applications, including those constructs that include a plasma membrane anchor domain and/or cytoplasmic tail polypeptide (including, for example, an apoptosis signaling sequence), are such constructs that include (1) an immunoglobulin variable region polypeptide sequence, including native or engineered V_H and/or V_L and/or single-chain variable region (sFv) sequences, and which include, for example, a mutation, alteration or deletion at an amino acid at a location or locations corresponding to one or more of amino acid positions 9, 10, 11, 12, 108, 110, 111, and 112, in a V_H region sequence (including in a V_H region sequence within an scFv or other construct), and/or (2) an immunoglobulin variable region polypeptide sequence, including native or engineered V_H and/or V_L and/or single-chain variable region (sFv) sequences, and which include, for example, a a mutation, alteration or deletion at a location or locations corresponding to one or more of amino acid positions 12, 80, 81, 82, 83, 105, 106, 107 and 108 in a light chain variable region sequence (including in a V_L region sequence within an scFv or other construct). Especially preferred are constructs that include an engineered V_H sequence (whether or not associated with one or more other sequences, including immunoglobulin-derived and other sequences contained, for example, within an sFv or scFv-containing construct), which includes a mutation, alteration or deletion at an amino acid at a location or locations corresponding to amino acid position 11. The V_H 11 amino acid, if substituted, may be substituted with another amino acid as described herein, or by another molecule as desired.

Once a construct, such as for example a binding domain-immunoglobulin fusion protein, as provided herein has been designed, polynucleotides including DNAs encoding the construct, where it or a relevant portion of it is a polypeptide, may be synthesized in whole or in part via oligonucleotide synthesis as described, for example, in Sinha *et al.*, *Nucleic Acids Res.*, **12**: 4539-4557 (1984); assembled via PCR as described, for example in Innis, Ed., *PCR Protocols*, Academic Press (1990) and also in Better *et al.* *J. Biol. Chem.* **267**: 16712-16118 (1992); cloned and expressed via standard procedures as described, for example, in Ausubel *et al.*, Eds., *Current Protocols in Molecular Biology*, John Wiley & Sons,

New York (1989) and also in Robinson *et al.*, *Hum. Antibod. Hybridomas*, **2**: 84-93 (1991); and tested for desired activity, for example, binding to a target, or specific antigen binding activity, as described, for example, in Harlow *et al.*, Eds., *Antibodies: A Laboratory Manual*, Chapter 14, Cold Spring Harbor Laboratory, Cold Spring Harbor (1988) and Munson *et al.*, *Anal. Biochem.*, **107**: 220-239 (1980).

The preparation of single polypeptide chain binding molecules of the Fv region, single-chain Fv molecules, is known in the art. See, *e.g.*, U.S. Pat. No. 4,946,778. In the present invention, single-chain Fv-like molecules that may be included in constructs of the invention may be synthesized by encoding a first variable region of the heavy or light chain, followed by one or more linkers to the variable region of the corresponding light or heavy chain, respectively. The selection of various appropriate linker(s) between the two variable regions is described in U.S. Pat. No. 4,946,778 (see also, *e.g.*, Huston *et al.*, 1993 *Int. Rev. Immunol.* **10**: 195). An exemplary linker described herein is (Gly-Gly-Gly-Gly-Ser)₃, but may be of any desired length. The linker is used to convert the naturally aggregated but chemically separate heavy and light chains into the amino terminal antigen binding portion of a single polypeptide chain, for example, wherein this antigen binding portion will fold into a structure similar to the original structure made of two polypeptide chains, or that otherwise has the ability to bind to a target, for example a target antigen. For those constructs that include an scFv as a binding region, a native or engineered immunoglobulin hinge as a connecting region, and one or more native or engineered heavy chain constant regions as a binding region, nucleotide sequences encoding the variable regions of native or engineered heavy and light chains, joined by a sequence encoding a linker, are joined to a nucleotide sequence encoding native or engineered antibody constant regions, as desired. The constant regions may be those that permit the resulting polypeptide to form interchain disulfide bonds to form a dimer, and which contain desired effector functions, such as the ability to mediate ADCC, CDC, or fix complement, although native or engineered constant regions that do not favor dimer or other multimer formation or aggregation are preferred. For a construct, such as an immunoglobulin-like molecule, of the invention that is intended for use in humans, the included sequences having constant regions and/or desired constant regions function(s) will typically be human or substantially human or humanized to minimize a potential anti-human immune response and to provide

appropriate or desired effector functions. Manipulation of sequences encoding antibody constant regions is referenced in the PCT publication of Morrison and Oi, WO 89/07142. In preferred embodiments, the CH1 domain is deleted in whole or in part from a tail region that includes, or consists essentially of, or consists of, a native or engineered immunoglobulin constant region(s) (for example, native or engineered CH2 and/or CH3 constant region(s), or native or engineered CH2 and/or CH3 and/or CH4 constant region(s)), and the carboxyl end of the binding region, for example, a binding domain polypeptide such as an immunoglobulin variable region polypeptide, is joined to the amino terminus of, for example, a CH2 via a connecting region, for example, a native or engineered hinge region polypeptide as provided herein.

As described above, the present invention provides recombinant expression constructs capable of directing the expression of constructs of the invention, including binding domain-immunoglobulin fusion proteins, as provided herein. The amino acids, which occur in the various amino acid sequences referred to herein, are identified according to their well known three-letter or single-letter abbreviations. The nucleotides, which occur in the various DNA sequences or fragments thereof referred herein, are designated with the standard single letter designations used routinely in the art. A given amino acid sequence may also encompass similar but changed amino acid sequences, such as those having only minor changes, for example by way of illustration and not limitation, covalent chemical modifications, insertions, deletions and substitutions, which may further include conservative substitutions or substitutions with non-naturally-occurring amino acids. Amino acid sequences that are similar to one another may share substantial regions of sequence homology. In like fashion, nucleotide sequences may encompass substantially similar nucleotide sequences having only minor changes, for example by way of illustration and not limitation, covalent chemical modifications, insertions, deletions and substitutions, which may further include silent mutations owing to degeneracy of the genetic code. Nucleotide sequences that are similar to one another may share substantial regions of sequence homology.

The presence of a malignant condition in a subject refers to the presence of dysplastic, cancerous, and/or transformed cells in the subject, including, for example, neoplastic, tumor, non-contact inhibited, or oncogenically transformed cells, or the like (*e.g.*, melanoma, carcinomas such as adenocarcinoma, squamous cell

carcinoma, small cell carcinoma, oat cell carcinoma, *etc.*, sarcomas such as chondrosarcoma, osteosarcoma, *etc.*) which are known to the art and for which criteria for diagnosis and classification are established. In preferred embodiments contemplated by the present invention, for example, such cancer cells are malignant
5 hematopoietic cells, such as transformed cells of lymphoid lineage and in particular, B cell lymphomas and the like; cancer cells may in certain preferred embodiments also be epithelial cells such as carcinoma cells. The invention also contemplates B cell disorders, which may include certain malignant conditions that affect B cells (*e.g.*, B cell lymphoma) but which is not intended to be so limited, and which is also intended
10 to encompass autoimmune diseases and in particular, diseases, disorders and conditions that are characterized by autoantibody production, for example.

Autoantibodies are antibodies that react with self antigens.

Autoantibodies are detected in several autoimmune diseases (*i.e.*, a disease, disorder or condition wherein a host immune system generates an inappropriate anti-"self"
15 immune reaction) where they are involved in disease activity. Current treatments for various autoimmune diseases include immunosuppressive drugs that require continuing administration, lack specificity, and cause significant side effects. New approaches that can eliminate autoantibody production with minimal toxicity will address an unmet medical need for a spectrum of diseases that affect many people.
20 Constructs of the subject invention, including binding domain-immunoglobulin fusion proteins, are designed, for example, for improved penetration into lymphoid tissues. Depletion of B lymphocytes interrupts the autoantibody production cycle, and allows the immune system to reset as new B lymphocytes are produced from precursors in the bone marrow.

25 A number of diseases, disorders, and conditions have been identified for which beneficial effects are believed, according to non-limiting theory, to result from B cell depletion therapy. Such diseases disorders, and conditions include, but are not limited to, Grave's disease, Hashimoto's disease, rheumatoid arthritis, systemic lupus erythematosus, Sjogrens Syndrome Immune Thrombocytopenic purpura,
30 multiple sclerosis, myasthenia gravis, scleroderma, psoriasis, Inflammatory Bowel Disease including Crohn's disease and ulcerative colitis. Inflammatory Bowel Disease including Crohn's disease and Ulcerative colitis, are autoimmune diseases of the digestive system.

The present invention further relates to nucleotide constructs encoding constructs of the invention, for example, binding domain-immunoglobulin fusion proteins, and in particular to methods for administering recombinant constructs encoding such proteins for gene therapy applications that may be expressed, for example, as fragments, analogs and derivatives of such polypeptides.

The terms "fragment," "derivative" and "analog" when referring to constructs of the invention including, for example, binding domain-immunoglobulin fusion polypeptides or fusion proteins, refers to any construct, such as a binding domain-immunoglobulin fusion polypeptide or fusion protein, that retains essentially the same biological function or activity as such polypeptide. Thus, an analog includes a pro- or prepro-form of a construct, for example, a pro-protein that can be activated by cleavage of the pro-protein portion to produce an active construct, such as a binding domain-immunoglobulin fusion polypeptide.

A fragment, derivative or analog of a construct of the invention, for example, a binding domain-immunoglobulin fusion polypeptide or fusion protein, including binding domain-immunoglobulin fusion polypeptides or fusion proteins encoded by the cDNAs referred to herein, may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which additional amino acids are fused or otherwise connected to the construct, *e.g.*, a binding domain-immunoglobulin fusion polypeptide, including amino acids that are employed for detection or specific functional alteration of the construct, including such constructs as a binding domain-immunoglobulin fusion polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

The constructs, including polypeptide constructs, of the present invention include, for example, binding domain-immunoglobulin fusion polypeptides and fusion proteins having binding regions such as binding domain polypeptide amino acid sequences that are identical or similar to sequences known in the art, or fragments or portions thereof. For example by way of additional illustration and not limitation, a human CD154 molecule extracellular domain is contemplated for use

according to the instant invention, as are portions of such polypeptides and/or polypeptides having at least about 70% similarity (preferably greater than a 70% identity) and more preferably about 90% similarity (more preferably greater than a 90% identity) to the reported polypeptide and still more preferably about 95% similarity (still more preferably greater than a 95% identity) to the reported polypeptides and to portions of such polypeptides, wherein such portions of a binding domain-immunoglobulin fusion polypeptide, for example, generally contain at least about 30 amino acids and more preferably at least about 50 amino acids. Extracellular domains include, for example, portions of a cell surface molecule, and in particularly preferred embodiments cell surface molecules that are integral membrane proteins or that comprise a plasma membrane spanning transmembrane domain, that are constructed to extend beyond the outer leaflet of the plasma membrane phospholipid bilayer when the molecule is expressed at a cell surface, preferably in a manner that exposes the extracellular domain portion of such a molecule to the external environment of the cell, also known as the extracellular milieu. Methods for determining whether a portion of a cell surface molecule comprises an extracellular domain are well known to the art and include, for example, experimental determination (*e.g.*, direct or indirect labeling of the molecule, evaluation of whether the molecule can be structurally altered by agents to which the plasma membrane is not permeable such as proteolytic or lipolytic enzymes) or topological prediction based on the structure of the molecule (*e.g.*, analysis of the amino acid sequence of a polypeptide) or other methodologies.

As used herein, an "amino acid" is a molecule having the structure wherein a central carbon atom (the alpha (α)-carbon atom) is linked to a hydrogen atom, a carboxylic acid group (the carbon atom of which is referred to herein as a "carboxyl carbon atom"), an amino group (the nitrogen atom of which is referred to herein as an "amino nitrogen atom"), and a side chain group, R. When incorporated into a peptide, polypeptide, or protein, an amino acid loses one or more atoms of its amino and carboxylic groups in the dehydration reaction that links one amino acid to another. As a result, when incorporated into a protein, an amino acid may also be referred to as an "amino acid residue." In the case of naturally occurring proteins, an amino acid residue's R group differentiates the 20 amino acids from which proteins are typically synthesized, although one or more amino acid residues in a protein may

be derivatized or modified following incorporation into protein in biological systems (e.g., by glycosylation and/or by the formation of cystine through the oxidation of the thiol side chains of two non-adjacent cysteine amino acid residues, resulting in a disulfide covalent bond that frequently plays an important role in stabilizing the folded conformation of a protein, *etc.*). As those in the art will appreciate, non-naturally occurring amino acids can also be incorporated into proteins, particularly those produced by synthetic methods, including solid state and other automated synthesis methods. Examples of such amino acids include, without limitation, α -amino isobutyric acid, 4-amino butyric acid, L-amino butyric acid, 6-amino hexanoic acid, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norlensine, norvaline, hydroxproline, sarcosine, citralline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids (e.g., β -methyl amino acids, α -methyl amino acids, N α -methyl amino acids) and amino acid analogs in general. In addition, when an α -carbon atom has four different groups (as is the case with the 20 amino acids used by biological systems to synthesize proteins, except for glycine, which has two hydrogen atoms bonded to the α carbon atom), two different enantiomeric forms of each amino acid exist, designated D and L. In mammals, only L-amino acids are incorporated into naturally occurring polypeptides. The instant invention envisions proteins incorporating one or more D- and L-amino acids, as well as proteins comprised of just D- or L-amino acid residues.

Herein, the following abbreviations may be used for the following amino acids (and residues thereof): alanine (Ala, A); arginine (Arg, R); asparagine (Asn, N); aspartic acid (Asp, D); cyteine (Cys, C); glycine (Gly, G); glutamic acid (Glu, E); glutamine (Gln, Q); histidine (His, H); isoleucine (Ile, I); leucine (Leu, L); lysine (Lys, K); methionine (Met, M); phenylalanine (Phe, F); proline (Pro, P); serine (Ser, S); threonine (Thr, T); tryptophan (Trp, W); tyrosine (Tyr, Y); and valine (Val, V). Non-polar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionines. Neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, esparagine, and glutamine. Positively charged (basic amino acids include arginine, lysine and histidine. Negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

"Protein" refers to any polymer of two or more individual amino acids (whether or not naturally occurring) linked via a peptide bond, and occurs when the carboxyl carbon atom of the carboxylic acid group bonded to the α -carbon of one amino acid (or amino acid residue) becomes covalently bound to the amino nitrogen atom of amino group bonded to the α -carbon of an adjacent amino acid. The term "protein" is understood to include the terms "polypeptide" and "peptide" (which, at times, may be used interchangeably herein) within its meaning. In addition, proteins comprising multiple polypeptide subunits or other components will also be understood to be included within the meaning of "protein" as used herein. Similarly, fragments of proteins, peptides, and polypeptides are also within the scope of the invention and may be referred to herein as "proteins."

In biological systems (be they in vivo or in vitro, including cell-free, systems), the particular amino acid sequence of a given protein (*i.e.*, the polypeptide's "primary structure," when written from the amino-terminus to carboxy-terminus) is determined by the nucleotide sequence of the coding portion of a mRNA, which is in turn specified by genetic information, typically genomic DNA (which, for purposes of this invention, is understood to include organelle DNA, for example, mitochondrial DNA and chloroplast DNA). Of course, any type of nucleic acid which constitutes the genome of a particular organism (*e.g.*, double-stranded DNA in the case of most animals and plants, single or double-stranded RNA in the case of some viruses, *etc.*) is understood to code for the gene product(s) of the particular organism. Messenger RNA is translated on a ribosome, which catalyzes the polymerization of a free amino acid, the particular identity of which is specified by the particular codon (with respect to mRNA, three adjacent A, G, C, or U ribonucleotides in the mRNA's coding region) of the mRNA then being translated, to a nascent polypeptide. Recombinant DNA techniques have enabled the large-scale synthesis of proteins and polypeptides (*e.g.*, human insulin, human growth hormone, erythropoietin, granulocyte colony stimulating factor, *etc.*) having the same primary sequence as when produced naturally in living organisms. In addition, such technology has allowed the synthesis of analogs of these and other proteins, which analogs may contain one or more amino acid deletions, insertions, and/or substitutions as compared to the native proteins. Recombinant DNA technology also enables the synthesis of entirely novel proteins.

In non-biological systems (*e.g.*, those employing solid state synthesis), the primary structure of a protein (which also includes disulfide (cystine) bond locations) can be determined by the user. As a result, polypeptides having a primary structure that duplicates that of a biologically produced protein can be achieved, as can analogs of such proteins. In addition, completely novel polypeptides can also be synthesized, as can protein incorporating non-naturally occurring amino acids.

As is known in the art, "similarity" between two polypeptides may be determined by comparing the amino acid sequence (including conserved amino acid substitutes therein) of one polypeptide to the sequence of a second polypeptide. Fragments or portions of the nucleic acids encoding polypeptides of the present invention may be used to synthesize full-length nucleic acids of the present invention. As used herein, "% identity" refers to the percentage of identical amino acids situated at corresponding amino acid residue positions when two or more polypeptide are aligned and their sequences analyzed using, for example, a gapped BLAST algorithm (*e.g.*, Altschul *et al.*, 1997 *Nucl. Ac. Res.* **25**: 3389; Altschul *et al.*, 1990 *J. Mol. Biol.*, **215**: 403-410) which weights sequence gaps and sequence mismatches according to the default weightings provided by the National Institutes of Health/NCBI database (Bethesda, MD). Other alignment methods include BLITZ (MPsrch) (Sturrock & Collins, 1993), and FASTA (Pearson and Lipman, 1988 *Proc. Natl. Acad. Sci. USA.* **85**: 2444-2448).

The term "isolated" means, in the case of a naturally occurring material, that the material is or has been removed from, or is no longer associated with, its natural or original environment. For example, a naturally occurring nucleic acid or protein or polypeptide present in a living animal is not isolated, but the same nucleic acid or polypeptide, separated from some or all of the co-existing materials in the natural system, is isolated. Such nucleic acids could be part of a vector and/or such nucleic acids or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment. The term "isolated", in the case of non-naturally occurring material, such as a recombinantly manufactured construct of the invention, includes material that is substantially or essentially free from components which normally accompany it during manufacture, such as, for example, proteins and peptides that have been purified to a desired degree, preferably, for example, so that they are at least about 80% pure, more

preferably at least about 90%, and still more preferably at least about 95% as measured by techniques known in the art.

The term "gene" means a segment of DNA involved in producing a polypeptide chain; it may also include regions preceding and following a polypeptide coding region, for example, a "leader and trailer" as well as intervening sequences (introns) between relevant individual coding segments (exons).

As described herein, the invention provides constructs, including binding domain-immunoglobulin fusion proteins, that may be encoded in whole or in part by nucleic acids that have a binding region coding sequence such as, for example, a binding domain coding sequence fused or otherwise connected in frame to an additional native or engineered immunoglobulin domain encoding sequence to provide for expression of, for example, a binding domain polypeptide sequence fused or otherwise connected to an additional functional polypeptide sequence that permits, for example by way of illustration and not limitation, detection, functional alteration, isolation and/or purification of the fusion protein. Such fusion proteins may permit functional alteration of a binding domain by containing additional immunoglobulin-derived polypeptide sequences that influence behavior of the fusion product, for example (and as described above) by reducing the availability of sufhdyryl groups for participation in disulfide bond formation, and by conferring the ability to potentiate ADCC and/or CDC and/or fix complement.

Modification of a polypeptide may be effected by any means known to those of skill in this art. The preferred methods herein rely on modification of DNA encoding, for example, a fusion protein and expression of the modified DNA. DNA encoding one of the constructs of the invention, for example, one of the binding domain-immunoglobulin fusions discussed herein, for example, may be altered or mutagenized using standard methodologies, including those described below. For example, cysteine residues that may otherwise facilitate multimer formation or promote particular molecular conformations can be deleted from a polypeptide or replaced, *e.g.*, cysteine residues that are responsible for or participate in aggregate formation. If necessary, for example, the identity of cysteine residues that contribute to aggregate formation may be determined empirically, by deleting and/or replacing a cysteine residue and ascertaining whether the resulting protein aggregates in solutions containing physiologically acceptable buffers and salts. In addition, fragments of, for

example, binding domain-immunoglobulin fusions may be constructed and used. As noted above, counterreceptor/ligand binding domains for many candidate binding domain-immunoglobulin fusion have been delineated, such that one having ordinary skill in the art may readily select appropriate polypeptide domains for inclusion in encoded products of the instant expression constructs.

Conservative substitutions of amino acids are well-known and may be made generally without altering the biological activity of the resulting binding domain-immunoglobulin fusion protein molecule. For example, such substitutions are generally made by interchanging within the groups of polar residues, charged residues, hydrophobic residues, small residues, and the like. If necessary, such substitutions may be determined empirically merely by testing the resulting modified protein for the ability to bind to the appropriate cell surface receptors in *in vitro* biological assays, or to bind to appropriate antigens or desired target molecules.

The present invention further relates to nucleic acids which hybridize to constructs of the invention, including for example, binding domain-immunoglobulin fusion protein encoding polynucleotide sequences as provided herein, or their complements, as will be readily apparent to those familiar with the art, if there is at least about 70%, preferably at least about 80-85%, more preferably at least about 90%, and still more preferably at least about 95%, 96%, 97%, 98% or 99% identity between the sequences.

The present invention particularly relates to nucleic acids which hybridize under stringent conditions to, for example, the binding domain-immunoglobulin fusion encoding nucleic acids referred to herein. As used herein, to "hybridize" under conditions of a specified stringency is used to describe the stability of hybrids formed between two single-stranded nucleic acid molecules. Stringency of hybridization is typically expressed in conditions of ionic strength and temperature at which such hybrids are annealed and washed. The term "stringent conditions" refers to conditions that permit hybridization between polynucleotides. Stringent conditions can be defined by salt concentration, the concentration of organic solvent (*e.g.*, formamide), temperature, and other conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of organic solvents (*e.g.*, formamide), or raising the hybridization temperature. For example, stringent salt concentration will ordinarily be less than

about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, *e.g.*, formamide, while high stringency hybridization can be obtained in the presence of an organic solvent (*e.g.*, at least about 35% formamide, most preferably at least about 50% formamide). Stringent temperature conditions will ordinarily include temperatures of at least about 30° C, more preferably of at least about 37° C, and most preferably of at least about 42° C Varying additional parameters, for example, hybridization time, the concentration of detergent, *e.g.*, sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed, and are within the skill in the art. Other typical "high", "medium" and "low" stringency encompass the following conditions or equivalent conditions thereto: high stringency: 0.1x SSPE or SSC, 0.1% SDS, 65-C; medium stringency: 0.2x SSPE or SSC, 0.1% SDS, 50° C; and low stringency: 1.0x SSPE or SSC, 0.1% SDS, 50° C As known to those having ordinary skill in the art, variations in stringency of hybridization conditions may be achieved by altering the time, temperature and/or concentration of the solutions used for prehybridization, hybridization and wash steps, and suitable conditions may also depend in part on the particular nucleotide sequences of the probe used, and of the blotted, proband nucleic acid sample. Accordingly, it will be appreciated that suitably stringent conditions can be readily selected without undue experimentation where a desired selectivity of the probe is identified, based on its ability to hybridize to one or more certain proband sequences while not hybridizing to certain other proband sequences.

As used herein, preferred "stringent conditions" generally refer to hybridization that will occur only if there is at least about 90-95% and more preferably at least about 97% identity between the sequences. The nucleic acid constructs which hybridize to, for example, binding domain-immunoglobulin fusion encoding nucleic acids referred to herein, in preferred embodiments, encode polypeptides which retain substantially the same biological function or activity as, for example, the binding domain-immunoglobulin fusion polypeptides encoded by the cDNAs.

The nucleic acids of the present invention, also referred to herein as polynucleotides, may be in the form of RNA, for example, mRNA, or in the form of DNA, which DNA includes cDNA (also called "complementary DNA", which is a DNA molecule that is complementary to a specific messenger RNA), genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. A coding sequence which encodes a construct of the invention, for example, a binding domain-immunoglobulin fusion polypeptide for use according to the invention may contain portions that are identical to the coding sequence known in the art or described herein for portions thereof, or may be a different coding sequence, which, as a result of the redundancy or degeneracy of the genetic code, encodes the same construct or portion thereof, including all or a portion of a binding domain-immunoglobulin fusion polypeptide.

The nucleic acids which encode constructs of the invention, for example, binding domain-immunoglobulin fusion polypeptides, for use according to the invention may include, but are not limited to: only the coding sequence for the construct, such as a binding domain-immunoglobulin fusion polypeptide; the coding sequence for the construct, such as a binding domain-immunoglobulin fusion polypeptide and additional coding sequence; the coding sequence for the construct, such as a binding domain-immunoglobulin fusion polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequences 5' and/or 3' of the coding sequence for the binding domain-immunoglobulin fusion polypeptide or a portion(s) thereof, which for example may further include but need not be limited to one or more regulatory nucleic acid sequences that may be a regulated or regulatable promoter, enhancer, other transcription regulatory sequence, repressor binding sequence, translation regulatory sequence or any other regulatory nucleic acid sequence. Thus, the term "nucleic acid encoding" or "polynucleotide encoding" a construct, for example, a binding domain-immunoglobulin fusion protein, encompasses a nucleic acid which includes only coding sequence for, for example, a binding domain-immunoglobulin fusion polypeptide as well as a nucleic acid which includes additional coding and/or non-coding sequence(s).

Nucleic acids and oligonucleotides for use as described herein can be synthesized by any method known to those of skill in this art (*see, e.g.*, WO 93/01286,

U.S. application Ser. No. 07/723,454; U.S. Pat. No. 5,218,088; U.S. Pat. No. 5,175,269; U.S. Pat. No. 5,109,124). Identification of various oligonucleotides and nucleic acid sequences also involves methods known in the art. For example, the desirable properties, lengths and other characteristics of oligonucleotides useful for cloning are well known. In certain embodiments, synthetic oligonucleotides and nucleic acid sequences may be designed that resist degradation by endogenous host cell nucleolytic enzymes by containing such linkages as: phosphorothioate, methylphosphonate, sulfone, sulfate, ketyl, phosphorodithioate, phosphoramidate, phosphate esters, and other such linkages that have proven useful in antisense applications. See, *e.g.*, Agrwal *et al.*, *Tetrahedron Lett.* **28**: 3539-3542 (1987); Miller *et al.*, *J. Am. Chem. Soc.* **93**: 6657-6665 (1971); Stec *et al.*, *Tetrahedron Lett.* **26**: 2191-2194 (1985); Moody *et al.*, *Nucl. Acids Res.* **12**: 4769-4782 (1989); Uznanski *et al.*, *Nucl. Acids Res.* (1989); Letsinger *et al.*, *Tetrahedron* **40**: 137-143 (1984); Eckstein, *Annu. Rev. Biochem.* **54**: 367-402 (1985); Eckstein, *Trends Biol. Sci.* **14**: 97-100 (1989); Stein In: *Oligodeoxynucleotides. Antisense Inhibitors of Gene Expression*, Cohen, Ed, Macmillan Press, London, pp. 97-117 (1989); Jager *et al.*, *Biochemistry* **27**: 7237-7246 (1988).

In one embodiment, the present invention provides truncated components (*e.g.*, binding domain polypeptide, hinge region polypeptide, linker, *etc.*) for use in a construct of the invention, for example, a binding domain-immunoglobulin fusion protein. In another embodiment the invention provides nucleic acids encoding a construct of the invention, for example, a binding domain-immunoglobulin fusion protein having such truncated components. A truncated molecule may be any molecule that comprises less than a full length version of the molecule of interest. Truncated molecules provided by the present invention may include truncated biological polymers, and in preferred embodiments of the invention such truncated molecules may be truncated nucleic acid molecules or truncated polypeptides. Truncated nucleic acid molecules have less than the full length nucleotide sequence of a known or described nucleic acid molecule, where such a known or described nucleic acid molecule may be a naturally occurring, a synthetic, or a recombinant nucleic acid molecule, so long as one skilled in the art would regard it as a full length molecule. Thus, for example, truncated nucleic acid molecules that correspond to a gene sequence contain less than the full length gene where the gene

comprises coding and non-coding sequences, promoters, enhancers and other regulatory sequences, flanking sequences and the like, and other functional and non-functional sequences that are recognized as part of the gene. In another example, truncated nucleic acid molecules that correspond to a mRNA sequence contain less than the full length mRNA transcript, which may include various translated and non-translated regions as well as other functional and non-functional sequences.

In other preferred embodiments, truncated molecules are polypeptides that comprise less than the full length amino acid sequence of a particular protein or polypeptide component. As used herein "deletion" has its common meaning as understood by those familiar with the art, and may refer to molecules that lack one or more portions of a sequence from either terminus or from a non-terminal region, relative to a corresponding full length molecule, for example, as in the case of truncated molecules provided herein. Truncated molecules that are linear biological polymers such as nucleic acid molecules or polypeptides may have one or more of a deletion from either terminus of the molecule and/or one or more deletions from a non-terminal region of the molecule, where such deletions may be deletions of from about 1-1500 contiguous nucleotide or amino acid residues, preferably about 1-500 contiguous nucleotide or amino acid residues and more preferably about 1-300 contiguous nucleotide or amino acid residues, including deletions of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31-40, 41-50, 51-74, 75-100, 101-150, 151-200, 201-250 or 251-299 contiguous nucleotide or amino acid residues. In certain particularly preferred embodiments truncated nucleic acid molecules may have a deletion of about 270-330 contiguous nucleotides. In certain more preferred embodiments, truncated polypeptide molecules may have a deletion, for example, of about 80-140 contiguous amino acids.

The present invention further relates to variants of the herein referenced nucleic acids that encode fragments, analogs and/or derivatives of a construct of the invention, for example, a binding domain-immunoglobulin fusion polypeptide. The variants of the nucleic acids encoding constructs of the invention, for example, binding domain-immunoglobulin fusion proteins, may be naturally occurring allelic variants of one or more portions of the the nucleic acid sequences included therein, or non-naturally occurring variants of such sequences or portions or sequences, including sequences varied by molecular engineering using, for example,

methods known in the art for varying sequence. As is known in the art, an allelic variant is an alternate form of a nucleic acid sequence which may have at least one of a substitution, a deletion or an addition of one or more nucleotides, any of which does not substantially or undesirably alter the function of the encoded binding domain-immunoglobulin fusion polypeptide.

Variants and derivatives of constructs of the invention, for example, binding domain-immunoglobulin fusion proteins, may be obtained by mutations of nucleotide sequences encoding, for example, binding domain-immunoglobulin fusion polypeptides or any portion thereof. Alterations of the native amino acid sequence may be accomplished by any of a number of conventional methods. Mutations can be introduced at particular loci, for example, by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, for example, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene wherein predetermined codons can be altered by substitution, deletion or insertion. Exemplary methods of making such alterations are disclosed by Walder *et al.*, 1986 *Gene* **42**: 133; Bauer *et al.*, 1985 *Gene* **37**: 73; Craik, January 1985 *BioTechniques* 12-19; Smith *et al.*, January 1985 *Genetic Engineering: Principles and Methods BioTechniques* 12-19; Costa GL, *et al.*, "Site-directed mutagenesis using a rapid PCR-based method," 1996 *Methods Mol. Biol.* **57**: 239-48; Rashtchian A., "Novel methods for cloning and engineering genes using the polymerase chain reaction," 1995 *Curr Opin Biotechnol.* **6**(1): 30-6; Sharon J, *et al.*, "Oligonucleotide-directed mutagenesis of antibody combining sites," 1993 *Int Rev Immunol.* **10**(2-3): 113-27; Kunkel, 1985 *Proc. Natl. Acad. Sci. USA* **82**: 488; Kunkel *et al.*, 1987 *Methods in Enzymol.* **154**: 367; and, U.S. Pat. Nos. 4,518,584 and 4,737,462.

As an example, modification of DNA may be performed by site-directed mutagenesis of DNA encoding a protein combined with the use of DNA amplification methods using primers to introduce and amplify alterations in the DNA template, such as PCR splicing by overlap extension (SOE). Site-directed mutagenesis is typically effected using a phage vector that has single- and double-stranded forms, such as M13 phage vectors, which are well-known and commercially available. Other

suitable vectors that contain a single-stranded phage origin of replication may be used. See, *e.g.*, Veira *et al.*, 1987 *Meth. Enzymol.* **15**: 3. In general, site-directed mutagenesis is performed by preparing a single-stranded vector that encodes the protein of interest (*e.g.*, all or a component portion of a given binding domain-

5 immunoglobulin fusion protein). An oligonucleotide primer that contains the desired mutation within a region of homology to the DNA in the single-stranded vector is annealed to the vector followed by addition of a DNA polymerase, such as *E. coli* DNA polymerase I (Klenow fragment), which uses the double stranded region as a primer to produce a heteroduplex in which one strand encodes the altered sequence
10 and the other the original sequence. The heteroduplex is introduced into appropriate bacterial cells and clones that include the desired mutation are selected. The resulting altered DNA molecules may be expressed recombinantly in appropriate host cells to produce the modified protein.

Equivalent DNA constructs that include code for additions or
15 substitutions of amino acid residues or sequences, or deletions of terminal or internal residues or sequences not needed or desired for biological activity, for example, are also encompassed by the invention. For example, and as discussed above, sequences encoding Cys residues that are not desirable or essential for biological activity can be altered to cause the Cys residues to be deleted or replaced with other amino acids, for
20 example, thus preventing formation of incorrect or undesired intramolecular disulfide bridges upon synthesis or renaturation.

A "host cell" or "recombinant host cell" is a cell that contains a vector, *e.g.*, an expression vector, or a cell that has otherwise been manipulated by recombinant techniques to express a protein of interest. Host organisms include those
25 organisms in which recombinant production of constructs of the invention, for example, binding domain-immunoglobulin fusion products encoded by the recombinant constructs of the present invention may occur, such as bacteria (for example, *E. coli*), yeast (for example, *Saccharomyces cerevisiae* and *Pichia pastoris*), insect cells, and mammalian cells, including *in vitro* and *in vivo* expression. Host
30 organisms thus may include organisms for the construction, propagation, expression or other steps in the production of the compositions provided herein. Hosts include subjects in which immune responses take place, as described herein. Presently preferred host organisms for production of constructs of the invention that produce

glycosylated proteins are mammalian cells or other cells systems that permit the expression and recovery of glycosylated proteins. Other cell lines include inbred murine strains and murine cell lines, and human cells and cell lines.

A DNA construct encoding a desired construct of the invention, for example, a binding domain-immunoglobulin fusion protein, is introduced into a vector, for example, a plasmid, for expression in an appropriate host. In preferred embodiments, the host is a mammalian host, for example, a mammalian cell line. The sequence encoding the ligand or nucleic acid binding domain is preferably codon-optimized for expression in the particular host. Thus, for example, if a construct, for example, is a human binding domain-immunoglobulin fusion and is expressed in bacteria, the codons may be optimized for bacterial usage. For small coding regions, the gene can be synthesized as a single oligonucleotide. For larger proteins, splicing of multiple oligonucleotides, mutagenesis, or other techniques known to those in the art may be used. The sequences of nucleotides in plasmids or other vectors that are regulatory regions, such as promoters and operators, are operationally associated with one another for transcription. The sequence of nucleotides encoding a binding domain-immunoglobulin fusion protein may also include DNA encoding a secretion signal, whereby the resulting peptide is a precursor protein. The resulting processed protein may be recovered from the periplasmic space or the fermentation medium.

In preferred embodiments, the DNA plasmids may also include a transcription terminator sequence. As used herein, a "transcription terminator region" is a sequence that signals transcription termination. The entire transcription terminator may be obtained from a protein-encoding gene, which may be the same or different from the inserted binding domain-immunoglobulin fusion encoding gene or the source of the promoter. Transcription terminators are optional components of the expression systems herein, but are employed in preferred embodiments.

The plasmids or other vectors used herein include a promoter in operative association with the DNA encoding the protein or polypeptide of interest and are designed for expression of proteins in a suitable host as described above (*e.g.*, bacterial, murine, or human) depending upon the desired use of the plasmid (*e.g.*, administration of a vaccine containing binding domain-immunoglobulin fusion encoding sequences). Suitable promoters for expression of proteins and polypeptides herein are widely available and are well known in the art. Inducible promoters or

constitutive promoters that are linked to regulatory regions are preferred. Such promoters include, for example, but are not limited to, the T7 phage promoter and other T7-like phage promoters, such as the T3, T5 and SP6 promoters, the trp, lpp, and lac promoters, such as the lacUV5, from *E. coli*; the P10 or polyhedrin gene promoter of baculovirus/insect cell expression systems (see, e.g., U.S. Pat. Nos. 5,243,041, 5,242,687, 5,266,317, 4,745,051, and 5,169,784) and inducible promoters from other eukaryotic expression systems. For expression of the proteins such promoters are inserted in a plasmid in operative linkage with a control region such as the lac operon.

10 Preferred promoter regions are those that are inducible and functional in mammalian cells, for example. Examples of suitable inducible promoters and promoter regions for bacterial expression include, but are not limited to: the *E. coli* lac operator responsive to isopropyl β -D-thiogalactopyranoside (IPTG; see Nakamura *et al.*, 1979 *Cell* **18**: 1109-1117); the metallothionein promoter metal-regulatory-elements responsive to heavy-metal (e.g., zinc) induction (see, e.g., U.S. Pat. No. 4,870,009);
15 the phage T7lac promoter responsive to IPTG (see, e.g., U.S. Pat. No. 4,952,496; and Studier *et al.*, 1990 *Meth. Enzymol.* **185**: 60-89) and the TAC promoter. Depending on the expression host system to be used, plasmids may optionally include a selectable marker gene or genes that are functional in the host. Thus, for example, a selectable
20 marker gene includes any gene that confers a phenotype on bacteria that allows transformed bacterial cells to be identified and selectively grown from among a vast majority of untransformed cells. Suitable selectable marker genes for bacterial hosts, for example, include the ampicillin resistance gene (Amp^r), tetracycline resistance gene (Tc^r) and the kanamycin resistance gene (Kan^r). The kanamycin resistance gene
25 is presently preferred for bacterial expression.

In various expression systems, plasmids or other vectors may also include DNA encoding a signal for secretion of the operably linked protein. Secretion signals suitable for use are widely available and are well known in the art. Prokaryotic and eukaryotic secretion signals functional in *E. coli* may be employed. Depending on
30 the expression systems, presently preferred secretion signals may include, but are not limited to, those encoded by the following *E. coli* genes: ompA, ompT, ompF, ompC, beta-lactamase, and alkaline phosphatase, and the like (von Heijne, *J. Mol. Biol.* **184**: 99-105, 1985). In addition, the bacterial pelB gene secretion signal (Lei et al., *J.*

Bacteriol. **169**: 4379, 1987), the *phoA* secretion signal, and the *cek2* functional in insect cell may be employed. The most preferred secretion signal for certain expression systems is the *E. coli* *ompA* secretion signal. Other prokaryotic and eukaryotic secretion signals known to those of skill in the art may also be employed (see, e.g., von Heijne, *J. Mol. Biol.* **184**: 99-105, 1985). Using the methods described herein, one of skill in the art can substitute secretion signals that are functional in either yeast, insect or mammalian cells to secrete proteins from those cells.

Preferred plasmids for transformation of *E. coli* cells include the pET expression vectors (e.g., pET-11a, pET-12a-c, pET-15b; see U.S. Pat. No. 4,952,496; available from Novagen, Madison, WI). Other preferred plasmids include the pKK plasmids, particularly pKK 223-3, which contains the *tac* promoter (Brosius *et al.*, 1984 *Proc. Natl. Acad. Sci.* **81**: 6929; Ausubel *et al.*, *Current Protocols in Molecular Biology*; U.S. Pat. Nos. 5,122,463, 5,173,403, 5,187,153, 5,204,254, 5,212,058, 5,212,286, 5,215,907, 5,220,013, 5,223,483, and 5,229,279). Plasmid pKK has been modified by replacement of the ampicillin resistance gene with a kanamycin resistance gene. (Available from Pharmacia; obtained from pUC4K, see, e.g., Vieira *et al.* (1982 *Gene* **19**: 259-268; and U.S. Pat. No. 4,719,179.) Baculovirus vectors, such as pBlueBac (also called pJVETL and derivatives thereof), particularly pBlueBac III (see, e.g., U.S. Pat. Nos. 5,278,050, 5,244,805, 5,243,041, 5,242,687, 5,266,317, 4,745,051, and 5,169,784; available from Invitrogen, San Diego) may also be used for expression of the polypeptides in insect cells. Other plasmids include the pIN-IIIompA plasmids (see U.S. Pat. No. 4,575,013; see also Duffaud *et al.*, *Meth. Enz.* **153**: 492-507, 1987), such as pIN-IIIompA2.

Preferably, if one or more DNA molecules is replicated in bacterial cells, the preferred host is *E. coli*. The preferred DNA molecule is such a system also includes a bacterial origin of replication, to ensure the maintenance of the DNA molecule from generation to generation of the bacteria. In this way, large quantities of the DNA molecule can be produced by replication in bacteria. In such expression systems, preferred bacterial origins of replication include, but are not limited to, the *fl*-ori and *col E1* origins of replication. Preferred hosts for such systems contain chromosomal copies of DNA encoding T7 RNA polymerase operably linked to an inducible promoter, such as the *lacUV* promoter (see U.S. Pat. No. 4,952,496). Such hosts include, but are not limited to, lysogens *E. coli* strains HMS174(DE3)pLysS,

BL21(DE3)pLysS, HMS174(DE3) and BL21(DE3). Strain BL21(DE3) is preferred. The pLys strains provide low levels of T7 lysozyme, a natural inhibitor of T7 RNA polymerase.

5 The DNA molecules provided may also contain a gene coding for a repressor protein. The repressor protein is capable of repressing the transcription of a promoter that contains sequences of nucleotides to which the repressor protein binds. The promoter can be derepressed by altering the physiological conditions of the cell. For example, the alteration can be accomplished by adding to the growth medium a molecule that inhibits the ability to interact with the operator or with regulatory
10 proteins or other regions of the DNA or by altering the temperature of the growth media. Preferred repressor proteins include, but are not limited to the *E. coli* lacI repressor responsive to IPTG induction, the temperature sensitive λ cI857 repressor, and the like. The *E. Coli* lacI repressor is preferred.

In general, recombinant constructs of the subject invention will also
15 contain elements necessary for transcription and translation. In particular, such elements are preferred where the recombinant expression construct containing nucleic acid sequences encoding binding domain-immunoglobulin fusion proteins is intended for expression in a host cell or organism. In certain embodiments of the present invention, cell type preferred or cell type specific expression of a cell binding domain-
20 immunoglobulin fusion encoding gene may be achieved by placing the gene under regulation of a promoter. The choice of the promoter will depend upon the cell type to be transformed and the degree or type of control desired. Promoters can be constitutive or active and may further be cell type specific, tissue specific, individual cell specific, event specific, temporally specific or inducible. Cell-type specific
25 promoters and event type specific promoters are preferred. Examples of constitutive or nonspecific promoters include the SV40 early promoter (U.S. Pat. No. 5,118,627), the SV40 late promoter (U.S. Pat. No. 5,118,627), CMV early gene promoter (U.S. Pat. No. 5,168,062), and adenovirus promoter. In addition to viral promoters, cellular promoters are also amenable within the context of this invention. In particular,
30 cellular promoters for the so-called housekeeping genes are useful. Viral promoters are preferred, because generally they are stronger promoters than cellular promoters. Promoter regions have been identified in the genes of many eukaryotes including

higher eukaryotes, such that suitable promoters for use in a particular host can be readily selected by those skilled in the art.

Inducible promoters may also be used. These promoters include MMTV LTR (PCT WO 91/13160), inducible by dexamethasone; metallothionein promoter, inducible by heavy metals; and promoters with cAMP response elements, inducible by cAMP. By using an inducible promoter, the nucleic acid sequence encoding a binding domain-immunoglobulin fusion protein may be delivered to a cell by the subject invention expression construct and will remain quiescent until the addition of the inducer. This allows further control on the timing of production of the gene product.

Event-type specific promoters are active or up-regulated only upon the occurrence of an event, such as tumorigenicity or viral infection. The HIV LTR is a well known example of an event-specific promoter. The promoter is inactive unless the *tat* gene product is present, which occurs upon viral infection. Some event-type promoters are also tissue-specific.

Additionally, promoters that are coordinately regulated with a particular cellular gene may be used. For example, promoters of genes that are coordinately expressed may be used when expression of a particular binding construct of the invention, for example, a binding domain-immunoglobulin fusion protein-encoding gene is desired in concert with expression of one or more additional endogenous or exogenously introduced genes. This type of promoter is especially useful when one knows the pattern of gene expression relevant to induction of an immune response in a particular tissue of the immune system, so that specific immunocompetent cells within that tissue may be activated or otherwise recruited to participate in the immune response.

In addition to the promoter, repressor sequences, negative regulators, or tissue-specific silencers may be inserted to reduce non-specific expression of binding domain-immunoglobulin fusion protein encoding genes in certain situations, such as, for example, a host that is transiently immunocompromised as part of a therapeutic strategy. Multiple repressor elements may be inserted in the promoter region. Repression of transcription is independent on the orientation of repressor elements or distance from the promoter. One type of repressor sequence is an

insulator sequence. Such sequences inhibit transcription (Dunaway *et al.*, 1997 *Mol Cell Biol* **17**: 182-9; Gdula *et al.*, 1996 *Proc Natl Acad Sci USA* **93**: 9378-83, Chan *et al.*, 1996 *J Virol* **70**: 5312-28; Scott and Geyer, 1995 *EMBO J.* **14**: 6258-67; Kalos and Fournier, 1995 *Mol Cell Biol* **15**: 198-207; Chung *et al.*, 1993 *Cell* **74**: 505-14) and will silence undesired background transcription.

Repressor elements have also been identified in the promoter regions of the genes for type II (cartilage) collagen, choline acetyltransferase, albumin (Hu *et al.*, 1992 *J Cell Growth Differ.* **3**(9): 577-588), phosphoglycerate kinase (PGK-2) (Misuno *et al.*, 1992 *Gene* **119**(2): 293-297), and in the 6-phosphofructo-2-kinase/fructose-2, 6-bisphosphatase gene. (Lemaigre *et al.*, *Mol. Cell Biol.* **11**(2): 1099-1106). Furthermore, the negative regulatory element Tse-1 has been identified in a number of liver specific genes, and has been shown to block cAMP response element(CRE)-mediated induction of gene activation in hepatocytes. (Boshart *et al.*, 1990 *Cell* **61**(5): 905-916,).

In preferred embodiments, elements that increase the expression of the desired product are incorporated into the construct. Such elements include internal ribosome binding sites (IRES; Wang and Siddiqui, 1995 *Curr. Top. Microbiol. Immunol* **203**: 99; Ehrenfeld and Semler, 1995 *Curr. Top. Microbiol. Immunol.* **203**: 65; Rees *et al.*, 1996 *Biotechniques* **20**: 102; Sugimoto *et al.*, 1994 *Biotechnology* **12**: 694). IRES increase translation efficiency. As well, other sequences may enhance expression. For some genes, sequences especially at the 5' end inhibit transcription and/or translation. These sequences are usually palindromes that can form hairpin structures. Any such sequences in the nucleic acid to be delivered are generally deleted. Expression levels of the transcript or translated product are assayed to confirm or ascertain which sequences affect expression. Transcript levels may be assayed by any known method, including Northern blot hybridization, RNase probe protection and the like. Protein levels may be assayed by any known method, including ELISA, western blot, immunocytochemistry or other well known techniques.

Other elements may be incorporated into the constructs of the invention, for example, into binding domain-immunoglobulin fusion protein encoding constructs of the present invention. In preferred embodiments, the construct includes a transcription terminator sequence, including a polyadenylation sequence, splice donor

and acceptor sites, and an enhancer. Other elements useful for expression and maintenance of the construct in mammalian cells or other eukaryotic cells may also be incorporated (*e.g.*, origin of replication). Because the constructs are conveniently produced in bacterial cells, elements that are necessary for, or that enhance, propagation in bacteria are incorporated. Such elements include an origin of replication, a selectable marker and the like.

As provided herein, an additional level of controlling the expression of nucleic acids encoding constructs of the invention, for example, binding domain-immunoglobulin fusion proteins, delivered to cells for gene therapy, for example, may be provided by simultaneously delivering two or more differentially regulated nucleic acid constructs. The use of such a multiple nucleic acid construct approach may permit coordinated regulation of an immune response such as, for example, spatiotemporal coordination that depends on the cell type and/or presence of another expressed encoded component. Those familiar with the art will appreciate that multiple levels of regulated gene expression may be achieved in a similar manner by selection of suitable regulatory sequences, including but not limited to promoters, enhancers and other well known gene regulatory elements.

The present invention also relates to vectors, and to constructs prepared from known vectors that include nucleic acids of the present invention, and in particular to "recombinant expression constructs", including any of various known constructs, including delivery constructs, useful for gene therapy, that include any nucleic acids encoding, for example, binding domain-immunoglobulin fusion proteins and polypeptides according to the invention as provided herein; to host cells which are genetically engineered with vectors and/or other constructs of the invention and to methods of administering expression or other constructs comprising nucleic acid sequences encoding, for example, binding domain-immunoglobulin fusion polypeptides and fusion proteins of the invention, or fragments or variants thereof, by recombinant techniques.

Various constructs of the invention, including for example, binding domain-immunoglobulin fusion proteins, can be expressed in virtually any host cell, including *in vivo* host cells in the case of use for gene therapy, under the control of appropriate promoters, depending on the nature of the construct (*e.g.*, type of promoter, as described above), and on the nature of the desired host cell (*e.g.*, whether

postmitotic terminally differentiated or actively dividing; *e.g.*, whether the expression construct occurs in host cell as an episome or is integrated into host cell genome).

Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described, for example, by Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, NY, (1989); as noted herein, in particularly preferred embodiments of the invention, recombinant expression is conducted in mammalian cells that have been transfected or transformed with the subject invention recombinant expression construct. See also, for example, Machida, C A., "Viral Vectors for Gene Therapy: Methods and Protocols"; Wolff, J A, "Gene Therapeutics: Methods and Applications of Direct Gene Transfer" (Birkhauser 1994); Stein, U and Walther, W (eds. P, "Gene Therapy of Cancer: Methods and Protocols" (Humana Press 2000); Robbins, PD (ed.), "Gene Therapy Protocols" (Humana Press 1997); Morgan, J R (ed.), "Gene Therapy Protocols" (Humana Press 2002); Meager, A (ed.), "Gene Therapy Technologies, Applications and Regulations: From Laboratory to Clinic" (John Wiley & Sons Inc. 1999); Machida, C A and Constant, J G, "Viral Vectors for Gene Therapy: Methods and Protocols" (Humana Press 2002); "New Methods Of Gene Therapy For Genetic Metabolic Diseases NIH Guide," Volume 22, Number 35, Oct. 1, 1993. See also recent U.S. patents relating to gene therapy, including vaccines, which include U.S. Pat. No. 6,384,210 ("Solvent for biopolymer synthesis, solvent microdroplets and methods of use"); U.S. Pat. No. 6,384,203 ("Family of immunoregulators designated leukocyte immunoglobulin-like receptors (LIR)"); U.S. Pat. No. 6,384,202 ("Cell-specific active compounds regulated by the cell cycle"); U.S. Pat. No. 6,384,018 ("Polynucleotide tuberculosis vaccine"); U.S. Pat. No. 6,383,814 ("Cationic amphiphiles for intracellular delivery of therapeutic molecules"); U.S. Pat. No. 6,383,811 ("Polyampholytes for delivering polyions to a cell"); U.S. Pat. No. 6,383,795 ("Efficient purification of adenovirus"); U.S. Pat. No. 6,383,794 ("Methods of producing high titer recombinant adeno-associated virus"); U.S. Pat. No. 6,383,785 ("Self-enhancing, pharmacologically controllable expression systems"); U.S. Pat. No. 6,383,753 ("Yeast mammalian regulators of cell proliferation"); U.S. Pat. No. 6,383,746 ("Functional promoter for CCR5"); U.S. Pat. No. 6,383,743 ("Method for serial analysis of gene expression"); U.S. Pat. No. 6,383,738 ("Herpes simplex virus ORF P is a repressor of viral protein synthesis"); U.S. Pat. No. 6,383,737 ("Human

oxalyl-CoA Decarboxylase"); U.S. Pat. No. 6,383,733 ("Methods of screening for pharmacologically active compounds for the treatment of tumour diseases"); U.S. Pat. No. 6,383,522 ("Toxicity reduced composition containing an anti-neoplastic agent and a shark cartilage extract"); U.S. Pat. No. 6,383,512 ("Vesicular complexes and methods of making and using the same"); U.S. Pat. No. 6,383,481 ("Method for transplantation of hemopoietic stem cells"); U.S. Pat. No. 6,383,478 ("Polymeric encapsulation system promoting angiogenesis"); U.S. Pat. No. 6,383,138 ("Method for transdermal sampling of analytes"); U.S. Pat. No. 6,380,382 ("Gene encoding a protein having diagnostic, preventive, therapeutic, and other uses"); U.S. Pat. No. 6,380,371 ("Endoglycan: a novel protein having selectin ligand and chemokine presentation activity"); U.S. Pat. No. 6,380,369 ("Human DNA mismatch repair proteins"); U.S. Pat. No. 6,380,362 ("Polynucleotides, polypeptides expressed by the polynucleotides and methods for their use"); U.S. Pat. No. 6,380,170 ("Nucleic acid construct for the cell cycle regulated expression of structural genes"); U.S. Pat. No. 6,380,169 ("Metal complex containing oligonucleoside cleavage compounds and therapies"); U.S. Pat. No. 6,379,967 ("Herpesvirus saimiri as viral vector"); U.S. Pat. No. 6,379,966 ("Intravascular delivery of non-viral nucleic acid protease proteins, and uses thereof").

Typically, for example, expression constructs are derived from plasmid vectors. One preferred construct is a modified pNASS vector (Clontech, Palo Alto, CA), which has nucleic acid sequences encoding an ampicillin resistance gene, a polyadenylation signal and a T7 promoter site. Other suitable mammalian expression vectors are well known (*see, e.g.,* Ausubel *et al.*, 1995; Sambrook *et al.*, *supra*; *see also, e.g.,* catalogues from Invitrogen, San Diego, CA; Novagen, Madison, WI; Pharmacia, Piscataway, NJ; and others). Presently preferred constructs may be prepared that include a dihydrofolate reductase (DHFR) encoding sequence under suitable regulatory control, for promoting enhanced production levels of the binding domain-immunoglobulin fusion protein, which levels result from gene amplification following application of an appropriate selection agent (*e.g.,* methotrexate).

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence, as described above. The heterologous structural

sequence is assembled in appropriate phase with translation initiation and termination sequences. Thus, for example, the binding domain-immunoglobulin fusion protein encoding nucleic acids as provided herein may be included in any one of a variety of expression vector constructs as a recombinant expression construct for expressing a
5 binding domain-immunoglobulin fusion polypeptide in a host cell. In certain preferred embodiments the constructs are included in formulations that are administered *in vivo*. Such vectors and constructs include chromosomal, nonchromosomal and synthetic DNA sequences, *e.g.*, derivatives of SV40; bacterial plasmids; phage DNA; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral
10 DNA, such as vaccinia, adenovirus, fowl pox virus, and pseudorabies, or replication deficient retroviruses as described below. However, any other vector may be used for preparation of a recombinant expression construct, and in preferred embodiments such a vector will be replicable and viable in the host.

The appropriate DNA sequence(s) may be inserted into a vector, for
15 example, by a variety of procedures. In general, a DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Standard techniques for cloning, DNA isolation, amplification and purification, for enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like, and various separation techniques are those known and commonly employed by
20 those skilled in the art. A number of standard techniques are described, for example, in Ausubel *et al.* (1993 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., Boston, MA); Sambrook *et al.* (1989 *Molecular Cloning*, Second Ed., Cold Spring Harbor Laboratory, Plainview, NY); Maniatis *et al.* (1982 *Molecular Cloning*, Cold Spring Harbor Laboratory, Plainview, NY); Glover
25 (Ed.) (1985 *DNA Cloning Vol. I and II*, IRL Press, Oxford, UK); Hames and Higgins (Eds.), (1985 *Nucleic Acid Hybridization*, IRL Press, Oxford, UK); and elsewhere.

The DNA sequence in the expression vector is operatively linked to at least one appropriate expression control sequence(s) (*e.g.*, a constitutive promoter or a regulated promoter) to direct mRNA synthesis. Representative examples of such
30 expression control sequences include promoters of eukaryotic cells or their viruses, as described above. Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early

and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art, and preparation of certain particularly preferred recombinant expression constructs comprising at least one promoter or regulated promoter operably linked to a nucleic acid encoding an binding domain-immunoglobulin fusion polypeptide is described
5 herein.

Transcription of the DNA encoding proteins and polypeptides included within the present invention by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are *cis*-acting elements of DNA, usually
10 about from 10 to 300 bp that act on a promoter to increase its transcription. Examples including the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Gene therapy is the use of genetic material to treat disease. It
15 comprises strategies to replace defective genes or add new genes to cells and/or tissues, and is being developed for application in the treatment of cancer, the correction of metabolic disorders and in the field of immunotherapy. Gene therapies of the invention include the use of various constructs of the invention, with or without a separate carrier or delivery vehicle or constructs, for treatment of the diseases,
20 disorders, and/or conditions noted herein. Such constructs may also be used as vaccines for treatment or prevention of the diseases, disorders, and/or conditions noted herein. DNA vaccines, for example, make use of polynucleotides encoding immunogenic protein and nucleic acid determinants to stimulate the immune system against pathogens or tumor cells. Such strategies can stimulate either acquired or
25 innate immunity or can involve the modification of immune function through cytokine expression. *In vivo* gene therapy involves the direct injection of genetic material into a patient or animal model of human disease. Vaccines and immune modulation are systemic therapies. With tissue-specific *in vivo* therapies, such as those that aim to treat cancer, localized gene delivery and/or expression/targeting
30 systems are preferred. Diverse gene therapy vectors have been designed to target specific tissues, and procedures have been developed to physically target specific tissues, for example, using catheter-based technologies, all of which are contemplated herein. Ex vivo approaches to gene therapy are also contemplated herein and involve

the removal, genetic modification, expansion and re-administration of a patient's own cells. Examples include bone marrow transplantation for cancer treatment or the genetic modification of lymphoid progenitor cells. *Ex vivo* gene therapy is preferably applied to the treatment of cells that are easily accessible and can survive in culture during the gene transfer process (such as blood or skin cells).

Useful gene therapy vectors include adenoviral vectors, lentiviral vectors, Adeno-associated virus (AAV) vectors, Herpes Simplex Virus (Hsv) vectors, and retroviral vectors. Gene therapies may also be carried out using "naked DNA," liposome-based delivery, lipid-based delivery (including DNA attached to positively charged lipids), and electroporation.

As provided herein, in certain embodiments, including but not limited to gene therapy embodiments, the vector may be a viral vector such as, for example, a retroviral vector. Miller *et al.*, 1989 *BioTechniques* 7: 980; Coffin and Varmus, 1996 *Retroviruses*, Cold Spring Harbor Laboratory Press, NY. For example, retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus:

Retroviruses are RNA viruses which can replicate and integrate into the genome of a host cell via a DNA intermediate. This DNA intermediate, or provirus, may be stably integrated into the host cell DNA. According to certain embodiments of the present invention, an expression construct may comprise a retrovirus into which a foreign gene that encodes a foreign protein is incorporated in place of normal retroviral RNA. When retroviral RNA enters a host cell coincident with infection, the foreign gene is also introduced into the cell, and may then be integrated into host cell DNA as if it were part of the retroviral genome. Expression of this foreign gene within the host results in expression of the foreign protein.

Most retroviral vector systems which have been developed for gene therapy are based on murine retroviruses. Such retroviruses exist in two forms, as free viral particles referred to as virions, or as proviruses integrated into host cell DNA. The virion form of the virus contains the structural and enzymatic proteins of the

retrovirus (including the enzyme reverse transcriptase), two RNA copies of the viral genome, and portions of the source cell plasma membrane containing viral envelope glycoprotein. The retroviral genome is organized into four main regions: the Long Terminal Repeat (LTR), which contains *cis*-acting elements necessary for the
5 initiation and termination of transcription and is situated both 5' and 3' of the coding genes, and the three coding genes *gag*, *pol*, and *env*. These three genes *gag*, *pol*, and *env* encode, respectively, internal viral structures, enzymatic proteins (such as integrase), and the envelope glycoprotein (designated gp70 and p15e) which confers infectivity and host range specificity of the virus, as well as the "R" peptide of
10 undetermined function.

Separate packaging cell lines and vector producing cell lines have been developed because of safety concerns regarding the uses of retroviruses, including their use in expression constructs as provided by the present invention. Briefly, this methodology employs the use of two components, a retroviral vector and a packaging
15 cell line (PCL). The retroviral vector contains long terminal repeats (LTRs), the foreign DNA to be transferred and a packaging sequence (*y*). This retroviral vector will not reproduce by itself because the genes which encode structural and envelope proteins are not included within the vector genome. The PCL contains genes encoding the *gag*, *pol*, and *env* proteins, but does not contain the packaging signal "*y*". Thus, a
20 PCL can only form empty virion particles by itself. Within this general method, the retroviral vector is introduced into the PCL, thereby creating a vector-producing cell line (VCL). This VCL manufactures virion particles containing only the retroviral vector's (foreign) genome, and therefore has previously been considered to be a safe retrovirus vector for therapeutic use.

25 "Retroviral vector construct" refers to an assembly which is, within preferred embodiments of the invention, capable of directing the expression of a sequence(s) or gene(s) of interest, such as binding domain-immunoglobulin fusion encoding nucleic acid sequences. Briefly, the retroviral vector construct must include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second strand DNA
30 synthesis and a 3' LTR. A wide variety of heterologous sequences may be included within the vector construct, including for example, sequences which encode a protein (*e.g.*, cytotoxic protein, disease-associated antigen, immune accessory molecule, or

replacement gene), or which are useful as a molecule itself (*e.g.*, as a ribozyme or antisense sequence).

5 Retroviral vector constructs of the present invention may be readily constructed from a wide variety of retroviruses, including for example, B, C, and D type retroviruses as well as spumaviruses and lentiviruses (see, *e.g.*, RNA Tumor Viruses, Second Edition, Cold Spring Harbor Laboratory, 1985). Such retroviruses may be readily obtained from depositories or collections such as the American Type Culture Collection ("ATCC"; Rockville, MD), or isolated from known sources using commonly available techniques. Any of the above retroviruses may be readily utilized
10 in order to assemble or construct retroviral vector constructs, packaging cells, or producer cells of the present invention given the disclosure provided herein, and standard recombinant techniques (*e.g.*, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, 1989; Kunkle, 1985 *PNAS* 82: 488).

15 Suitable promoters for use in viral vectors generally may include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller, *et al.*, 1989 *Biotechniques* 7: 980-990, or any other promoter (*e.g.*, cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and P-actin promoters).
20 Other viral promoters which may be employed include, but are not limited to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein, and may be from among either regulated promoters or promoters as described above.

25 As described above, the retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, ψ -2, ψ -AM, PA12, T19-14x, VT-19-17-H2, ψ CRE, ψ CRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, *Human Gene Therapy*, 1: 5-14 (1990). The
30 vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO_4 precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include the nucleic acid sequence(s) encoding the binding domain-immunoglobulin fusion polypeptides or fusion proteins. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the binding domain-immunoglobulin fusion polypeptide or fusion protein. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, circulating peripheral blood mononuclear and polymorphonuclear cells including myelomonocytic cells, lymphocytes, myoblasts, tissue macrophages, dendritic cells, Kupffer cells, lymphoid and reticuloendothelia cells of the lymph nodes and spleen, keratinocytes, endothelial cells, and bronchial epithelial cells.

As another example of an embodiment of the invention in which a viral vector is used to prepare, for example, a recombinant binding domain-immunoglobulin fusion expression construct, in one preferred embodiment, host cells transduced by a recombinant viral construct directing the expression of binding domain-immunoglobulin fusion polypeptides or fusion proteins may produce viral particles containing expressed binding domain-immunoglobulin fusion polypeptides or fusion proteins that are derived from portions of a host cell membrane incorporated by the viral particles during viral budding.

In another aspect, the present invention relates to host cells containing the herein described nucleic acid constructs, such as, for example, recombinant binding domain-immunoglobulin fusion expression constructs. Host cells are genetically engineered (transduced, transformed or transfected) with the vectors and/or expression constructs of this invention which may be, for example, a cloning vector, a shuttle vector, or an expression construct. The vector or construct may be, for example, in the form of a plasmid, a viral particle, a phage, *etc.* The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying particular genes such as genes encoding binding domain-immunoglobulin fusion polypeptides or binding domain-immunoglobulin fusion fusion proteins. The culture conditions for particular host cells selected for expression, such as temperature, pH and the like, will be readily apparent to the ordinarily skilled artisan.

The host cell for production or expression of a construct of the invention, for example, can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Representative examples of appropriate host cells according to the present invention include, but need not be limited to, bacterial cells, such as *E. coli*, *Streptomyces*, *Salmonella typhimurium*; fungal cells, such as yeast; insect cells, such as *Drosophila S2* and *Spodoptera Sf9*; animal cells, such as CHO, COS or 293 cells; adenoviruses; plant cells, or any suitable cell already adapted to *in vitro* propagation or so established *de novo*. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the CQS-7 lines of monkey kidney fibroblasts, described by Gluzman, 1981 *Cell* 23: 175, and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences, for example as described herein regarding the preparation of binding domain-immunoglobulin fusion expression constructs. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Introduction of the construct into the host cell can be effected by a variety of methods with which those skilled in the art will be familiar, including but not limited to, for example, calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis *et al.*, 1986 *Basic Methods in Molecular Biology*).

The present invention constructs, for example, binding domain-immunoglobulin fusion proteins, or compositions comprising one or more polynucleotides encoding same as described herein (for example, to be administered under conditions and for a time sufficient to permit expression of a binding domain-immunoglobulin fusion protein in a host cell *in vivo* or *in vitro*, for gene therapy, for example, among other things), may be formulated into pharmaceutical compositions for administration according to well known methodologies. Pharmaceutical

compositions generally comprise one or more recombinant expression constructs, and/or expression products of such constructs, in combination with a pharmaceutically acceptable carrier, excipient or diluent. Such carriers will be nontoxic to recipients at the dosages and concentrations employed. For nucleic acid-based formulations, or for formulations comprising expression products of the subject invention recombinant constructs, about 0.01 ag/kg to about 100 mg/kg body weight will be administered, for example, typically by the intradermal, subcutaneous, intramuscular or intravenous route, or by other routes. A preferred dosage, for example, is about 1 µg/kg to about 1 mg/kg, with about 5 µg/kg to about 200 µg/kg particularly preferred. It will be evident to those skilled in the art that the number and frequency of administration will be dependent upon the response of the host.

"Pharmaceutically acceptable carriers" for therapeutic use are well known in the pharmaceutical art, and are described, for example, in *Remingtons Pharmaceutical Sciences*, Mack Publishing Co. (A. R. Gennaro edit. 1985). For example, sterile saline and phosphate-buffered saline at physiological pH may be used. Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. For example, sodium benzoate, sorbic acid and esters of *p*-hydroxybenzoic acid may be added as preservatives. *Id.* at 1449. In addition, antioxidants and suspending agents may be used. *Id.*

"Pharmaceutically acceptable salt" refers to salts of the compounds of the present invention derived from the combination of such compounds and an organic or inorganic acid (acid addition salts) or an organic or inorganic base (base addition salts). The compounds of the present invention may be used in either the free base or salt forms, with both forms being considered as being within the scope of the present invention.

The pharmaceutical compositions that contain one or more nucleic acid constructs of the invention, for example, binding domain-immunoglobulin fusion protein encoding constructs (or their expressed products) may be in any form which allows for the composition to be administered to a patient. For example, the composition may be in the form of a solid, liquid or gas (aerosol). Typical routes of administration include, without limitation, oral, topical, parenteral (*e.g.*, sublingually or buccally), sublingual, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal,

intracavernous, intrathecal, intrameatal, intraurethral injection or infusion techniques. The pharmaceutical composition is formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Compositions that will be administered to a patient take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of one or more compounds of the invention in aerosol form may hold a plurality of dosage units.

For oral administration, an excipient and/or binder may be present. Examples are sucrose, kaolin, glycerin, starch dextrins, sodium alginate, carboxymethylcellulose and ethyl cellulose. Coloring and/or flavoring agents may be present. A coating shell may be employed.

The composition may be in the form of a liquid, *e.g.*, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred compositions contain, in addition to one or more binding domain-immunoglobulin fusion construct or expressed product, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

A liquid pharmaceutical composition as used herein, whether in the form of a solution, suspension or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

It may also be desirable to include other components in the preparation, such as delivery vehicles including but not limited to aluminum salts, water-in-oil emulsions, biodegradable oil vehicles, oil-in-water emulsions, biodegradable microcapsules, and liposomes. Examples of immunostimulatory substances (adjuvants) for use in such vehicles include N-acetylmuramyl-L-alanine-D-isoglutamine (MDP), lipopoly-saccharides (LPS), glucan, IL-12, GM-CSF, gamma interferon and IL-15.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration and whether a sustained release is desired. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Pat. Nos. 4,897,268 and 5,075,109. In this regard, it is preferable that the microsphere be larger than approximately 25 microns.

Pharmaceutical compositions may also contain diluents such as buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with nonspecific serum albumin are exemplary appropriate diluents. Preferably, product is formulated as a lyophilizate using appropriate excipient solutions (*e.g.*, sucrose) as diluents.

As described above, the subject invention includes compositions capable of delivering nucleic acid molecules encoding binding domain-immunoglobulin fusion proteins. Such compositions include recombinant viral vectors (*e.g.*, retroviruses (*see* WO 90/07936, WO 91/02805, WO 93/25234, WO 93/25698, and WO 94/03622), adenovirus (*see* Berkner, 1988 *Biotechniques* 6: 616-627; Li *et al.*, 1993 *Hum. Gene Ther.* 4: 403-409; Vincent *et al.*, *Nat. Genet.* 5: 130-134; and Kolls *et al.*, 1994 *Proc. Natl. Acad. Sci. USA* 91: 215-219), pox virus (*see*

U.S. Pat. No. 4,769,330; U.S. Pat. No. 5,017,487; and WO 89/01973)), recombinant expression construct nucleic acid molecules complexed to a polycationic molecule (see WO 93/03709), and nucleic acids associated with liposomes (see Wang *et al.*, 1987 *Proc. Natl. Acad. Sci. USA* **84**: 7851). In certain embodiments, the DNA may be linked to killed or inactivated adenovirus (see Curiel *et al.*, 1992 *Hum. Gene Ther.* **3**: 147-154; Cotton *et al.*, 1992 *Proc. Natl. Acad. Sci. USA* **89**: 6094). Other suitable compositions include DNA-ligand (see Wu *et al.*, 1989 *J. Biol. Chem.* **264**: 16985-16987) and lipid-DNA combinations (see Felgner *et al.*, 1989 *Proc. Natl. Acad. Sci. USA* **84**: 7413-7417).

In addition to direct *in vivo* procedures, *ex vivo* procedures may be used in which cells are removed from a host, modified, and placed into the same or another host animal. It will be evident that one can utilize any of the compositions noted above for introduction of constructs of the invention, for example, binding domain-immunoglobulin fusion proteins or of binding domain-immunoglobulin fusion protein encoding nucleic acid molecules into tissue cells in an *ex vivo* context. Protocols for viral, physical and chemical methods of uptake are well known in the art.

Accordingly, the present invention is useful for treating a patient having a B cell disorder or a malignant condition, or for treating a cell culture derived from such a patient. As used herein, the term "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with cancer or a malignant condition, such as B cell lymphoma, or may be normal (*i.e.*, free of detectable disease and infection). A "cell culture" includes any preparation amenable to *ex vivo* treatment, for example a preparation containing immunocompetent cells or isolated cells of the immune system (including, but not limited to, T cells, macrophages, monocytes, B cells and dendritic cells). Such cells may be isolated by any of a variety of techniques well known to those of ordinary skill in the art (*e.g.*, Ficoll-hypaque density centrifugation). The cells may (but need not) have been isolated from a patient afflicted with a B cell disorder or a malignant condition, and may be reintroduced into a patient after treatment.

A liquid composition intended for either parenteral or oral administration should contain an amount of a construct of the invention, for example, a binding domain-immunoglobulin fusion protein encoding construct or expressed

product, such that a suitable dosage will be obtained. Typically, this amount is at least 0.01 wt % of a binding domain-immunoglobulin fusion construct or expressed product in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the composition.

- 5 Preferred oral compositions contain between about 4% and about 50% of binding domain-immunoglobulin fusion construct or expressed product(s). Preferred compositions and preparations are prepared so that, for example, a parenteral dosage unit contains between 0.01 to 1% by weight of active compound.

The pharmaceutical composition may be intended for topical
10 administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, beeswax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for
15 transdermal administration, the composition may include a transdermal patch or iontophoresis device. Topical formulations may contain a concentration of a construct of the invention, for example, a binding domain-immunoglobulin fusion construct or expressed product, of from about 0.1 to about 10% w/v (weight per unit volume).

The composition may be intended for rectal administration, in the
20 form, *e.g.*, of a suppository which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

In the methods of the invention, a construct of the invention, for
25 example, a binding domain-immunoglobulin fusion encoding constructs or expressed product(s), may be administered through use of insert(s), bead(s), timed-release formulation(s), patch(es) or fast-release formulation(s).

Constructs of the invention, for example, antigen-binding constructs of
the invention, may be administered or co-administered to an animal or patient in
30 combination with, or at the same or about the same time, as other compounds. In one aspect, one or more constructs, including for example one or more antigen-binding constructs, are administered to an animal or patient in conjunction with one or more

chemotherapeutic compounds such as alkylating agents, nucleoside analogues, and the like. The administration or co-administration of one or more constructs, including one or more antigen-binding constructs, of the invention and one or more chemotherapeutic agents can be used for the treatment of tumors or cancer in an animal or patient.

5 Exemplary cancers include, but are not limited to, head and neck cancer, breast cancer, colorectal cancer, gastric cancer, hepatic cancer, bladder cancer, cervical cancer, endometrial cancer, lung cancer (non-small cell), ovarian cancer, pancreatic cancer, prostate cancer; choriocarcinoma (lung cancer); hairy cell leukemia, chronic lymphocytic leukemia, acute lymphocytic leukemia (breast & bladder), acute
10 myelogenous leukemia, meningeal leukemia, chronic myelogenous leukemia, erythroleukemia. More commonly the cancers treated include non-Hodgkin's lymphoma (osteogenic sarcoma, adult soft tissue sarcoma), T-cell lymphoma, chronic lymphocytic leukaemia, slowly growing non-Hodgkin's lymphomas, Hodgkin's lymphoma and ovarian cancer.

15 Examples of an alkylating agents that can be co-administered with one or more constructs, including one or more antigen-binding constructs, of the invention include mechlorethamine, chlorambucil, ifosfamide, melphalan, busulfan, carmustine, lomustine, procarbazine, dacardazine, cisplatin, carboplatin, mitomycin C, cyclophosphamide, isosfamide, hexamethylmelamine, thiotepa, and dacarbazine, and
20 analogues thereof. See for example U.S. Pat. No. 3,046,301 describing the synthesis of chlorambucil, U.S. Pat. No. 3,732,340 describing the synthesis of ifosfamide, U.S. Pat. No. 3,018,302 for the synthesis of cyclophosphamide, U.S. Pat. No. 3,032,584 describing the synthesis of melphalan, and Braunwald et al., "Harrison's Principles of Internal Medicine," 15th Ed., McGraw-Hill, New York, NY, pp. 536-544 (2001) for
25 clinical aspects of cyclophosphamide, chlorambucil, melphalan, ifosfamide, procarbazine, hexamethylmelamine, cisplatin, and carboplatin. Examples of nucleoside analogues, include, but are not limited to, fludarabine pentostatin, methotrexate, fluorouracil, fluorodeoxyuridine, CB3717, azacitidine, cytarabine, floxuridine, mercaptopurine, 6-thioguanine, cladribine and analogues thereof. One
30 example is the combination of constructs, including antigen-binding constructs, that bind CD20. This construct acts as a chemosensitising agent and works together with chemotherapeutic agents, such that less chemotherapeutic agents are necessary to achieve anti-tumor or anti-cancer effects. For example, U.S. Pat. No. 3,923,785

describing the synthesis of pentostatin, U.S. Pat. No. 4,080,325 describing the synthesis of methotrexate, U.S. Pat. No. 2,802,005 describing the synthesis of fluorouracil, and Braunwald et al., "Harrison's Principles of Internal Medicine," 15th Ed., McGraw-Hill, New York, NY, pp. 536-544 (2001) for clinical aspects of
5 methotrexate, 5-fluorouracil, cytosine arabinoside, 6-mercaptopurine, 6-thioguanine, and fludarabine phosphate.

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered compounds that inhibit topoisomerase II or compounds that otherwise interact with
10 nucleic acids in cells. Such compounds include, for example, doxorubicin, epirubicin, etoposide, teniposide, mitoxantrone, and analogues thereof. In one example, this combination is used in treatment to reduce tumor cell contamination of peripheral blood progenitor cells (PBSC) in conjunction with high-dose chemotherapy and autologous stem cell support (HDC-ASCT). See U.S. Pat. No. 6,586,428 to Geroni *et*
15 *al.*

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with therapeutic drugs. For example, Virulizin (Lorus Therapeutics), which is believed to stimulate the release of tumour necrosis factor, TNF-alpha, by tumour
20 cells in vitro and stimulate activation of macrophage cells. This can be used in combination with one or more constructs, including one or more antigen-binding constructs, of the invention to increase cancer cell apoptosis and treat various types of cancers including Pancreatic Cancer, Malignant Melanoma, Kaposi's Sarcoma (KS), Lung Cancer, Breast Cancer, Uterine, Ovarian and Cervical Cancer. Another example
25 is CpG 7909 (Coley Pharmaceutical Group), which is believed to activate NK cells and monocytes and enhance ADCC. This drug can be used in combination with cancer or tumor specific constructs, including antigen-binding constructs, of the invention, such as an anti-CD20 construct, to treat non-Hodgkin's lymphoma and other cancers.

30 One or more constructs, including one or more antigen-binding constructs, of the invention can also be combined with angiogenesis inhibitors to increase anti-tumor effects. Angiogenesis is the growth of new blood vessels. This process allows tumors to grow and metastasize. Inhibiting angiogenesis can help

prevent metastasis, and stop the spread of tumors cells. Angiogenesis inhibitors include, but are not limited to, angiostatin, endostatin, thrombospondin, platelet factor 4, Cartilage-derived inhibitor (CDI), retinoids, Interleukin-12, tissue inhibitor of metalloproteinase 1, 2 and 3 (TIMP-1, TIMP-2, and TIMP-3) and proteins that block the angiogenesis signaling cascade, such as anti-VEGF (Vascular Endothelial Growth Factor) and IFN-alpha. Angiogenesis inhibitors can be administered or co-administered with tumor specific constructs, including antigen-binding constructs capable of mediating, for example, ADCC and/or complement fixation or chemotherapy-conjugated antigen-binding of the invention to combat various types of cancers, for example, solid tumor cancers such as lung and breast cancer.

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with disease modifying anti-rheumatic agents (DMAR agents) for the treatment of rheumatoid arthritis, psoriasis, ulcerative colitis, systemic lupus erythematosus (SLE), Crohn's disease, ankylosing spondylitis, and various inflammatory disease processes. In such treatment, the constructs, for example, antigen-binding constructs, of the invention are commonly administered in conjunction with compounds such as azathioprine, cyclosporin, gold, hydroxychloroquine, methotrexate, penicillamine, sulphasalazine, and the like.

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with agents or compounds that counteract the biological effects of interleukin-1, including for example interleukin-1 inhibitors and interleukin-1 receptor antagonist. It is thought that interleukin-1 has a role in the generation of rheumatoid arthritis (RA), inflammation, and the destruction of joints. IL-1 inhibitors can also be used in conjunction with the constructs, including antigen-binding constructs, of the invention to treat arthritis, inflammatory bowel disease, sepsis and septic shock, ischemic injury, reperfusion, ischemic brain injury such as cerebral palsy and multiple sclerosis. See U.S. Pat. No. 6,159,460 to Thompson *et al.* In another aspect, for example, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered to an animal or patient in conjunction with one or more glucocorticoids for example, methylprednisilone, dexamethasone, hydrocortisone, and the like. Glucocorticoids have been used to

induce apoptosis and inhibit growth, independent of ADCC and CDC. These compounds can be combined with constructs, including antigen-binding constructs, of the invention capable of inducing apoptosis in cancer cells. In one example is the anti-CD20, and anti-CD40 antigen-binding constructs, which can be used to induce apoptosis in B-cells, are combined with glucocorticoids to treat B-cell non-Hodgkin's lymphoma (NHL).

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with p38 inhibitors or antagonists. The p38 mitogen-activated protein kinase pathway is involved in a number of cellular processes instrumental to the development of rheumatoid arthritis. For example, the activation and infiltration of leukocytes as well as the production of inflammatory cytokines are p38-dependent processes.

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention are administered or co-administered with compounds that promote the differentiation and proliferation of B-cells. Cytokines such as interleukin-4 (IL-4) and interleukin-6 (IL-6), in addition to other biological activities, have been shown to stimulate antibody synthesis and secretion by activated B lymphocytes. In a particular aspect of the invention, constructs, including antigen-binding constructs, that recognize and bind CD20 are co-administered with one or more of interleukin-4 (IL-4) and interleukin-6 (IL-6).

In another aspect one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with Interleukin-2 (IL-2). Interleukin 2 (IL-2) is a lymphokine that increases production of effector cells, such as CD4+ T-helper cells, CD8 cytotoxic cells, antibody producing B cells, natural killer cells (NK), and monocytes/macrophages. IL-2 helps produce T-cells, which in turn secrete more of the IL-2 (an "autocrine loop"). IL-2 can be used to augment antibody-dependent cell-mediated cytotoxicity (ADCC) and immunotherapies associated with constructs of the invention. In one example, an anti-CD20 construct of the invention and IL-2 are used to treat patients with relapsed or refractory follicular non-Hodgkin's lymphoma. In another example IL-2 is administered or co-administered with HIV immunotherapies to help with T cell recovery.

In another aspect one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with Interleukin-12 (IL-12). IL-12 is known to enhance cytolytic T-cell responses, promote the development of helper T cells, enhance the activity of natural killer (NK) cells, and induces the secretion of IFN- γ in T and NK cells. IL-12 also increases many helper and effector cells that mediate apoptosis. In another aspect of the invention, one or more constructs, including one or more antigen-binding constructs, are administered or co-administered with IL-12 in the treatment of an animal or patient with a tumor or cancer. For example, a construct, including an antigen-binding construct, of the invention that binds CD20 combined with IL-2 for the treatment of a patient with B-cell non-Hodgkin's lymphoma (NHL).

One or more constructs, including one or more antigen-binding constructs, of the invention can also be combined with immunomodulators to boost the efficacy of the antigen-binding constructs of the invention. Immunomodulators include, but are not limited to, Colony Stimulating Factors (CSF), Tumor necrosis Factors (TNF), and Interferons (IFN).

CSFs can include granulocyte-macrophage CSF (GM-CSF), granulocyte-CSF (G-CSF), and macrophage CSF (M-CSF). GM-CSF is thought to regulate the development of neutrophils, macrophages, monocytes and eosinophils. G-CSF has been shown to induce neutrophil production, and M-CSF production. M-CSF has been shown to stimulate macrophages and monocytes. The use of CSFs to treat neutropenia in cancer patients has been long established. In one example, constructs, including antigen-binding constructs, of the invention can be combined with GM-CSF, G-CSF or combinations thereof in order to accelerate recovery from neutropenia in patients after bone marrow transplantation and chemotherapy. Neutrophils play a major role in fighting microbes such as bacterial, fungi and parasites. Patients with neutropenia are particularly susceptible to bacterial and wide spread fungal infections. In another example, a construct, including an antigen binding construct, of the invention can be combined with GM-CSF-treated neutrophils, monocytes and macrophages to increase activity against bacteria, fungi, etc, including the dreaded *Pneumocystis carinii*.

An example of an IFN is interferon alpha (IFN- α). IFN- α is made naturally by some types of white blood cell as part of the immune response when the

body reacts to cancers or viral infections. It has two main modes of attack, interfering with growth and proliferation of cancer cells and it boosting the production of killer T cells and other cells that attack cancer cells. Interferon is also thought to facilitate cancer cells to put out chemical signals that make them better targets for the immune system, and has been used in recent years for several different types of cancer, particularly kidney cancer, melanoma, multiple myeloma, and some types of leukemia. It is also used to treat viral infections such as hepatitis. Interferon-alpha2a, for example, enhances ADCC and can be combined with one or more constructs, including antigen-binding constructs, of the invention to increase the efficiency of ADCC activity associated with the construct. In another example, one or more constructs, including one or more antigen-binding constructs of the invention are administered or co-administered to an animal or patient with interferon-gamma (IFN- γ), which has been show to increase the number of anti-CD20 antigens on B cells and bone marrow plasma cells (BMPC). This is particularly useful for the treatment of patients with multiple myelomas, which have a reduced expression of CD20 in their B cells and bone marrow plasma cells (BMPC). Accordingly, the treatment of multiple myeloma patients with constructs, including antigen-binding constructs of the invention, in particular constructs that bind CD20, may be usefully co-administered in conjunction with IFN- γ .

TNF is a class of natural chemicals with anticancer properties. One example of a TNF is TNF-alpha. TNF-alpha has also been shown to have synergistic effects with IFN-gamma and IL-12. In another example, TNF can be administered or co-administered with one or more tumor specific constructs, including one or more antigen-binding constructs, of the invention, and include chemotherapy-conjugated antigen binding constructs of the invention, together with IFN-gamma, IL-12 or various combinations thereof. TNF is also known to be an inflammatory regulation molecule. TNF-alpha antibodies or antagonist(s) can be combined with anti-T cell constructs, including antigen-binding constructs, of the invention to treat patients with rheumatoid arthritis, psoriasis, ulcerative colitus, systemic lupus erythematosus (SLE), Crohn's disease, ankylosing spondylitis, and various inflammatory disease processes.

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered

with another antibody or antigen-binding construct of the invention. One example is a construct, for example, an antigen-binding construct of the invention capable of binding CD20 combined with a construct capable of binding CD22, CD19 or combinations thereof. This combination is effective as a treatment for indolent and aggressive forms of B-cell lymphomas, and acute and chronic forms of lymphatic leukemias. See U.S. Pat. No. 6,306,393 to Goldberg. In another example, constructs, including antigen-binding constructs, of the invention are co-administered with other constructs such as antigen-binding constructs of the invention that aid in mediating apoptosis. For example, a combination of one or more constructs, including one or more antigen-binding constructs of the invention capable of binding CD28, CD3, CD20 or a combination thereof. The combination of anti-CD28 and CD3 provides a method for prolonged proliferation of T-cells. See U.S. Pat. No. 6,352,694 to June et al. This prolonged T-cell proliferation increases the efficiency immune dependent cytotoxicity, particularly those associated with anti-CD20.

In another aspect, constructs, including antigen-binding constructs, of the invention can be administered or co-administered with one or more T-cell regulatory molecules. One example is a combination with interleukin-12 (IL-12). The IL-12 cytokine stimulates cell-mediated immunity, has angiostatic activity, and possesses significant anti-tumor effects in a variety of tumor models. IL-12 has also been shown to stimulate the production of interferon-gamma (IFN- γ). Accordingly, the treatment of multiple myeloma patients with one or more constructs, including one or more antigen-binding constructs, of the invention, in particular those that bind CD20, is expected to be more efficacious when co-administered in conjunction with IL-12. In another example, one or more constructs, including one or more antigen-binding constructs, of the invention can be administered or co-administered with a binding-domain construct of the invention other protein capable of binding CTLA-4 to enhance the anti-tumor immune response, by inhibiting the downregulation of T-cell activation.

In another aspect, one or more constructs, including one or more antigen-binding constructs, of the invention can be combined with gene therapies. In one example, a chemotherapy-conjugated construct of the invention is administered or co-administered with the Bcl-2 antisense oligonucleotide. Bcl-2 is associated with tumor resistance to anti-cancer therapies, and its believed to blocking chemotherapy-

induced cell death. In another example one or more constructs, including one or more antigen-binding constructs, of the invention is administered or co-administered with an adenovirus for delivery of a "suicide gene." The adenovirus inserts the gene directly into the tumor cells, which makes these cells sensitive to an otherwise
5 ineffective drug. Drug treatment then destroys the tumor cells, while leaving healthy cells untouched. However, once therapy is complete stray cancer cells that escaped therapy can reestablish and metastasize. Combining gene therapy with one or more constructs, including one or more antigen-binding constructs, will help kill stray cancer cells and minimize cancer reoccurrence.

10 A similar combination can be used with palliative (non-radical) operations to surgically remove tumors. In this example one or more constructs, including one or more antigen-binding constructs, of the invention can be administered before and after surgical extractions of tumors in order to increase the immune response and reduce the likelihood of reoccurrence by killing any cancer
15 cells that were not removed during the surgery.

 Another aspect combines a cancer or antigen vaccine and T-cell regulator molecules. For example, the binding portion, for example, an antigen-binding portion, of a construct can be specific for a cancer cell or antigen, or a protein fragment from a cancer cell or antigen. This can help mediate an immune response
20 against a particular tumor or antigen. Such constructs can be combined with T-cell regulators to increase the efficiency of the immune response.

 In another example, one or more constructs, including one or more antigen-binding constructs, of the invention is administered or co-administered with retinoids. Retinoids include Vitamin A and its derivatives, which have the ability to
25 stop cells from dividing and cause them to differentiate. Vitamin A is combined with an anti-cancer construct(s), including antigen-binding construct(s), of the invention to combat various forms of cancer.

 The terms "binding construct" and "antigen-binding construct" as used herein may refer to, for example, engineered polypeptides, recombinant polypeptides,
30 synthetic, semi-synthetic or other fusion proteins that are capable of binding a target, for example, an antigen. Antigen-binding constructs of the invention may be used in various applications, including those within the variety of uses to which antibodies or

related immunoglobulin-type constructs may be put. Constructs, including antigen-binding constructs of the invention can be used in *in vivo* and *in vitro* experiments for therapeutic, diagnostic, research, and other purposes. Such uses include, for example, the following.

5 Constructs, including antigen-binding constructs of the invention may be used for immunohistochemistry applications. For example, they may be used for immunolocalization of a particular antigen or group of antigens in a tissue. Tissue can be fixed and incubated with antigen-binding constructs of interest. These constructs can then be localized using a secondary antibody or binding construct of the invention
10 coupled to a label, for example, to a gold particle or an enzyme that gives a chemical reaction, like horseradish peroxidase or beta-galactosidase. A secondary antibody or binding construct is frequently made that is reactive against, for example, a portion of the primary binding construct. Thus, for example, if the primary binding construct has a human tail portion, the secondary antibody or binding construct could be, for
15 example, a rabbit anti-mouse antibody or antigen-binding construct that has been linked to beta-galactosidase. Alternatively the antibody or binding construct of the invention can be purified and then conjugated to another molecule to produce a fluorescent antibody or binding construct.

 Constructs, including antigen-binding constructs of the invention can
20 also be used to detect the location of an antigen or antigens on the surface of cells or to detect the location of intracellular materials using, for example, Immunoelectron Microscopy. Electron dense materials such as ferritin or colloidal gold, for example, can be conjugated to an antigen-binding construct. Scanning electron microscopy can be used to detect the localization of the antigen/binding construct complex.

25 Constructs, including antigen-binding constructs of the invention may also be used to quantitate the presence of an antigen or antigens using one of a variety of immunoassay formats, for example, a radioimmunoassay (RIA) format or an enzyme-linked immunosorbent assay (ELISA) format. There are many variants of these approaches, but those are based on a similar idea. For example, if an antigen can
30 be bound to a solid support or surface, or is in solution, it can be detected by reacting it with a specific antigen-binding construct of the invention. The presence or amount of the construct can then be detected or quantitated by reacting it with, for example, either a secondary antibody or a second antigen-binding construct of the invention by

incorporating a label directly into the primary antibody. Alternatively, for example, an antigen-binding polypeptide of the invention can be bound to a solid surface and the antigen added. A second antibody or antigen-binding polypeptide(s) of the invention that recognizes a distinct epitope on the antigen can then be added and detected. This
5 technique is commonly referred to as a "sandwich assay", which is frequently used to avoid problems of high background or non-specific reactions, among other reasons.

Because the binding constructs of the invention can have high affinity/affinities and/or selectivity/selectivities for a particular epitope or epitopes, they can also be used as affinity reagents, for example, in protein or antigen
10 purification. In one example of such a process, antigen-binding constructs of the invention are immobilized on a suitable support, for example, Sephadex resin or filter paper. The immobilized construct is exposed to a sample containing, or suspected of containing, a target protein(s) or antigen(s). The support is rinsed with a suitable buffer that will remove unwanted materials. The support is washed with another
15 buffer that will release the bound protein(s) or antigen(s).

Because particular binding constructs of the invention can bind to proteins or other antigens with high affinity and selectivity they can also be used as a criterion for the importance of a particular enzyme or other macromolecule in a particular reaction. If an antigen-binding construct of the invention can interfere with
20 a reaction in a solution, this will indicate that the construct may be binding specifically to a protein or other antigenic material involved in that reaction.

Constructs, including antigen-binding constructs of the invention can also be used as receptor blockers or inhibitors or antagonists.

Constructs, including antigen-binding constructs of the invention can
25 also be used in identifying and studying the function(s) of proteins. If an antigen-binding construct of the invention reacts with a specific protein, for example, that protein can subsequently be precipitated from solution, for example. Precipitation is typically performed by using a secondary antibody or antigen-binding construct of the invention that links primary complexes together. Alternatively, the complex can be
30 removed by reacting the solution with either protein A or, for example, depending on the construct, an anti-Fc antibody, for example, which has been attached to beads, for example, so that can be easily removed from the solution.

Constructs, including antigen-binding constructs of the invention can also be used in conjunction with gel-shift experiments to identify specific nucleic acid-binding proteins such as DNA-binding proteins. For example, DNA-binding proteins can be assayed by their ability to bind with high affinity to a particular
5 oligonucleotide. The mobility of an oligonucleotide associated with the protein is far different than the mobility of a free oligonucleotide and results in a gel migration pattern and signal that is commonly referred to as a gel shift. The addition of the construct to the binding assay can have either of two effects. If the construct binds to a region of a protein not involved in DNA binding it can result in a complex that has
10 even a slower mobility and is detected as a greater shift in mobility (a super-shift). Alternatively, if the construct binds to a region of the protein involved in recognizing the DNA then it can disrupt the binding and eliminate the shift. In either case, the data from these experiments can serve as a criterion to identify a DNA-binding protein, for example.

15 It is also possible to use constructs, including antigen-binding constructs of the invention to detect a protein by western blotting after fractionation by SDS-PAGE, for example. Once fractionated proteins are transferred to a membrane such as a nitrocellulose sheet, they are exposed to a particular antigen-binding construct of the invention that specifically recognizes, or recognizes to a
20 desired degree of selectivity, proteins immobilized to the blot. This allows particular proteins to be identified. This approach is particularly useful if the mobility of the protein changes during an experiment. For example, incorporation of a phosphate or a carbohydrate, or cleavage of the protein, results in a change in mobility that can be followed in straight forward manner by western analysis. With appropriate controls,
25 this approach can be used to measure the abundance of a protein in response to experimental manipulations.

The combination of SDS gels and immunoprecipitation can also be extremely effective. If a particular protein can be immunoprecipitated in a solution, both supernatant and precipitated fractions can be separated on an SDS gel and
30 studied using an antigen-binding constructs of the invention.

Sometimes a binding construct of the invention directed against one protein will also precipitate a second protein that interacts with the first protein. The second protein, as well as the first, can then be seen by staining the gel or by

autoradiography. This relationship is frequently the first indication that a protein functions as part of a complex and it can also be used to demonstrate a physical interaction of two proteins that are hypothesized to interact on the basis of other evidence (*e.g.*, a two hybrid screen or a supressor mutation). This approach can be
5 combined with western blotting analysis in several extremely effective ways.

Thus, for example, antigen-binding constructs of the invention can be used in a combination of immunoprecipitation and western analysis in the study, for example, of signal transduction and protein processing. For example, an immunoprecipitated protein can be subsequently studied by western analysis using a
10 different antibody or antigen-binding construct of the invention that binds to the protein. The most useful of are those that are directed against particular structural determinants that may be present in a protein. Thus, an antibody or antigen-binding construct of the invention directed against a region of the protein that undergoes proteolytic processing can be useful to follow proteolytic processing. Additionally, a
15 construct of the invention or a mixture of antigen-binding constructs of the invention that recognize phosphorylated peptides (*e.g.*, anti PY (phosphorylated tyrosine) can be used to follow the extent of phosphorylation of a protein (using western analysis) after it is precipitated, or *visa versa*. Glycosylation reactions can also be followed by antigen-binding constructs of the invention directed against a carbohydrate epitope (or
20 by lectins, *i.e.*, proteins that recognize carbohydrates). Likewise, some antigen-binding constructs of the invention can be made that specifically recognize a phosphorylated epitope, for example, that will recognize a tyrosine or a serine residue after phosphorylation, but will not bind (or detectably bind) the epitope in the absence of phosphate. This approach can be used to determine the phosphorylation state of a
25 particular protein. For example, the phosphorylation of CREB (the cAMP response element binding protein) can be followed by an antibody that specifically recognizes an epitope in a way that is dependent on the phosphorylation of serine 133.

Constructs, including antigen-binding constructs of the invention can also be used to screen expression libraries to isolate candidate polynucleotides that
30 express or present a particular epitope, or that have a particular affinity or expression characteristic.

Constructs, including antigen-binding constructs of the invention that bind to a cell surface can also be used as a marker to quantitate the fraction of cells

expressing that marker using flow cytometry. If different antigen-binding constructs of the invention/fluorescent dye combinations are used, for example, the fraction of cells expressing several antigens can be determined.

Constructs, including antigen-binding constructs of the invention that
5 function like anti-idiotypic antibodies, *i.e.*, antibodies against the binding domain of another antibody, can be used in any of a number of methods in which it would be desirable or useful to mimic the structure of an antigen. Such uses include, for example, uses as cancer vaccines (including antigen-binding constructs of the invention that incorporate a molecular adjuvant), as probes for receptors, as receptor
10 agonists, as receptor antagonists, as receptor blockers or inhibitors, and so on.

In another aspect, constructs, including antigen-binding constructs of the invention may be bispecific and thus capable of binding to two distinct epitopes, which may be present on the same or different cell types.

In vivo uses of constructs of the invention, including antigen-binding
15 constructs, include therapy, alone or in combination with one or more other therapies, for various diseases including cancers as well as B-cell disorders including autoimmune diseases. In some cases the constructs of the invention are administered to a patient. In other cases, the construct may be coupled to another molecule by techniques known in the art, for example, a fluorescent molecule to aid in imaging a
20 target, or a therapeutic drug and/or a toxin to aid in killing a target.

For example, a labeling molecule or atom can be conjugated or otherwise linked to the antigen-binding construct of the invention to aid in imaging or as a diagnostic agent. These include, but are not limited to enzymatic labels, radioisotopes or radioactive compounds or elements, fluorescent compounds or
25 metals, chemiluminescent compounds and bioluminescent compounds. Thus, binding constructs or antigen-binding constructs of the invention can be conjugated to a drug, which allows specific drug targeting and increased efficiency once the drug reaches the target. This facilitates drug therapy while reducing systemic toxicity and side effects. This allows use of drugs that would otherwise be unacceptable when
30 administered systemically. Dosage will depend on the potency of the drug and the efficiency of the carrier construct. Other examples of *in vivo* uses include the use of binding constructs or antigen-binding constructs of the invention in which a toxin is

chemically linked or conjugated to an polypeptide of the invention to form, for example, molecules that may be termed "immunoconjugates" or "immunotoxins." Typically, for example, such a toxin may include one ore more radioisotopes (for example, Iodine-131, Yttrium-90, Rhenium-186, Copper-67, and/or Bishmuth-212),
5 natural toxins, chemotherapy agents, biological response modifiers, or any other substance that is capable of assisting in damaging or killing a target cell, inhibiting target cell replication, or is effective in disrupting a desired cellular function in a target cell.

The toxin portion of the immunotoxin can be derived form various
10 sources. Toxins are commonly derived from plants or bacteria, but toxins of human origin or synthetic toxins can be used as well, for example. Examples of toxins derived from bacteria or plants include, but are not limited to, abrin, α -sarcin, diptheria toxin, ricin, saporin, and *pseudomonas* exotoxin. Examples of mammalian enzymes include, but are not limited to, ribonucleases (RNAse) and
15 deoxyribonucleases. Numerous immunotoxins that may be used with one or more constructs of the invention have been described in the art. See, for example, U.S. Pat. No. 4,753,894 to Frankel *et al.*; U.S. Pat. No. 6,099,842 to Pastan *et al.*; Nevelle, *et al.*, 1982 *Immunol Rev.* **62**: 75-91; Pastan *et al.*, 1992 *Ann Rev Biochem* **61**: 331-354; Chaudary *et al.*, 1989 *Nature* **339**: 394; and Batra *et al.*, 1991 *Mol. Cell. Biol.* **11**:
20 2200. Modified toxins described herein and those described in the various publications are also within the scope of the instant invention.

Generally, the immunotoxins and other therapeutic agents of this invention are administered at a concentration that is therapeutically effective to treat or prevent a particular disease, disorder, or condition, such as for the treatment of
25 tumors and malignancies, the treatment of autoimmune diseases, allergies and inflammation, *etc.* This effective dosage and mode of administration will depend on the animal or patient being treated, the disease or condition being treated, the strength of the immunoconjugates or immunotoxins and the efficiency of the conjugate. To accomplish this goal, the immunotoxins may be formulated using a variety of
30 acceptable formulations and excipients known in the art. Typically, for example, the immunotoxins are administered by injection, either intravenously or intraperitoneally. Methods to accomplish this administration are known to those of ordinary skill in the art. It another aspect, the invention includes topically or orally administered

compositions such as an aerosol or cream or patch that may be capable of transmission across mucous membranes.

Formulants may be added to an immunoconjugates or immunotoxins of the invention before administration to a patients being treated. A liquid formulation is most common, but other formulations are within the scope of the invention. The formulants may include for example oils, polymers, vitamins, carbohydrates, amino acids, salts, buffers, albumin, surfactants, or bulking agents. Carbohydrates can include sugar or sugar alcohols such as mono, di, or polysaccharides, or water-soluble glucans. The saccharides or glucans can include for example fructose, dextrose, lactose, glucose, mannose, sorbose, xylose, maltose, sucrose, dextran, pullulan, dextrin, alpha and beta cyclodextrin, soluble starch, hydroxethyl starch and carboxymethylcellulose, or mixtures thereof. "Sugar alcohol" may be defined as a C₄ to C₈ hydrocarbon having an -OH group and includes, for example, galactitol, inositol, mannitol, xylitol, sorbitol, glycerol, and arabitol. These sugars or sugar alcohols mentioned above may be used individually or in combination. There is no fixed limit to the amount used as long as the sugar or sugar alcohol is soluble in the aqueous preparation. In one aspect, the sugar or sugar alcohol concentration is between 0.5 w/v % and 15 w/v %, typically between 1.0 w/v % and 7.0 w/v %, more typically between 2.0 and 6.0w/v %.

Exemplary amino acids include levorotary (L) forms of camitine, arginine, and betaine; however, other amino acids may be added. Commonly used polymers include polyvinylpyrrolidone (PVP) with an average molecular weight between 2,000 and 3,000, for example, or polyethylene glycol (PEG) with an average molecular weight between 3,000 and 5,000, for example. A buffer can be used in the composition to minimize pH changes in the solution before lyophilization or after reconstitution. Any physiological buffer may be used, but citrate, phosphate, succinate, and glutamate buffers or mixtures thereof are more commonly utilized. The concentration can be, for example, from 0.01 to 0.3 molar. Higher or lower concentrations may be used.

Immunotoxins of the invention can be chemically modified by covalent conjugation to a polymer to increase their circulating half-life, for example. Exemplary polymers and methods to attach them to peptides are referenced in U.S.

Pat. No. 4,766,106 to Katre *et al.*; 4,179,337 to Davis *et al.*; 4,495,285 to Shimizu *et al.*; and U.S. Pat. No. 4,609,546 to Hiratani.

The following Examples are offered by way of illustration and not by way of limitation.

5

EXAMPLES

EXAMPLE 1

CLONING OF THE 2H7 VARIABLE REGIONS AND CONSTRUCTION AND SEQUENCING OF 2H7ScFv-Ig

10 This Example illustrates the cloning of cDNA molecules that encode the heavy chain and light chain variable regions of the monoclonal antibody 2H7. This Example also demonstrates the construction, sequencing, and expression of 2H7scFv-Ig.

Hybridoma cells expressing 2H7 monoclonal antibody that specifically bound to CD20 were provided by Ed Clark at the University of Washington, Seattle, WA. Prior to harvesting, hybridoma cells were kept in log phase growth for several days in RPMI 1640 media Invitrogen/Life Technologies, Gaithersburg, MD) supplemented with glutamine, pyruvate, DMEM non-essential amino acids, and penicillin-streptomycin. Cells were pelleted by centrifugation from the culture medium, and 2×10^7 cells were used to prepare RNA. RNA was isolated from the 2H7-producing hybridoma cells using the Pharmingen (San Diego, CA) total RNA isolation kit (Catalog # 45520K) according to the manufacturer's instructions accompanying the kit. One microgram (1 μ g) of total RNA was used as template to prepare cDNA by reverse transcription. The RNA and 300 ng random primers were combined and denatured at 72° C for 10 minutes prior to addition of enzyme. Superscript II reverse transcriptase (Life Technologies) was added to the RNA plus-primer mixture in a total volume of 25 μ l in the presence of 5x second strand buffer and 0.1 M DTT provided with the enzyme. The reverse transcription reaction was allowed to proceed at 42° C for one hour.

30 The 2H7 cDNA generated in the randomly primed reverse transcriptase reaction and V region specific primers were used to amplify by PCR the variable regions for the light and heavy chain of the 2H7 antibody. The V region specific primers were designed using the published sequence (Genbank accession

numbers M17954 for V_L and M17953 for V_H) as a guide. The two variable chains were designed with compatible end sequences so that an scFv could be assembled by ligation of the two V regions after amplification and restriction enzyme digestion.

A (gly₄ser)₃ peptide linker to be inserted between the two V regions was incorporated by adding the extra nucleotides to the antisense primer for the V_L of 2H7. A Sac I restriction site was also introduced at the junction between the two V regions. The sense primer used to amplify the 2H7 V_L, that included a HindIII restriction site and the light chain leader peptide was 5'-gtc aag ctt gcc gcc atg gat ttt caa gtg cag att ttt cag c-3' (SEQ ID NO: 530). The antisense primer was 5'-gtc gtc gag ctc cca cct cct cca gat cca cca ccg ccc gag cca ccg cca cct ttc agc tcc agc ttg gtc cc-3' (SEQ ID NO: 531). The reading frame of the V region is indicated as a bold, underlined codon. The Hind III and SacI sites are indicated by underlined italicized sequences.

The V_H domain was amplified without a leader peptide, but included a 5' SacI restriction site for fusion to the V_L and a BclI restriction site at the 3' end for fusion to various tails, including the human IgG1 Fc domain and the truncated forms of CD40 ligand, CD154. The sense primer was 5'-gct gct gag ctc tca ggc tta tct aca gca agt ctg g-3' (SEQ ID NO: 532). The SacI site is indicated in italicized and underlined font, and the reading frame of the codon for the first amino acid of the V_H domain is indicated in bold, underlined type. The antisense primer was 5'-gtt gtc tga tca gag acg gtg acc gtg gtc cc-3' (SEQ ID NO: 533). The BclI site is indicated in italicized, underlined type, and the last serine of the V_H domain sequence is indicated in bold, underlined type.

The scFv-Ig was assembled by inserting the 2H7 scFv HindIII-BclI fragment into pUC 19 containing the human IgG1 hinge, CH2, and CH3 regions, which was digested with restriction enzymes, HindIII and BclI. After ligation, the ligation products were transformed into DH5α bacteria. Positive clones were screened for the properly inserted fragments using the SacI site at the V_L-V_H junction of 2H7 as a diagnostic site. The 2H7scFv-Ig cDNA was subjected to cycle sequencing on a PE 9700 thermocycler using a 25-cycle program by denaturing at 96° C for 10 seconds, annealing at 50° C for 30 seconds, and extending at 72° C for 4 minutes. The sequencing primers were pUC forward and reverse primers and an internal primer that annealed to the CH2 domain human in the IgG constant region portion. Sequencing

reactions were performed using the Big Dye Terminator Ready Sequencing Mix (PE-Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. Samples were subsequently purified using Centriseq columns (Catalog # CS-901, Princeton Separations, Adelphia, NJ), the eluates dried in a Savant vacuum dryer, denatured in Template Suppression Reagent (PE-ABI), and analyzed on an ABI 310 Genetic Analyzer (PE-Applied Biosystems). The sequence was edited, translated, and analyzed using Vector Nti version 6.0 (Informax, North Bethesda, MD). Figure 1 shows the cDNA and predicted amino acid sequence of the 2H7scFv-Ig construct.

EXAMPLE 2

10 EXPRESSION OF 2H7 scFv-IG 1N STABLE CHO CELL LINES

This Example illustrates expression of 2H7scFv-Ig in a eukaryotic cell line and characterization of the expressed 2H7scFv-Ig by SDS-PAGE and by functional assays, including ADCC and complement fixation.

The 2H7scFv-Ig HindIII-XbaI (-1.6 kb) fragment with correct sequence was inserted into the mammalian expression vector pD18, and DNA from positive clones was amplified using QIAGEN plasmid preparation kits (QIAGEN, Valencia, CA). The recombinant plasmid DNA (100 µg) was then linearized in a nonessential region by digestion with AscI, purified by phenol extraction, and resuspended in tissue culture media, Excell 302 (Catalog # 14312-79P, JRH Biosciences, Lenexa, KS). Cells for transfection, CHO DG44 cells, were kept in logarithmic growth, and 10⁷ cells harvested for each transfection reaction. Linearized DNA was added to the CHO cells in a total volume of 0.8 ml for electroporation.

Stable production of the 2H7 scFv-Ig fusion protein (SEQ. ID NO: 10) was achieved by electroporation of a selectable, amplifiable plasmid, pD18, containing the 2H7 scFv-Ig cDNA under the control of the CMV promoter, into Chinese Hamster Ovary (CHO) cells (all cell lines from American Type Culture Collection, Manassas, VA, unless otherwise noted). The 2H7 expression cassette was subcloned downstream of the CMV promoter into the vector multiple cloning site as a .about.1.6 kb HindIII-XbaI fragment. The pD18 vector is a modified version of pcDNA3 encoding the DHFR selectable marker with an attenuated promoter to increase selection pressure for the plasmid. Plasmid DNA was prepared using Qiagen maxiprep kits, and purified plasmid was linearized at a unique AscI site prior to

phenol extraction and ethanol precipitation. Salmon sperm DNA (Sigma-Aldrich, St. Louis, MO) was added as carrier DNA, and 1001g each of plasmid and carrier DNA was used to transfect 10^7 CHO DG44 cells by electroporation. Cells were grown to logarithmic phase in Excell 302 media (JRH Biosciences) containing glutamine (4 mM), pyruvate, recombinant insulin, penicillin-streptomycin, and 2x DMEM nonessential amino acids (all from Life Technologies, Gaithersburg, MD), hereafter referred to as "Excell 302 complete" media. Media for untransfected cells also contained HT (diluted from a 100x solution of hypoxanthine and thymidine) (Invitrogen/Life Technologies). Media for transfections under selection contained varying levels of methotrexate (Sigma-Aldrich) as selective agent, ranging from 50 nM to 5 μ M. Electroporations were performed at 275 volts, 950 μ F. Transfected cells were allowed to recover overnight in non-selective media prior to selective plating in 96 well flat bottom plates (Costar) at varying serial dilutions ranging from 125 cells/well to 2000 cells/well. Culture media for cell cloning was Excell 302 complete, containing 100 nM methotrexate. Once clonal outgrowth was sufficient, serial dilutions of culture supernatants from master wells were screened for binding to CD20-CHO transfected cells. The clones with the highest production of the fusion protein were expanded into T25 and then T75 flasks to provide adequate numbers of cells for freezing and for scaling up production of the 2H7scFvIg. Production levels were further increased in cultures from three clones by progressive amplification in methotrexate containing culture media. At each successive passage of cells, the Excell 302 complete media contained an increased concentration of methotrexate, such that only the cells that amplified the DHFR plasmid could survive.

Supernatants were collected from CHO cells expressing the 2H7scFv-Ig, filtered through 0.2 μ m PES express filters (Nalgene, Rochester, NY) and were passed over a Protein A-agarose (IPA 300 crosslinked agarose) column (Repligen, Needham, MA). The column was washed with PBS, and then bound protein was eluted using 0.1 M citrate buffer, pH 3.0. Fractions were collected and eluted protein was neutralized using 1M Tris, pH 8.0, prior to dialysis overnight in PBS. Concentration of the purified 2H7scFv-Ig was determined by absorption at 280 nm. An extinction coefficient of 1.77 was determined using the protein analysis tools in the Vector Nti Version 6.0 Software package (Informax, North Bethesda, MD). This program uses the amino acid composition data to calculate extinction coefficients.

Production levels of 2H7scFv-Ig by transfected, stable CHO cells were analyzed by flow cytometry. Purified 2H7scFv-Ig to CHO cells was allowed to bind to CHO cells that expressed CD20 (CD20 CHO) and analyzed by flow cytometry using a fluorescein-conjugated anti-human IgG second step reagent (Catalog Numbers
5 H10101 and H10501, CalTag, Burlingame, CA). Figure 2 (top) shows a standard curve generated by titration of 2H7scFv-Ig binding to CD20 CHO. At each concentration of 2H7scFv-Ig, the mean brightness of the fluorescein signal in linear units is shown. Supernatants collected from T flasks containing stable CHO cell clones expressing 2H7scFv-1 g were then allowed to bind to CD20 CHO and the
10 binding was analyzed by flow cytometry. The fluorescein signal generated by 2H7scFv-Ig contained in the supernatants was measured and the 2H7scFv-Ig concentration in the supernatants was calculated from the standard curve (Figure 2, bottom).

Purified 2H7scFv-Ig was analyzed by electrophoresis on SDS-
15 Polyacrylamide gels. Samples of 2H7scFv-Ig, purified by independent Protein A Agarose column runs, were boiled in SDS sample buffer without reduction of disulfide bonds and applied to SDS 10% Tris-BIS gels (Catalog # NP0301, Novex, Carlsbad, CA). Twenty micrograms of each purified batch was loaded on the gels. The proteins were visualized after electrophoresis by Coomassie Blue staining (Pierce
20 Gel Code Blue Stain Reagent, Catalog #24590, Pierce, Rockford, IL), and destaining in distilled water. Molecular weight markers were included on the same gel (Kaleidoscope Prestained Standards, Catalog # 161-0324, Bio-Rad, Hercules, CA). The results are presented in Figure 3. The numbers above the lanes designate independent purification batches. The molecular weights in kilodaltons of the size
25 markers are indicated on the left side of the figure. Further experiments with alternative sample preparation conditions indicated that reduction of disulfide bonds by boiling the protein in SDS sample buffer containing DTT or 2-mercaptoethanol caused the 2H7scFv-Ig to aggregate.

Any number of other immunological parameters may be monitored
30 using routine assays that are well known in the art. These may include, for example, antibody dependent cell-mediated cytotoxicity (ADCC) assays, secondary *in vitro* antibody responses, flow immunocytofluorimetric analysis of various peripheral blood or lymphoid mononuclear cell subpopulations using well established marker

antigen systems, immunohistochemistry or other relevant assays. These and other assays may be found, for example, in Rose *et al.* (Eds.), *Manual of Clinical Laboratory Immunology*, 5th Ed., 1997 American Society of Microbiology, Washington, DC.

5 The ability of 2H7scFv-Ig to kill CD20 positive cells in the presence of complement was tested using B cell lines Ramos and Bjab. Rabbit complement (Pel-Freez, Rogers, AZ) was used in the assay at a final dilution of {fraction (1/10)}. Purified 2H7scFv-Ig was incubated with B cells and complement for 45 minutes at 37° C, followed by counting of live and dead cells by trypan blue exclusion. The
10 results in Figure 4A show that in the presence of rabbit complement, 2H7scFv-Ig lysed B cells expressing CD20.

 The ability of 2H7scFv-Ig to kill CD20 positive cells in the presence of peripheral blood mononuclear cells (PBMC) was tested by measuring the release of ⁵¹Cr from labeled Bjab cells in a 4-hour assay using a 100:1 ratio of PBMC to Bjab
15 cells. The results shown in Figure 4B indicated that 2H7scFv-Ig can mediate antibody dependent cellular cytotoxicity (ADCC) because the release of ⁵¹Cr was higher in the presence of both PBMC and 2H7scFv-Ig than in the presence of either PBMC or 2H7scFv-Ig alone.

EXAMPLE 3

20 **EFFECT OF SIMULTANEOUS LIGATION OF CD20 AND CD40 ON GROWTH OF NORMAL B CELLS, AND ON CD95 EXPRESSION, AND INDUCTION OF APOPTOSIS**

 This example illustrates the effect of cross-linking of CD20 and CD40 expressed on the cell surface on cell proliferation.

 Dense resting B cells were isolated from human tonsil by a Percoll step
25 gradient and T cells were removed by E-rosetting. Proliferation of resting, dense tonsillar B cells was measured by uptake of ³[H]-thymidine during the last 12 hours of a 4-day experiment. Proliferation was measured in quadruplicate cultures with means and standard deviations as shown. Murine anti-human CD20 monoclonal antibody 1F5 (anti-CD20) was used alone or was cross-linked with anti-murine κ monoclonal
30 antibody 187.1 (anti-CD20XL). CD40 activation was accomplished using soluble human CD154 fused with murine CD8 (CD154) (Hollenbaugh *et al.*, *EMBO J.* 11: 4212-21 (1992)), and CD40 cross-linking was accomplished using anti-murine CD8

monoclonal antibody 53-6 (CD154XL). This procedure allowed simultaneous cross-linking of CD20 and CD40 on the cell surface. The results are presented in Figure 5.

The effect of CD20 and CD40 cross-linking on Ramos cells, a B lymphoma cell line, was examined. Ramos cells were analyzed for CD95 (Fas) expression and percent apoptosis eighteen hours after treatment (no goat anti-mouse IgG (GAM)) and/or cross-linking (+GAM) using murine monoclonal antibodies that specifically bind CD20 (1F5) and CD40 (G28-5). Control cells were treated with a non-binding isotype control (64.1) specific for CD3.

Treated Ramos cells were harvested, incubated with FITC-anti-CD95, and analyzed by flow cytometry to determine the relative expression level of Fas on the cell surface after CD20 or CD40 cross-linking. Data is plotted as mean fluorescence of cells after treatment with the stimuli indicated (Figure 6A).

Treated Ramos cells from the same experiment were harvested and binding of annexin V was measured to indicate the percentage apoptosis in the treated cultures. Apoptosis was measured by binding of Annexin V 18 hours after cross-linking of CD20 and CD40 using 1F5 and G28-5 followed by cross-linking with GAM. Binding of Annexin V was measured using a FITC-Annexin V kit (Catalog # PN-IM2376, Immunotech, Marseille, France,). Annexin V binding is known to be an early event in progression of cells into apoptosis. Apoptosis, or, programmed cell death, is a process characterized by a cascade of catabolic reactions leading to cell death by suicide. In the early phase of apoptosis, before cells change morphology and hydrolyze DNA, the integrity of the cell membrane is maintained but cells lose the asymmetry of their membrane phospholipids, exposing negatively charged phospholipids, such as phosphatidylserine, at the cell surface. Annexin V, a calcium and phospholipid binding protein, binds preferentially and with high affinity to phosphatidylserine. Results demonstrating the effect of cross-linking both CD20 and CD40 on expression of the FAS receptor (CD95) are presented in Figure 6B. The effect of cross-linking of both CD20 and CD40 on Annexin V binding to cells is shown in Figure 6B.

EXAMPLE 4

CONSTRUCTION AND CHARACTERIZATION OF 2H7 scFv-CD154 FUSION PROTEINS

To construct a molecule capable of binding to both CD20 and CD40, cDNA encoding the 2H7 scFv was fused with cDNA encoding CD154, the CD40 ligand. The 2H7 scFv cDNA encoded on the HindIII-BclI fragment was removed from the 2H7 scFvIg construct, and inserted into a pD18 vector along with a BamHI-XbaI cDNA fragment encoding the extracellular domain of human CD154. The extracellular domain is encoded at the carboxy terminus of CD154, similar to other type II membrane proteins.

The extracellular domain of human CD154 was PCR amplified using cDNA generated with random primers and RNA from human T lymphocytes activated with PHA (phytohemagglutinin). The primer sets included two different 5' or sense primers that created fusion junctions at two different positions within the extracellular domain of CD 154. Two different fusion junctions were designed that resulted in a short or truncated form (form S4) including amino acids 108 (Glu)-261 (Leu)+(Glu), and a long or complete form (form L2) including amino acids 48 (Arg)-261 (Leu)+(Glu), of the extracellular domain of CD154, both constructed as BamHI-XbaI fragments. The sense primer which fuses the two different truncated extracellular domains to the 2H7scFv includes a BamHI site for cloning. The sense primer for the S4 form of the CD154 cDNA is designated SEQUENCE ID NO: 535 or CD154BAM108 and encodes a 34 mer with the following sequence: 5'-gtt gtc gga tcc aga aaa cag ctt tga aat gca a-3', while the antisense primer is designated SEQUENCE ID NO: 536 or CD154XBA and encodes a 44 mer with the following sequence: 5'-gtt gtt tct aga tta tca ctc gag ttt gag taa gcc aaa gga cg-3'.

The oligonucleotide primers used in amplifying the long form (L2) of the CD154 extracellular domain encoding amino acids 48 (Arg)-261 (Leu)+(Glu), were as follows: The sense primer designated CD154 BAM48 (SEQUENCE ID NO: 537) encoded a 35-mer with the following sequence: 5'-gtt gtc gga tcc aag aag gtt gga caa gat aga ag-3'. The antisense primer designated or CD154XBA (SEQUENCE ID NO: 538) encoded the 44-mer: 5'-gtt gtt tct aga tta tca ctc gag ttt gag taa gcc aaa gga cg-3'. Other PCR reaction conditions were identical to those used for amplifying the 2H7 scFv (see Example 1). PCR fragments were purified by PCR quick kits

(QIAGEN, San Diego, CA), eluted in 30 µl ddH₂O, and digested with BamHI and XbaI (Roche) restriction endonucleases in a 40 µl reaction volume at 37° C for 3 hours. Fragments were gel purified, purified using QIAEX kits according to the manufacturer's instructions (QIAGEN), and ligated along with the 2H7 HindIII-BclI
5 fragment into the pD 18 expression vector digested with HindIII+XbaI. Ligation reactions were transformed into DH5-alpha chemically competent bacteria and plated onto LB plates containing 100 µg/ml ampicillin. Transformants were grown overnight at 37° C, and isolated colonies used to inoculate 3 ml liquid cultures in Luria Broth containing 100 µg/ml ampicillin. Clones were screened after mini-
10 plasmid preparations (QIAGEN) for insertion of both the 2H7 scFv and the CD154 extracellular domain fragments.

The 2H7scFv-CD154 construct cDNAs were subjected to cycle sequencing on a PE 9700 thermocycler using a 25-cycle program that included denaturing at 96° C, 10 seconds, annealing at 50° C for 5 seconds, and extension at
15 60° C, for 4 minutes. The sequencing primers used were pD18 forward (SEQ ID NO: 539 5'-gtctatataagcagagctctggc-3') and pD18 reverse (SEQ ID NO: 540 5'-cgaggctgatcagcgagctctagca-3') primers. In addition, an internal primer was used that had homology to the human CD154 sequence (SEQ ID NO: 541 5'-ccgcaatttgaggattctgacacc-3'). Sequencing reactions included primers at 3.2 pmol,
20 approximately 200 ng DNA template, and 8 µl sequencing mix. Sequencing reactions were performed using the BIGDYE® Terminator Ready Sequencing Mix (PE-Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. Samples were subsequently purified using Centriscp columns (Princeton Separations, Adelphia, NJ). The eluates were dried in a Savant speed-vacuum dryer, denatured in
25 20 µl template Suppression Reagent (ABI) at 95° C for 2 minutes, and analyzed on an ABI 310 Genetic Analyzer (PE-Applied Biosystems). The sequence was edited, translated, and analyzed using Vector Nti version 6.0 (Informax, North Bethesda, MD). The 2H7scFv-CD154 L2 cDNA sequence and predicted amino acid sequence is presented in Figure 7A, and 2H7scFv-CD154 S4 cDNA sequence and predicted
30 amino acid sequence is presented in Figure 7B.

The binding activity of the 2H7 scFv-CD154 fusion proteins (SEQ. ID NO: 691 and 693) to CD20 and CD40 simultaneously was determined by flow cytometry. The assay used CHO cell targets that express CD20. After a 45-minute

incubation of CD20 CHO cells with supernatants from cells transfected with the 2H7 scFv-CD154 expression plasmid, the CD20 CHO cells were washed twice and incubated with biotin-conjugated CD40-Ig fusion protein in PBS/2% FBS. After 45 min, cells were washed twice and incubated with phycoerythrin (PE)-labeled streptavidin at 1:100 in PBS/2% FBS (Molecular Probes, Eugene OR). After an additional 30 min incubation, cells were washed 2x and were analyzed by flow cytometry. The results show that the 2H7 scFv-CD154 molecule was able to bind to CD20 on the cell surface and to capture biotin-conjugated CD40 from solution (Figure 8).

To determine the effect of the 2H7scFv-CD154 on growth and viability of B lymphoma and lymphoblastoid cell lines, cells were incubated with 2H7scFv-CD154 L2 (SEQ ID NO: 691) for 12 hours and then examined for binding of Annexin V. Binding of Annexin V was measured using a FITC-Annexin V kit (Immunotech, Marseille, France, Catalog # PN-IM2376). B cell lines were incubated in 1 ml cultures with dilutions of concentrated, dialyzed supernatants from cells expressing secreted forms of the 2H7scFv-CD154 fusion proteins. The results are presented in Figure 9.

The growth rate of the Ramos B lymphoma cell line in the presence of 2H7scFv-CD 154 was examined by uptake of ³H-thymidine for the last 6 hours of a 24-hour culture. The effect of 2H7scFv-CD154 on cell proliferation is shown in Figure 10.

EXAMPLE 5

CONSTRUCTION AND CHARACTERIZATION OF CYTOXB ANTIBODY DERIVATIVES

CytoxB antibodies were derived from the 2H7 scFv-IgG polypeptide. The 2H7 scFv (see Example 1) was linked to the human IgG1 Fc domain via an altered hinge domain (see Figure 11). Cysteine residues in the hinge region were substituted with serine residues by site-directed mutagenesis and other methods known in the art. The mutant hinge was fused either to a wild-type Fc domain to create one construct, designated CytoxB-MHWTG1C, or was fused to a mutated Fc domain (CytoxB-MHMG1C) that had additional mutations introduced into the CH2 domain. Amino acid residues in CH2 that are implicated in effector function are illustrated in Figure 11. Mutations of one or more of these residues may reduce FcR binding and mediation of effector functions. In this example, the leucine residue 234

known in the art to be important to Fc receptor binding, was mutated in the 2H7 scFv fusion protein, CytoxB-[MG1H/MG1C]. In another construct, the human IgG1 hinge region was substituted with a portion of the human IgA hinge, which was fused to wild-type human Fc domain (CytoxB-IgAHWTHGI C). (See Figure 11). This mutated hinge region allows expression of a mixture of monomeric and dimeric molecules that retain functional properties of the human IgG1 CH2 and CH3 domains. Synthetic, recombinant cDNA expression cassettes for these molecules were constructed and polypeptides were expressed in CHODG44 cells according to methods described in Example 2.

Purified fusion protein derivatives of CytoxB-scFvIg molecules were analyzed by SDS-PAGE according to the methods described in Example 2. Polyacrylamide gels were run under non-reducing and reducing conditions. Two different molecule weight marker sets, BioRad prestained markers, (BioRad, Hercules, CA) and Novex MULTIMARK® molecular weight markers were loaded onto each gel. The migration patterns of the different constructs and of Rituximab are presented in Figure 12.

The ability of the different derivatives of CytoxB-scFvIg molecules to mediate ADCC was measured using the Bjab B lymphoma cells as the target and freshly prepared human PBMCs as effector cells. (See Example 2). Effector to target ratios were varied as follows: 70:1, 35:1, and 18:1, with the number of Bjab cells per well remaining constant but the number of PBMCs were varied. Bjab cells were labeled for 2 hours with ^{51}Cr and aliquoted at a cell density of 5×10^4 cells/well to each well of flat-bottom 96 well plates. Purified fusion proteins or rituximab were added at a concentration of 10 $\mu\text{g/ml}$ to the various dilutions of PBMCs. Spontaneous release was measured without addition of PBMC or fusion protein, and maximal release was measured by the addition of detergent (1% NP-40) to the appropriate wells. Reactions were incubated for 4 hours, and 100 μl of culture supernatant was harvested to a Lumaplate (Packard Instruments) and allowed to dry overnight prior to counting cpm released. The results are presented in Figure 13.

Complement dependent cytotoxicity (CDC) activity of the CytoxB derivatives was also measured. Reactions were performed essentially as described in Example 2. The results are presented in Figure 14 as percent of dead cells to total cells for each concentration of fusion protein.

EXAMPLE 6

IN VIVO STUDIES IN MACAQUES

Initial *in vivo* studies with CytoxB derivatives have been performed in nonhuman primates. Figure 15 shows data characterizing the serum half-life of
5 CytoxB in monkeys. Measurements were performed on serum samples obtained from two different macaques (J99231 and K99334) after doses of 6 mg/kg were administered to each monkey on the days indicated by arrows. For each sample, the level of 2H7scFvIg present was estimated by comparison to a standard curve generated by binding of purified CytoxB-(MHWTG1C)-1 g fusion protein to CD20
10 CHO cells (see Example 2). The data are tabulated in the bottom panel of the Figure 15.

The effect of CytoxB-(MHWTG1 C)Ig fusion protein on levels of circulating CD40+ cells in macaques was investigated. Complete blood counts were performed at each of the days indicated in Figure 16. In addition, FACS (fluorescence
15 activated cell sorter) assays were performed on peripheral blood lymphocytes using a CD40-specific fluorescein conjugated antibody to detect B cells among the cell population. The percentage of positive cells was then used to calculate the number of B cells in the original samples. The data are graphed as thousands of B cells per microliter of blood measured at the days indicated after injection (Figure 16).

20 EXAMPLE 7

CONSTRUCTION AND EXPRESSION OF AN ANTI-CD19 scFv-IG FUSION PROTEIN

An anti-CD 19 scFv-Ig fusion protein was constructed, transfected into eukaryotic cells, and expressed according to methods presented in Examples 1, 2, and 5 and standard in the art. The variable heavy chain regions and variable light chain
25 regions were cloned from RNA isolated from hybridoma cells producing antibody HD37, which specifically binds to CD19. Expression levels of a HD37scFv-IgAHWTG1C and a HD37scFv-IgMHWTG1C were measured and compared to a standard curve generated using purified HD37 scFvIg. The results are presented in Figure 17.

EXAMPLE 8

CONSTRUCTION AND EXPRESSION OF AN ANTI-L6 scFv-IG FUSION PROTEIN

An scFv-Ig fusion protein was constructed using variable regions derived from an anti-carcinoma monoclonal antibody, L6. The fusion protein was constructed, transfected into eukaryotic cells, and expressed according to methods presented in Examples 1, 2, and 5 and standard in the art. Expression levels of L6 scFv-IgAH WCH2 CH3 and L6 scFv-(SSS-S)H WCH2 WCH3 were measured and compared to a standard curve generated using purified L6 scFvIg. The results are presented in Figure 18.

EXAMPLE 9

CHARACTERIZATION OF VARIOUS scFv-IG FUSION PROTEINS

In addition to the scFv-Ig fusion protein already described, G28-1 (anti-CD37) scFv-Ig fusion proteins were prepared essentially as described in Examples 1 and 5. The variable regions of the heavy and light chains were cloned according to methods known in the art. ADCC activity of 2H7-MHWTG1C, 2H7-IgAHWTG1C, G28-1-MHWTG1C, G28-1 IgAHWTG1C, HD37-MHWTG1C, and HD37--IgAHWTG1C was determined according to methods described above (see Example 2). Results are presented in Figure 19. ADCC activity of L6scFv-IgAHWTG1C and L6scFv-IgMHWTG1C was measured using the 2981 human lung carcinoma cell line. The results are presented in Figure 20. The murine L6 monoclonal antibody is known not to exhibit ADCC activity.

The purified proteins were analyzed by SDS-PAGE under reducing and non-reducing conditions. Samples were prepared and gels run essentially as described in Examples 2 and 5. The results for the L6 and 2H7 scFv-Ig fusion proteins are presented in Figure 21 and the results for the G28-1 and HD37 scFv-Ig fusion proteins are presented in Figure 22.

EXAMPLE 10

CONSTRUCTION AND EXPRESSION OF ANTI-CD20 scFv-LLAMA Ig FUSION

PROTEINS

This Example illustrates the cloning of llama IgG1, IgG2, and IgG3
5 constant region domains and the construction of immunoglobulin fusion proteins with
each of the three constant regions and anti-CD20 scFv.

The constant regions of llama IgG1, IgG2, and IgG3 immunoglobulins
were cloned and inserted into mammalian vector constructs containing an anti-CD20
single chain Fv, 2H7 scFv. Total RNA was isolated from peripheral blood
10 mononuclear cells (PBMC) from llama blood (Triple J Farms, Bellingham, WA) by
lysing the lymphocytes in TRIZOL® (Invitrogen Life Technologies, Carlsbad, CA)
according to the manufacturer's instructions. One microgram (1 µg) of total RNA was
used as template to prepare cDNA by reverse transcription. The RNA and 200 ng
random primers were combined and denatured at 72° C for 10 minutes prior to
15 addition of enzyme. Superscript II reverse transcriptase (Invitrogen Life
Technologies) was added to the RNA plus primer mixture in a total volume of 25 µl
in the presence of 5x second strand buffer and 0.1 M DTT provided with the enzyme.
The reverse transcription reaction was allowed to proceed at 42° C for one hour. The
cDNA was amplified by PCR using sequence specific primers. The 5' primers were
20 designed according to published sequences for the V_HH and V_H domains of camelids.
The 3' primer, which was used to amplify all three isotypes, was designed using
mammalian CH3 domain sequences as a guide. The following specific primers were
used. The Bcl and XbaI sites are indicated by underlined italicized sequences.

5' primer for llama IgG1 constant region

25 LLG1-5'bgl: 5'-gtt gtg gat caa gaa cca cat gga gga tgc acg tg-3'
(SEQ ID NO: 542)

5' primer for llama IgG2 constant region

LLG2-5'bgl: 5'-gtt gtg gat caa gaa ccc aag aca cca aaa cc-3'
(SEQ ID NO: 543)

30 5' primer for llama IgG3 constant region

LLG3-5'bgl: 5'-gtt gtg gat caa gcg cac cac agc gaa gac ccc-3'
(SEQ ID NO: 544)

3' primer for llama IgG1, IgG2, and IgG3 constant regions

LLG123-3'X: 5'-gtt gtt tct aga tta cta ttt acc cga aga ctg ggt gat gga-3'
(SEQ ID NO: 545)

5 PCR fragments of the expected size were cloned into TOPO® cloning
vectors (Invitrogen Life Technologies) and then were sequenced. The sense
sequencing primer, LLseqsense, had the sequence 5'-ctg aga tcg agt tca gct g-3' (SEQ
ID NO: 546), and the antisense primer, LLseqAS, had the sequence 5'-cct cct ttg gct
ttg tct c-3' (SEQ ID NO: 547). Sequencing was performed as described in Example 1.
Figure 23 compares the amino acid sequence of the three isotype llama constant
10 regions containing the hinge, CH2, and CH3 domains with the amino acid sequence
of human IgG1 hinge, CH2, and CH3 domains.

After verifying the sequence, the amplified PCR products were
digested with restriction enzymes BclI and XbaI to create compatible restriction sites.
The digested fragments were then gel-purified, and the DNA was eluted using a
15 QIAquick Gel Extraction Kit (QIAGEN, Valencia, CA). The 2H7scFv-Ig pD18
mammalian expression vector construct (see Example 2) was digested with BclI and
XbaI to remove the human IgG hinge, CH2, and CH3 domains. The pD18 vector is a
modified derivative of pcDNA3 that contains an attenuated DHFR gene, which serves
as a selectable marker for mammalian expression (Hayden *et al.*, *Tissue Antigens* 48:
20 242-54 (1996)). The purified llama IgG1, IgG2, and IgG3 constant region PCR
products were ligated by T4 DNA ligase (Roche Molecular Biochemicals,
Indianapolis, IN) into the double-digested 2H7 scFv-pD18 vector at room temperature
overnight according to the manufacturer's instructions. After ligation, the ligation
products were transformed into *E. coli* DH5α bacteria (BD Biosciences, Palo Alto,
25 CA) and plated according to standard molecular biology procedures and
manufacturer's instructions. Isolated colonies were chosen to screen for transformants
containing the correct inserts.

For expression of the encoded polypeptides, plasmid DNA from
positive clones was transiently transfected into COS-7 cells using DEAE-dextran
30 (Hayden *et al.*, *The Immunol.* 1: 3-15 (1994)). COS-7 cells were seeded at
approximately 3x 10⁵ cells per 150 mm plate and grown overnight until the cells were
about 75% confluent. Cells were then washed once with serum-free DMEM
(Invitrogen Life Technologies, Grand Island, NY). Transfection supernatant (10 ml)

containing 400 µg/ml DEAE-dextran, 0.1 mM chloroquine, and 5 µg/ml of the DNA constructs were added to the cells, which were then incubated at 37° C for 3-4 hrs. After incubation, cells were pulsed with 10 ml of 10% dimethyl sulfoxide (DMSO) in 1x PBS at room temperature for 2 minutes. Cells were then placed back into fully
5 supplemented DMEM/10% FBS (1% L-glutamine, 1% penicillin/streptomycin, 1% sodium pyruvate, 1% MEM essential amino acids) (Invitrogen Life Technologies). After 24 hours, the media was replaced with serum-free fully supplemented DMEM (Invitrogen Life Technologies), and the cells were maintained up to 21 days with media changes every 3-4 days.

10 Ig-fusion proteins were purified by passing COS cell culture supernatants through Protein A Agarose (Repligen, Cambridge, MA) columns. After application of the culture supernatant, the Protein A columns were then washed with 1x PBS (Invitrogen Life Technologies). Bound Ig-fusion proteins were eluted with 0.1 M citric acid (pH 2.8), and the collected fractions were immediately neutralized
15 with Tris base (pH 10.85). The fractions containing protein were identified by measuring the optical density (A₂₈₀) and then were pooled, dialyzed against 1x PBS, (Invitrogen Life Technologies) and filtered through a 0.2 µm filter.

The purified Ig-fusion proteins were analyzed by SDS-PAGE. Aliquots of 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, 2H7 scFv-llama IgG3, and
20 RITUXAN® (Rituximab, anti-CD20 antibody, Genentech, Inc. and IDEC Pharmaceuticals Corp.) (provided by Dr. Oliver W. Press, Fred Hutchison Cancer Research Center, Seattle, WA) (5 µg protein) were combined with 25 µl 2x NUPAGE® SDS Sample Buffer (Invitrogen Life Technologies) (non-reduced samples). Samples of each protein were also prepared in reducing sample buffer
25 containing 5% 2-mercaptoethanol (Sigma-Aldrich, St. Louis, MO). Molecular weight markers (Invitrogen Life Technologies) were applied to the gels in non-reducing buffer only. The proteins were fractionated on NUPAGE® 10% Bis-Tris gels (Invitrogen Life Technologies). After electrophoresis (approximately 1 hour), the gels were washed three times, five minutes each, with Distilled Water (Invitrogen Life
30 Technologies) and then stained in 50 ml Bio-Safe Coomassie Stain (BioRad, Hercules, CA) overnight at room temperature. After a wash in Distilled Water, the gels were photographed. The migration pattern of each Ig-fusion protein is presented in Figure 24.

The ability of the 2H7 scFv-llama Ig fusion proteins to bind to cells expressing CD20 was demonstrated by flow cytometry. Serial dilutions starting at 25 µg/ml of purified 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, and 2H7 scFv-llama IgG3 were prepared and incubated with CD20-transfected (CD20+) CHO cells (from the laboratory of Dr. S. Skov, Institute of Medical Microbiology and Immunology, Copenhagen Denmark in 1% FBS 1x PBS media (Invitrogen Life Technologies) for one hour on ice. After the incubation, the cells were then centrifuged and washed with 1% FBS in 1x PBS. To detect bound 2H7 scFv-llama Ig, the cells were incubated for one hour on ice with fluorescein-conjugated goat anti-camelid IgG (heavy and light chain) (1:100) (Triple J Farms). The cells were then centrifuged and resuspended in 1% FBS-1x PBS and analyzed using a Coulter Epics XL cell sorter (Beckman Coulter, Miami, FL). The data (percent of maximum brightness) are presented in Figure 25.

EXAMPLE 11

EFFECTOR FUNCTION OF ANTI-CD20 scFV-LLAMA IG FUSION PROTEINS

This Example demonstrates the ability of anti-CD20 llama IgG1, IgG2, and IgG3 fusion proteins to mediate complement dependent cytotoxicity (CDC) and antibody dependent cell-mediated cytotoxicity (ADCC).

The ability of the 2H7 scFv-llama IgG fusion proteins to kill CD20 positive cells in the presence of complement was tested using the BJAB human B cell line. Rabbit complement was obtained from 3-4 week old rabbits (Pel-Freez, Brown Deer, WI). BJAB cells (2×10^6 cells/ml) were combined with rabbit complement (final dilution 1:10) and purified 2H7 Ig fusion proteins. 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, 2H7 scFv-llama IgG3, and 2H7 scFv-human IgG1 wild type hinge-CH2-CH3) (Example 1) were added at 1:3 serial dilutions beginning at a concentration of 30 µg/ml. After one hour at 37° C, cell viability was determined by counting live and dead cells by trypan blue exclusion (0.4%) (Invitrogen Life Technologies) using a hemacytometer (Bright-line, Horsham, PA). The percent killing was calculated by dividing the number of dead cells by the number of total cells (dead+live cells). The data presented in Figure 26 show that all Ig fusion proteins had CDC activity.

The ADCC activity of the 2H7 scFv-llama IgG fusion proteins was determined using BJAB cells as target cells and human or llama peripheral blood mononuclear cells (PBMC) as effector cells. BJAB cells were pre-incubated for approximately 2 hours with ^{51}Cr (100 μCi) (Amersham Biosciences, Piscataway, NJ) in fully supplemented IMDM (Invitrogen Life Technologies) containing 15% FBS. The cells were mixed intermittently during the pre-incubation period. Fresh, resting human PBMC were purified from whole blood using Lymphocyte Separation Media (LSM) (ICN Pharmaceuticals, New York, NY). PBMC were combined with labeled BJAB cells (5×10^4 cells per well of 96 well tissue culture plate) at ratios of 25:1, 50:1, and 100:1. To each combination was added 10 $\mu\text{g/ml}$ of purified 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, 2H7 scFv-llama IgG3, Rituximab, or no anti-CD20 antibody. The mixtures were incubated for 6 hours at 37°C . Supernatant from each reaction containing ^{51}Cr released from lysed cells was collected onto a LumaPlate-96 filter plate (Packard, Meriden, CT), which was dried overnight. The amount of ^{51}Cr was measured by a TopCount NXT plate reader (Packard). Figure 27 shows that the 2H7 scFv-llama IgG2 fusion protein was the most effective llama fusion protein in mediating ADCC. Each data point represents the average measurement of triplicate wells.

ADCC activity was affected by the source of effector cells. Llama PBMC were isolated from llama blood (Triple J Farms) using LSM. Llama effector cells were added at the same ratios to BJAB target cells as described for the ADCC assay using human effector cells. The cells were combined with 10 $\mu\text{g/ml}$ of purified 2H7 scFv-llama IgG1, 2H7 scFv-llama IgG2, 2H7 scFv-llama IgG3, Rituximab, or no anti-CD20 antibody. The results are presented in Figure 28.

EXAMPLE 12

CONSTRUCTION AND CHARACTERIZATION OF scFv Ig FUSION PROTEINS EXPRESSED ON THE CELL SURFACE

This Example describes a retroviral transfection system for ectopic surface expression of genetically engineered cell surface receptors composed of scFvs that bind costimulatory receptors. The Example also demonstrates the effector function of these various scFv Ig fusion proteins expressed on the surface of target cells.

The heavy and light chain variable regions were cloned from murine monoclonal antibodies specific for various costimulatory receptors, and single chain Fv constructs were prepared essentially as described in Example 1. Antibodies included 2H7, anti-human CD20; 40.2.220, anti-human CD40; 2E12, anti-human CD28; 10A8, anti-human CD152 (anti-CTLA-4); and 500A2, anti-murine CD3. The heavy chain and light chain variable regions of each antibody were cloned according to standard methods for cloning immunoglobulin genes and as described in Example 1. Single chain Fv constructs were prepared as described in Example 1 by inserting a nucleotide sequence encoding a (gly₄ser)₃ peptide linker between the V_L region nucleotide sequence of 40.2.220, 2E12, 10A8, and 500A2, respectively (SEQ ID NOs: 31, 39, 43, respectively) and the V_H region nucleotide sequence of 40.2.220, 2E12, 10A8, and 500A2, respectively (SEQ ID NOs: 33, 39, 45, respectively). The polypeptide sequence for V_L of 40.2.220, 2E12, 10A8, and 500A2 are set forth in SEQ ID NOs: 32, 38, 44, respectively, and the polypeptide sequence for V_H of 40.2.220, 2E12, 10A8, and 500A2 are set forth in SEQ ID NOs: 34, 40, 46, respectively. Each scFv polynucleotide (SEQ ID NOs: 36, 42, 47 for 40.2.220, 2E12, 10A8, and 500A2, respectively) was then fused to human IgG1 mutant hinge (CCC→SSS) and mutant CH2 (proline to serine mutation at residue 238 (238 numbering according to EU nomenclature, Ward *et al.*, 1995 *Therap. Immunol.* 2:77-94; residue 251 according to Kabat *et al.*) and wild type CH3 domains according to the methods described in Example 5 and 11. Each scFv mutant IgG1 fusion polynucleotide sequence was then fused in frame to sequences encoding the transmembrane domain and cytoplasmic tail of human CD80 (SEQ ID NO: 29), such that when the fusion protein was expressed in the transfected cell, CD80 provided an anchor for surface expression of the scFv Ig fusion protein. cDNAs encoding the scFv-IgG-CD80 fusion proteins (SEQ ID NOs: 49, 51, 53 and 55, for 40.2.220-, 2E12-, 10A8-, and 500A2-scFv-IgG-CD80, respectively) were inserted into the retroviral vector pLNCX (BD Biosciences Clontech, Palo Alto, CA) according to standard molecular biology procedures and vendor instructions. The scFv-Ig-CD80 cDNA was inserted between the 5'LTR-neomycin resistance gene-CMV promoter sequences and the 3'LTR sequence. The retroviral constructs were transfected into Reh, an acute lymphocytic leukemia cell line (ATCC CRL-8286). Transfected cells were screened to select clones that were expressing scFv-Ig fusion proteins on the cell surface.

CDC and ADCC assays were performed with the transfected Reh cells to determine if expression of the scFv-Ig polypeptides on the cell surface augmented effector cell function. Reh cells expressing anti-human CD152 scFv-mutant IgG-CD80 (SEQ ID NO: 56); Reh anti-human CD28 scFv-mutant IgG-CD80 (SEQ ID NO: 52); Reh anti-human CD40 scFv-mutant IgG-CD80 (SEQ ID NO: 50); Reh anti-human CD20 scFv-mutant IgG-CD80 were combined with human PBMC (see Example 11) and rabbit complement (10 µg/ml) for one hour at 37° C. Untransfected Reh cells were included as a control. Viability of the cells was determined by trypan blue exclusion, and the percent of killed cells was calculated (see Example 11). Figure 29 shows the effectiveness of the scFv-IgG-CD80 fusion proteins when expressed on the cell surface of tumor cells to mediate complement dependent cytotoxicity.

The same transfected Reh cells tested in the CDC assay plus Reh cells transfected with the polynucleotide construct that encodes anti-murine CD3-scFv-Ig-CD80 (SEQ ID NO: 110) were analyzed for ADCC activity (see Example 11). Untransfected and transfected Reh cells were pre-labeled with ⁵¹Cr (100 µCi) (Amersham) for two hours at 37° C. Human PBMC served as effector cells and were added to the Reh target cells (5 x 10⁴ cells per well of 96 well plate) at ratios of 5:1, 2.5:1, and 1.25:1. After five hours at 37° C, culture supernatants were harvested and analyzed as described in Example 11. Percent specific killing was calculated according to the following equation: ((experiment release minus spontaneous release)/(maximum release minus spontaneous release)) x 100. The data are presented in Figure 30. Each data point represents the average of quadruplicate samples.

Using the same procedures described above, the same results with other binding domains were obtained using the following monoclonal antibodies as sources of scFv: for CD20, IF5 (Genbank AY 058907 and AY058906); for CD40, 2.36 and G28.5; for CD28, 9.3.

Cell surface expression of antibody binding domains is accomplished by fusing antibody scFvs to IgA hinge and constant regions and IgE hinge-acting region, *i.e.*, IgE CH2, and constant regions. Polynucleotides encoding an anti-4-1BB scFv, 5B9 (anti-human 4-1BB) scFv, and 2e12 (anti-human CD40) fused to IgAH IgA T4 (four terminal CH3 residues deleted) fused to the CD80 transmembrane and cytoplasmic domains and IgE Fc regions are shown in SEQ ID NOs: 177,181, 179

and 183. The encoded polypeptides are shown in SEQ ID NOs: 178, 182, 180, and 184.

EXAMPLE 13

CONSTRUCTION AND SEQUENCE OF HUMAN 1G HINGE-CH2-CH3 MUTANTS AND 2H7 VARIABLE REGION MUTANTS

5

This Example describes construction of scFv fusion proteins containing mutant human IgG1 and IgA constant regions. This Example also describes construction of a 2H7 scFv mutant with a single point mutation in the variable heavy chain region. Mutations were introduced into variable and constant region domains according to methods described herein and known in the molecular biology arts. Figure 31 presents nomenclature for the Ig constant region constructs.

10

The human IgG1 hinge region of the 2H7 scFv human IgG1 hinge-CH2-CH3 fusion proteins was mutated to substitute cysteine residues that in a whole immunoglobulin are involved in forming disulfide bonds between two heavy chain molecules. One mutant, 2H7 scFv fused to a human IgG1 hinge region in which all three cysteine residues were mutated to serine residues (MTH (SSS)), was prepared as described in Example 5 (designated in Example 5 as CytoxB-MHWTG1C (includes wild type IgG1 CH2 and CH3 domains)) (now referred to as 2H7 scFv MTH (SSS) WTCH2CH3) and comprises the polynucleotide sequence SEQ ID NO: 57 encoding the polypeptide as set forth in SEQ ID NO: 58. The polynucleotide sequence encoding this mutant (SEQ ID NO: 57) was used as a template to create mutant hinge regions in which the first two cysteine residues were substituted with serine residues (IgG MTH (SSC)). An oligonucleotide was designed to substitute the third serine residue with a cysteine and had the following sequence: 5'-gtt gtt gat cag gag ccc aaa tct tct gac aaa act cac aca tct cca ccg tgc cca gca cct g-3' (HuIgGMHncs3, SEQ ID NO: 548). A second mutant was prepared in which the mutant hinge had serine residues substituting the first and third cysteine residues (IgG MTH (SCS)). The sequence of the oligonucleotide to create this mutant was as follows: 5'-gtt gtt gat cag gag ccc aaa tct tct gac aaa act cac aca tgc cca ccg-3' (HuIgGMHncs2, SEQ ID NO: 549). A third mutant was prepared with cysteine residues substituted at the second and third positions (IgG MTH (CSS)), also using the IgG MTH (SSS) mutant as template, and

15

20

25

30

an oligonucleotide having the sequence, 5'-gtt gtt gat cag gag ccc aaa tct tgt gac aaa act cac-3' (HuIgGMHncs1, SEQ ID NO: 550).

The oligonucleotides introducing the mutations into the hinge region were combined with template and a 3' oligonucleotide containing an XbaI site (underlined and italicized) (5'-gtt gtt tct aga tca ttt acc cgg aga cag gga gag gct ctt ctg cgt gta g-3' (SEQ ID NO: 551) to amplify the mutant hinge-wild type (WT)-CH2-CH3 sequences by PCR. The IgG MTH CSS and IgG MTH SCS mutant sequences were amplified for 25 cycles with a denaturation profile of 94° C, annealing at 52° C for 30 seconds, and extension at 72° C for 30 seconds. The IgG MTH SSC mutant sequence was amplified under slightly different conditions: denaturation profile of 94° C, annealing at 45° C for 30 seconds, and extension at 72° C for 45 seconds. The amplified polynucleotides were inserted into the TOPO® cloning vector (Invitrogen Life Technologies) and then were sequenced as described in Example 1 to confirm the presence of the mutation. pD18 vector containing 2H7 scFv was digested to remove the constant region sequences essentially as described in Example 10. The mutant hinge-wild type CH2-CH3 regions were inserted in frame into the digested vector DNA to obtain vectors comprising 2H7 scFv MTH (CSS) WTCH2CH3 encoding DNA (SEQ ID NO: 134); 2H7 scFv MTH (SCS) WTCH2CH3 encoding DNA (SEQ ID NO: 136); and 2H7 scFv MTH (SSC) WTCH2CH3 encoding DNA (SEQ ID NO: 138).

A mutation of leucine to serine at position 11 in the first framework region of the heavy chain variable region (numbering according to Kabat et al., Sequences of Proteins of Immunological Interest, 5th ed. Bethesda, MD: Public Health Service, National Institutes of Health (1991)) was introduced into the 2H7 scFv MTH (SSS) WTCH2CH3 fusion protein (SEQ ID NO: 58). The wild type leucine residue was substituted with serine by site-directed mutagenesis using the oligonucleotide Vhser11: 5'-gga ggt ggg agc tet cag gct tat cta cag cag tct ggg get gag teg gtg agg cc-3' (SEQ ID NO: 552). The 3'-primer for PCR was huIgG1-3' having the sequence 5'-gtc tct aga cta tca ttt acc cgg aga cag-3' (SEQ ID NO: 553) (XbaI site underlined and italicized). After PCR amplification, the fragments were inserted into the TOPO® cloning vector and sequenced to confirm the presence of the VH11 leucine to serine mutation. The 2H7 scFv-IgG MTH (SSS) WTCH2CH3 encoding DNA was shuttled into the PSL1180 cloning vector (Pharmacia Biotech, Inc., Piscataway, NJ). The

construct PSL1180-2H7 scFv-IgG MTH (SSS) WTCH2CH3 was digested with Sac and XbaI to remove the wild type V_H domain and the hinge and CH2 and CH3 domains. The PCR product comprising the VH11 mutant was digested with Sac and XbaI and then inserted into the digested PSL 1180 construct according to standard molecular biology procedures. The construct was then digested with Hind III and XbaI, and inserted into the mammalian expression vector pD18 (see methods described in Example 1 and Example 10). The mutant is designated 2H7 scFv VH11SER IgG MTH (SSS) WTCH2CH3 (Figure 31). The polynucleotide sequence is provided in SEQ ID NO: 369, and the encoded polypeptide sequence is provided in SEQ ID NO: 370.

Four constructs containing IgA constant region domains were prepared. One construct contained wild type IgA hinge fused to human IgG1 CH2 and CH3 (IgAH IgG WTCH2CH3) (Figure 31). Sequential PCR amplifications were performed to substitute the human IgG1 hinge of the 2H7 scFv construct with nucleotide sequences encoding the IgA hinge. The 5' oligonucleotide primer (hulgA/Gchim5) for the first PCR reaction had the sequence, 5'-cca tct ccc tca act cca cct acc cca tct ccc tca tgc gca cct gaa ctc ctg-3' (SEQ ID NO: 554). The primer (hulgAhg-5') for the second PCR reaction to add more IgA specific hinge sequence and add a BclI restriction enzyme site (*italicized and underlined*) had the sequence, 5'-gtt gtt gat cag cca gtt ccc tca act cca cct acc cca tct ccc caa ct-3' (SEQ ID NO: 555). The 3' primer for both amplification steps was huIgG1-3' having the sequence, 5'-gtc tct aga cta tca ttt acc cgg aga cag-3' (SEQ ID NO: 556). The sequence of the PCR product was confirmed by TOPO® cloning as described above. The gel-purified fragment was digested with BclI and XbaI and then inserted into the 2H7 scFv-pD 18 vector that had been digested BclI and XbaI to remove all the IgG1 constant region domains. Ligation was performed as described in Example 10 to provide a mammalian expression vector comprising the nucleotide sequence (SEQ ID NO: 59) encoding a 2H7 scFv IgA hinge-IgG1-CH2-CH3 polypeptide (SEQ ID NO: 60).

A second pD18 mammalian expression vector was constructed that had a polynucleotide sequence (SEQ ID NO: 61) that encoded a 2H7 scFv fused to wild type IgA hinge, CH2, and CH3 domains (SEQ ID NO: 62). Human IgA constant regions sequences were obtained by using random primers to reverse transcribe total RNA isolated from human tonsil followed by PCR amplification of the cDNA using

sequence specific primers, essentially as described in Example 10. Human IgA hinge-CH2-CH3 nucleotide sequence (SEQ ID NO: 63) encoding the IgA-CH2-CH3 polypeptide (IgAH IgACH2CH3, Figure 31) (SEQ ID NO: 64) was amplified using the 5' oligonucleotide hulgAhg-5' (SEQ ID NO: 555) and a 3' oligonucleotide hulgA3' having the sequence, 5'-gtt gtt tct aga tta tca gta gca ggt gcc gtc cac ctc cgc cat gac aac-3' (SEQ ID NO: 557). Secretion of a 2H7-IgA hinge-IgA CH2-CH3 polypeptide from transfected mammalian cells required co-expression of human J chain that covalently binds to two IgA CH3 domains via disulfide bonds. Total RNA was isolated from tonsil B cells and was reversed transcribed to generate cDNA as described above. PCR amplification of the nucleotide sequence encoding the J chain was performed with J chain specific primers. The 5' PCR primer, HUIJCH5nl, had the sequence, 5'-gtt gtt aga tct caa gaa gat gaa agg att gtt ctt-3' (SEQ ID NO: 558), and sequence of the 3' primer, HUIJCH3, was 5'-gtt gtt tct aga tta gtc agg ata gca ggc atc tgg-3' (SEQ ID NO: 559). The cDNA was cloned into TOPO(V for sequencing as described in Example 10. J chain encoding cDNA (SEQ ID NO: 65) was then inserted into pD18 and pCDNA3-Hygro (+) (Invitrogen Life Technology) vectors for co-transfection with 2H7 scFv IgA hinge-CH2-CH3 constructs. The J chain has the predicted amino acid sequence set forth in SEQ ID NO: 66.

Secretion of an scFv IgA constant region construct in the absence of J chain was accomplished by engineering a truncated CH3 domain with a deletion of the four carboxy terminal amino acids (GTCY, SEQ ID NO: 67) (IgAH IgA-T4, Figure 31), which include a cysteine residue that forms a disulfide bond with the J chain. The IgA hinge-CH2-CH3 nucleotide sequence containing the deletion in CH3 (SEQ ID NO: 68) was prepared using a 5' PCR primer (hulgAhg-5') having the sequence 5'-gtt gtt gat cag cca gtt ccc tca act cca cct acc cca tct ccc tca act-3' (SEQ ID NO: 561) (BclI site is underlined and italicized), and a 3' PCR primer (HUIGA3T1) having the sequence 5'-gtt gtt tct aga tta tca gtc cac ctc cgc cat gac aac aga cac-3' (SEQ ID NO: 562). This mutated IgA constant region nucleotide sequence was inserted into a 2H7 scFv pD18 vector as described for the generation of the previous 2H7 scFv-Ig constructs (see Example 1 and this example) that comprises the polynucleotide sequence (SEQ ID NO: 70) encoding a 2H7 IgAH IgA-T4 polynucleotide (SEQ ID NO: 71).

A fourth construct was prepared that encoded a 2H7 scFv-IgA constant region fusion protein with a deletion of 14 additional amino acids, most of which are hydrophobic residues, from the carboxy terminus of IgA CH3. The 2H7 scFv-IgAH IgA-T4 encoding polynucleotide was used as template to engineer a deletion of the nucleotide sequence encoding PTHVNVSVVMAEVD (SEQ ID NO: 72). The 5' oligonucleotide primer had the sequence 5'-ggt ggt gat cag cca gtt ccc tca act cca cct acc cca tct ccc tca act-3' (SEQ ID NO: 564) (BclI site shown as underlined and italicized). The 3' oligonucleotide sequence was 5'-ggt gtt tct aga tta tca ttt acc cgc caa gcg gtc gat ggt ctt-3' (SEQ ID NO: 565). The deleted IgA CH3 region was amplified by using the above oligonucleotides to amplify the IgA constant region from RNA isolated from human tonsil such that the cDNA contained the deleted carboxyl terminal encoding region for the 18 amino acids. The IgAH IgA-T18 constant region was inserted into a 2H7 scFv pD18 vector that comprises the polynucleotide sequence (SEQ ID NO: 75) encoding a 2H7 IgAH IgA-T18 polynucleotide (SEQ ID NO: 76) as described above.

EXAMPLE 14

EFFECTOR FUNCTION OF CTLA-4 IGG FUSION PROTEINS

The Example compares the effector functions of CTLA-4 Ig fusion proteins in CDC and ADCC assays.

Two CTLA-4 IgG fusion proteins were constructed. One fusion protein comprises the extracellular domain of CTLA-4 fused to human IgG1 wild type hinge, CH2, and CH3 domains and is designated CTLA-4 IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 78). A pD18 mammalian expression vector comprising a polynucleotide sequence encoding CTLA-4 IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 77) was prepared by fusing in frame the nucleotide sequence encoding the extracellular domain of CTLA-4 (SEQ ID NO: 83) (see U.S. Pat. No. 5,844,095) to the nucleotide sequence encoding IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 1) according to the methods described in Examples 1 and 10. The extracellular domain nucleotide sequence also comprises a BclI restriction enzyme site at the 3' end, and a leader peptide nucleotide sequence (SEQ ID NO: 79) that encodes an oncoM leader peptide (SEQ ID NO: 80). A second CTLA-4 IgG fusion protein, designated CTLA-4 IgG MTH (SSS) MTCH2WTCH3, contained the extracellular domain of CTLA-4

(plus the oncoM leader peptide sequence) fused to a mutant IgG hinge in which all three cysteine residues were replaced with serine residues. The hinge region was fused to a mutant IgG1 CH2 domain that had a mutation at isotype position 238 (EU numbering, Ward et al., *supra*, (position 251 using numbering according to Kabat *et al.*, *supra*; position 209 where numbering commences with first residue of IgG1 CH1; *i.e.*, PAPELLDGPS (SEQ ID NO: 1) of wild type IgG1 CH2 is modified to PAPELLDGSS (SEQ ID NO: 1)), which was fused to IgG1 wild type CH3 (U.S. Pat. No. 5,844,095). The CTLA-4 IgG MTH (SSS) MTCH2WTCH3 polynucleotide comprises the nucleotide sequence in SEQ ID NO: 85 and the deduced amino acid sequence comprises the sequence provided in SEQ ID NO: 86. CTLA-4 fusion proteins were also prepared using CTLA-4 extracellular membrane encoding sequences without the leader peptide (SEQ ID NO: 84).

To measure CDC activity, purified CTLA-4 IgG WTH (CCC) WTCH2CH3 (2 µg/ml) or CTLA-4 IgG MTH (SSS) MTCH2WTCH3 (2 µg/ml) was added to Reh cells (see Example 12) and to Reh cells transfected with the costimulatory molecule CD80 such that CD80 was expressed on the cell surface (Reh CD80.10, obtained from Dr. E. Clark, University of Washington, Seattle, WA; see Doty *et al.*, 1998 *J. Immunol.* 161: 2700; Doty *et al.*, 1996 *J. Immunol.* 157: 3270), in the presence or absence of rabbit complement (10 µg/ml). Purified CTLA Ig fusion proteins were prepared from culture supernatants of transiently transfected COS cells according to methods described in Example 10. The assays were performed essentially as described in Example 11 and 12. The data presented in Figure 32 show that only CD80-transfected Reh cells were killed in the presence of complement and CTLA-4 IgG WTH (CCC) WTCH2CH3 fusion protein.

The purified CTLA-4 Ig fusion proteins were also tested in ADCC assays. Human PBMC, serving as effector cells, were added to Reh or Reh CD80.1 target cells at a ratio of 1.25:1, 2.5:1, 5.0:1, and 10:1. Cells were labeled and the assays performed essentially as described in Examples 11 and 12. The results are presented in Figure 33. Each data point represents the average of four independent culture wells at each effector:target cell ratio. The data show that only CTLA-4 IgG WTH (CCC) WTCH2CH3 mediated significant ADCC of Reh CD80.10 cells.

EXAMPLE 15

EFFECTOR FUNCTION OF CTLA-4 IGA FUSION PROTEINS

CTLA-4 IgA fusion proteins are prepared as described for the IgG fusion proteins (see Examples 1, 13, and 14). CTLA-4 extracellular domain nucleotide sequence (SEQ ID NO: 84) is fused in open reading frame to nucleotides encoding IgAH IgACH2CH3 (SEQ ID NO: 63) to provide the nucleotide sequence (SEQ ID NO: 87) encoding a CTLA-4 IgAH IgACH2CH3 fusion protein (SEQ ID NO: 88). The fusion protein is transiently expressed in COS cells (see Example 10) or stably expressed in CHO cells (see Example 1). Secretion of the CTLA-4 IgAH IgACH2CH3 fusion protein requires co-transfection with a construct containing a polynucleotide sequence (SEQ ID NO: 65) that encodes human J chain (SEQ ID NO: 66). The CTLA-4 IgAH IgACH2CH3 fusion protein is isolated as described in Examples 10 and 14. To express a CTLA-4 IgA construct without the presence of J chain, a CTLA-4 IgAH IgA-T4 construct is prepared and transfected into mammalian cells. In a similar manner as described for the CTLA-4 extracellular fragment fused to wild type IgA hinge-CH2CH3, the CTLA-4 extracellular domain nucleotide sequence (SEQ ID NO: 84) is fused in open reading frame to a nucleotide sequence (SEQ ID NO: 68) encoding a IgAH IgA-T4 polypeptide (SEQ ID NO: 69) to provide a nucleotide sequence comprising SEQ ID NO: 89 encoding a CTLA-4 IgAH IgA-T4 polypeptide (SEQ ID NO: 90). Effector function of each construct is evaluated by CDC and ADCC as described in Example 14.

EXAMPLE 16

BINDING OF ANTI-CD20 SCFV HUMAN IG FUSION PROTEINS TO CHO CELLS

EXPRESSING CD20

This Example describes binding of 2H7 scFv Ig fusion proteins to CHO cells that express CD20. The analysis was performed by flow cytometry. Culture supernatants were collected from transiently transfected COS cells expressing 2H7 scFv IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 28); 2H7 scFv IgG MTH (CSS) WTCH2CH3 (SEQ ID NO: 135); 2H7 scFv IgG MTH (SCS) WTCH2CH3 (SEQ ID NO: 137); and 2H7 scFv VHSE11 WTH WTCH2CH3, and two-fold serial dilutions were prepared. Serial two-fold dilutions of purified 2H7 scFv IgG MTH (SSC) WTCH2CH3 (SEQ ID NO: 139) were prepared starting at a concentration of 5

µg/ml. The culture supernatants and purified fusion protein samples were incubated with (CD20+) CHO cells for one hour on ice. The cells were washed twice and then incubated with 1:100 FITC-conjugated goat anti-human IgG (CalTag) for 40 minutes. The unbound conjugate was then removed by washing the cells and flow cytometry analysis was performed using a Coulter Epics XL cell sorter. Results are presented in Figure 34.

EXAMPLE 17

IMMUNOBLOT ANALYSIS OF ANTI-CD20 scFv HUMAN IgG AND IgA FUSION

PROTEINS

10 This Example describes immunoblot analysis of 2H7 scFv IgG and 2H7 scFv IgA fusion proteins that were immunoprecipitated from transfected cell culture supernatants.

COS cells were transiently transfected with plasmids comprising nucleotide sequences for 2H7 scFv IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 28); 15 2H7 scFv IgG MTH (CSS) WTCH2CH3 (SEQ ID NO: 135); 2H7 scFv IgG MTH (SCS) WTCH2CH3 (SEQ ID NO: 137); 2H7 scFv IgA H IgG WTCH2CH3 (SEQ ID NO: 60); and scFv IgG MTH (SSS) WTCH2CH3 (SEQ ID NO: 58) essentially according to the method described in Example 10. Cells were also transfected with vector only. After 48-72 hours at 37° C, cell culture supernatants were harvested and 20 combined with protein A-agarose beads (Repligen) for one hour at 4° C. The beads were centrifuged and washed several times in TNEN [20 mM Tris base, 100 mM NaCl, 1 mM EDTA, and 0.05% NP-40, pH 8.0). The immunoprecipitates were combined with 25 µl 2x NUPAGE® SDS Sample Buffer (Invitrogen Life Technologies) (non-reduced samples). The proteins were fractionated on NUPAGE® 25 10% Bis-Tris gels (Invitrogen Life Technologies). After electrophoresis (approximately 1 hour), the proteins were transferred from the gel onto a Immobilon P polyvinylidene fluoride (PVDF) membrane (Millipore, Bedford, MA) using a semi-dry blotter (Ellard Instrumentation, Monroe, WA). The PVDF membrane was blocked in PBS containing 5% nonfat milk and then probed with HRP-conjugated goat anti- 30 human IgG (Fc specific) (CalTag). After washing the immunoblot several times in PBS, the blot was developed using ECL (Amersham Biosciences). The results are shown in Figure 35.

EXAMPLE 18

BINDING OF ANTI-CD20 scFv HUMAN IGA FUSION PROTEINS TO CD20+CHO CELLS

This Example describes flow immunocytofluorimetry analysis of
5 binding of 2H7 scFv IgAH IgACH2CH3 (SEQ ID NO: 62) and 2H7 scFv IgAH
IgAT4 (SEQ ID NO: 70) fusion proteins to (CD20+) CHO cells.

COS cells were transiently co-transfected as described in Example 10
with plasmid DNA comprising a polynucleotide sequence (SEQ ID NO: 61) encoding
2H7 scFv IgAH IgACH2CH3 polypeptide (SEQ ID NO: 62) and with a separate
10 plasmid comprising a polynucleotide sequence (SEQ ID NO: 65) encoding a human J
chain polypeptide (SEQ ID NO: 66). COS cells were also transfected with a
polynucleotide sequence (SEQ ID NO: 70) encoding a 2H7 scFv IgA fusion protein
that had a deletion of four amino acids at the carboxy terminus of CH3 (2H7 scFv
IgAH IgA-T4, SEQ ID NO: 71). The transfections were performed as described in
15 Example 10, Culture supernatants from transfected COS cells were combined with
(CD20+) CHO cells (see Example 1) and incubated for one hour on ice. The cells
were washed twice with PBS-2% FBS and then combined with FITC-conjugated goat
anti-human IgA chain (CalTag) (1:100) for 40 minutes. The cells were again washed
and then analyzed by flow cytometry using a Coulter Epics XL cell sorter. Figure 36
20 shows that co-transfection with J chain was not required for secretion of 2H7 scFv
IgAH IgAT4, the 2H7 IgA fusion protein with the truncated CH3 carboxy end (SEQ
ID NO: 71).

EXAMPLE 19

EFFECTOR FUNCTION OF ANTI-CD20 scFv HUMAN IGA FUSION PROTEINS

25 This Example illustrates ADCC activity of 2H7 IgG and IgA fusion
proteins against cells that express CD20. BJAB cells were prelabeled with ⁵¹Cr (100
μCi) (Amersham) for two hours at 37° C Effector cells were obtained from fresh,
resting human whole blood, which was diluted in an equal volume of Alsever's
solution to prevent coagulation. 2H7 scFv IgG MTH (SSS) WTCH2CH3 (SEQ ID
30 NO: 58); 2H7 scFv IgG MTH (SCS) WTCH2CH3 (SEQ ID NO: 137); 2H7 scFv IgG
WTH (CCC) WTCH2CH3 (SEQ ID NO: 28); and 2H7 scFv IgAH IgACH2CH3
(SEQ ID NO: 62) fusion proteins were purified from transiently transfected COS cell

supernatants (100-200 ml) by protein A chromatography as described in Example 10. COS cells transfected with the plasmid encoding 2H7 scFv IgAH IgACH2CH3 were co-transfected with a plasmid encoding human J chain as described in Example 18. Two-fold serial dilutions of the purified 2H7 Ig fusion proteins starting at 5 µg/ml were added to the labeled BJAB cells (5×10^4 cells per well of 96 well tissue culture plate) in the presence of whole blood (100 µL of whole blood diluted 1:1 in Alsever's solution, final dilution 1:4) and incubated for five hours at 37° C. Culture supernatants were harvested and analyzed as described in Example 1. Percent specific killing was calculated according to the following equation: $((\text{experiment release} - \text{spontaneous release}) / (\text{maximum release} - \text{spontaneous release})) \times 100$. The data are presented in Figure 37. Each data point represents the average of quadruplicate samples.

In a second ADCC assay, the number of labeled BJAB target cells was held constant in each sample, and whole blood was added at dilutions of 0.25, 0.125, and 0.0625. Purified 2H7 IgG and IgA fusion proteins were added at a concentration of 5 µg/ml. The BJAB cells, whole blood, and fusion proteins were incubated, the supernatants harvested, and the percent specific killing was calculated as described above. Percent specific killing for each of the 2H7 fusion proteins is presented in Figure 38.

The ADCC activity of purified 2H7 scFv IgG MTH (SSS) WTCH2CH3 (5 µg/ml) and of purified 2H7 scFv IgAH IgACH2CH3 (5 µg/ml) was compared in the presence of different effector cell populations. PBMC were isolated from whole blood as described in Examples 11 and 12. PBMC were combined with labeled BJAB target cells (5×10^4 per well of 96 well tissue culture plate) at ratios of 50:1, 25:1, and 12.5:1. The assay was performed and the data analyzed as described above. Figure 39A shows that only the 2H7 scFv IgG MTH (SSS) WTCH2CH3 fusion protein had ADCC activity when PBMC served as the effector cells. Figure 39B shows that both 2H7 scFv IgG MTH (SSS) WTCH2CH3 and 2H7 scFv IgAH IgACH2CH3 exhibit ADCC activity when whole blood was the source of effector cells (as illustrated in Figure 38).

EXAMPLE 20

EXPRESSION LEVEL OF 2H7 scFv VH11SER IgG MTH (SSS) WTCH2CH3 FUSION PROTEIN

This Example compares the expression level of 2H7 scFv VH11Ser
5 IgG MTH (SSS) WTCH2CH3 fusion protein (SEQ ID NO: 370) with other 2H7 scFv
IgG constructs that do not contain the mutation in the variable heavy chain domain.
The mammalian expression vector pD 18 comprising nucleotide sequences 2H7 scFv
IgG MTH (SSS) WTCH2CH3 (SEQ ID NO: 370); 2H7 scFv IgG MTH (CSS)
WTCH2CH3 (SEQ ID NO: 372); 2H7 scFv IgG MTH (SCS) WTCH2CH3 (SEQ ID
10 NO: 137); 2H7 scFv IgG WTH (CCC) WTCH2CH3 (SEQ ID NO: 28); and 2H7 scFv
VHSER11 IgG MTH (SSS) WTCH2CH3 (see Examples 1 and 13) were transiently
transfected into COS cells as described in Example 10. After 72 hours at 37° C,
culture supernatants were harvested, and 1 µl of each supernatant was combined with
non-reducing sample buffer (see method described in Example 10). The culture
15 supernatant samples and aliquots of purified 2H7 scFv IgG MTH (SSS) WTCH2CH3
(40 ng, 20 ng, 10 ng/5 ng, and 2.5 ng) were fractionated on 10% Bis-Tris (MOPS)
NUPAGE® gels (Invitrogen Life Technologies). MULTIMARK® protein standards
(Invitrogen Life Technologies) were also separated on the gel. The proteins were
transferred to a PDVF membrane and immunoblotted as described in Example 17.
20 The immunoblot is presented in Figure 40. The amounts of the fusion proteins were
quantified by densitometry analysis of the blots using the ScionImage for Windows
software and comparison with the standard curve. The 2H7 scFv IgG WTH (CCC)
WTCH2CH3 construct produced approximately 12 ng/ul or 12 micrograms/ml, the
2H7 scFv IgG MTH (CSS) WTCH2CH3 produced approximately 10 ng/ul or 10
25 micrograms/ml, the 2H7 scFv IgG MTH (SCS) WTCH2CH3 construct produced
approximately 1 ng/ul or 1 microgram/ml, and the 2H7 scFv VHSER11 IgG MTH
(SSS) WTCH2CH3 construct produced approximately 30 ng/ml or 30 micrograms/ml.

EXAMPLE 21

CONSTRUCTION OF A 2H7 scFv IgG FUSION PROTEIN WITH A MUTANT CH3 DOMAIN

30 Amino acid mutations were introduced into the CH3 domain of a 2H7
IgG fusion protein. The pD18 vector comprising 2H7 scFv IgG MTH (SSS)
WTCH2CH3 (SEQ ID NO: 135) was digested with BclI and XbaI to remove the

MTH WTCH2CH3 fragment, which was then subcloned into pShuttle vector (BD Biosciences Clontech, Palo Alto, CA) that was double-digested with BclI and XbaI. Subcloning was performed in a kanamycin resistant vector because the ampicillin resistance gene has an XmnI site, which is required for this cloning procedure. Five constructs were prepared with the following substitutions: (1) a phenylalanine residue at position 405 (numbering according to Kabat *et al. supra*) was substituted with tyrosine using the oligonucleotide CH3Y405; (2) the phenylalanine position at 405 was substituted with an alanine residue using the oligonucleotide CH3A405; (3) the tyrosine residue at position 407 was substituted with an alanine using the oligonucleotide CH3A407; (4) both wild type amino acids at positions 405 and 407 were substituted with tyrosine and alanine, respectively using the oligonucleotide CH3Y405a407; and (5) both wild type amino acids at positions 405 and 407 were substituted with alanine using the oligonucleotide CH3A405a407. The oligonucleotides were the 3' primers for PCR amplification of a portion of the CH3 domain. The nucleotide sequences for each 3' oligonucleotide were as follows.

CH3Y405: 5'-gtt gtt gaa gac gtt ccc ctg ctg cca cct gct ctt gtc cac ggt gag ctt gct gta gag gta gaa gga gcc-3' (SEQ ID NO: 568)

CH3A405: 5'-gtt gtt gaa gac gtt ccc ctg ctg cca cct gct ctt gtc cac ggt gag ctt gct gta gag ggc gaa gga gcc-3' (SEQ ID NO: 569)

CH3A407: 5'-gtt gtt gaa gac gtt ccc ctg ctg cca cct gct ctt gtc cac ggt gag ctt gct ggc gag gaa gga gcc-3' (SEQ ID NO: 570)

CH3Y405A407: 5'-gtt gtt gaa gac gtt ccc ctg ctg cca cct gct ctt gtc cac ggt gag ctt gct ggc gag gta gaa gga gcc-3' (SEQ ID NO: 571)

CH3A405A407: 5'-gtt gtt gaa gac gtt ccc ctg ctg cca cct gct ctt gtc cac ggt gag ctt gct ggc gag ggc gaa gga gcc-3' (SEQ ID NO: 572)

The template was the mutant hinge MHWTC2CH3 human IgG1. The 5' PCR oligonucleotide primer was huIgGMHWC. The amplified products were TOPO® cloned and sequenced as described in Examples 1 and 10. DNA from the clones with the correct sequence was digested with BclI and XmnI and transferred to pShuttle containing the MTH WTCH2CH3 sequence, which was also digested with the same restriction enzymes. The mutated IgG sequences were then removed by

digestion with BclI and XbaI and inserted into a pD18 vector containing 2H7 scFv that was also digested with BclI and XbaI. The polynucleotide sequences for mutated the CH3 domains, MTCH3 Y405, MTCH3 A405, MTCH3 A407, MTCH3 Y405a407, and MTCH3 A405a407 are shown in SEQ ID NOs: 143, 145, 147, 149, respectively, and the polypeptide sequences for each are shown in SEQ ID NOs: 144, 146, 148, 150, respectively. The polynucleotide sequences for the 2H7 scFv MTH WTCH2 MTCH3 Y405, 2H7 scFv MTH WTCH2 MTCH3 A405, scFv MTH WTCH2 MTCH3 A407, scFv MTH WTCH2 MTCH3 Y405a407, and scFv MTH WTCH2 MTCH3 A405a407, respectively, and the deduced amino acid sequences are shown in SEQ ID NOs: 154, 156, 158, 160, 162, and the deduced nucleotide sequences are shown in SEQ ID NOs: 153, 155, 157, 159, 161, respectively.

EXAMPLE 22

CONSTRUCTION OF 2H7 scFv IgG FUSION PROTEINS WITH HINGE MUTATIONS

A 2H7 scFv IgG fusion protein was constructed with the third cysteine residue in the IgG1 hinge region substituted with a serine residue. The template for introduction of the mutations was a polynucleotide encoding 2H7 scFv WTH WTCH2CH3 (SEQ ID NO: 28). The oligonucleotide introducing the mutations was a 5' PCR primer oligonucleotide HIgGMHcys3 having the sequence 5'-gtt gtt gat cag gag ccc aaa tct tgt gac aaa act cac aca tgt cca ccg tcc cca gca cct-3'. The oligonucleotide introducing the mutation into the hinge region was combined with template and a 3' oligonucleotide containing an XbaI site (underlined and italicized) (5'-gtt gtt *tct aga* tca ttt acc cgg aga cag gga gag gct ctt ctg cgt gta g-3' (SEQ ID NO: 574)) to amplify the mutant hinge-wild type (WT)-CH2-CH3 sequences by PCR. The IgG MTH CCS mutant sequence was amplified for 30 cycles with a denaturation profile of 94° C, annealing at 50° C for 30 seconds, and extension at 72° C for 30 seconds. The amplified polynucleotides were inserted into the TOPO® cloning vector (Invitrogen Life Technologies) and then were sequenced as described in Example 1 to confirm the presence of the mutation. pD18 vector containing 2H7 scFv was digested to remove the constant region sequences essentially as described in Example 10. The mutant hinge-wild type CH2-CH3 regions were inserted in frame into the digested vector DNA to obtain vectors comprising 2H7 scFv MTH (CCS) WTCH2CH3 encoding DNA (SEQ ID NO: 167). The deduced polypeptide sequence is shown in SEQ ID NO: 168.

EXAMPLE 23

CONSTRUCTION OF ANTI-CD20 IGE FUSION PROTEINS

A binding domain is fused to IgE constant region sequences such that the expressed polypeptide is capable of inducing an allergic response mechanism. The single chain Fv nucleotide sequence of 40.2.220 (SEQ ID NO: 35), an anti-CD40 antibody, is fused to IgE CH2--CH3-CH4 according to methods described for other scFv immunoglobulin constant region constructs (see Examples 1, 5, 10, and 13). To PCR amplify the IgE CH2-CH3-CH4 domains, a 5' oligonucleotide primer, hIgE5Bcl, having the sequence 5'-gtt gtt gat cac gtc tgc tcc agg gac ttc acc cc-3', and a 3' (SEQ ID NO: 575) oligonucleotide primer, hIgE3stop, having the sequence 5'-gtt gtt tct aga tta act ttt acc ggg att tac aga cac cgc tgc ctg g-3' (SEQ ID NO: 576) are used.

The retroviral transfection system for ectopic surface expression of genetically engineered cell surface receptors composed of scFvs that bind costimulatory receptors described in Example 12 is used to construct a 40.2.220 scFv IgE-CD80 fusion protein. The 40.2.220 scFv IgE fusion polynucleotide sequence is fused in frame to sequences encoding the transmembrane domain and cytoplasmic tail of human CD80 (SEQ ID NO: 30), such that when the fusion protein is expressed in the transfected cell, CD80 provided an anchor for surface expression of the scFv Ig fusion protein. cDNA encoding the anti-CD40 scFv-IgE-CD80 fusion proteins is inserted into the retroviral vector pLNCX (BD Biosciences Clontech) according to standard molecular biology procedures and vendor instructions. The 40.2.220 scFv-Ig-CD80 cDNA is inserted between the 5'LTR-neomycin resistance gene-CMV promoter sequences and the 3'LTR sequence. The retroviral constructs are transfected into a carcinoma cell line, and transfected cells are screened to select clones that are expressing the 40.2.220 scFv-Ig-CD80 fusion protein on the cell surface.

EXAMPLE 24

CONSTRUCTION OF IGA-T4 MUTANTS THAT ARE EXPRESSED ON THE CELL SURFACE

The retroviral transfection system for ectopic surface expression of genetically engineered cell surface receptors composed of scFvs that bind costimulatory receptors described in Example 12 is used to construct a 2H7 scFv IgA hinge IgA-T4-CD80 fusion protein. The 2H7 scFv IgAH IgA-T4 fusion

polynucleotide sequence (SEQ ID NO: 70) is fused in frame to sequences encoding the transmembrane domain and cytoplasmic tail of human CD80 (SEQ ID NO: 29), such that when the fusion protein is expressed in the transfected cell, CD80 provided an anchor for surface expression of the scFv Ig fusion protein. cDNA encoding the 2H7 scFv IgAH IgA-T4-CD80 fusion protein is inserted into the retroviral vector pLNCX (BD Biosciences Clontech) according to standard molecular biology procedures and vendor instructions. The 2H7 scFv IgAH IgA-T4-CD80 cDNA is inserted between the 5'LTR-neomycin resistance gene-CMV promoter sequences and the 3'LTR sequence. The retroviral construct is transfected into Reh, an acute lymphocytic leukemia cell line (ATCC CRL-8286). Transfected cells are screened to select clones that are expressing 2H7 scFv-Ig fusion proteins on the cell surface.

EXAMPLE 25

CHARACTERIZATION OF AN ANTI-4-1BB scFv Ig-CD80 FUSION PROTEIN EXPRESSED ON THE CELL SURFACE OF TUMOR CELLS AND GROWTH OF THE TUMOR CELLS IN VIVO

This example describes construction of an anti-murine 4-1 BB (CD 137) scFv fusion protein that has an IgG wild type hinge and CH2 and CH3 domains that is fused to the CD80 transmembrane and cytoplasmic domains. The Example also illustrates the effect of the cell surface expression of the anti-4-1BB scFv IgG CD80 polypeptide when the transfected tumor cells are transplanted into mice.

The heavy and light chain variable regions of a rat anti-4-1BB (CD137) monoclonal antibody (1D8) were cloned, and a single chain Fv construct was prepared essentially as described in Example 1. The heavy chain and light chain variable regions of each antibody were cloned according to standard methods for cloning immunoglobulin genes and as described in Example 1. A single chain Fv construct was prepared as described in Example 1 by inserting a nucleotide sequence encoding a (gly₄ser)₃ peptide linker between the V_L region nucleotide sequence of 1D8 (SEQ ID NO: 102) and the V_H region nucleotide sequence of 1D8 (SEQ ID NO: 100). The polypeptide sequence for 1D8 V_L is shown in SEQ ID NO: 103, and the polypeptide sequence for the V_H domain is shown in SEQ ID NO: 101. The scFv polynucleotide (SEQ ID NO: 104) was then fused to human IgG1 wild-type hinge-CH2-CH3 domains according to the methods described in Example 1. The scFv IgG1 fusion polynucleotide sequence was then fused in frame to sequences encoding the

transmembrane domain and cytoplasmic tail of human CD80 (SEQ ID NO: 29) essentially as described in Example 12, such that when the fusion protein was expressed in the transfected cell, CD80 provided an anchor for surface expression of the scFv Ig fusion protein. cDNA encoding the scFv-IgG-CD80 fusion protein (SEQ ID NO: 110) was inserted into the retroviral vector pLNCX (BD Biosciences Clontech) according to standard molecular biology procedures and vendor instructions. The scFv-Ig-CD80 cDNA was inserted between the 5'LTR-neomycin resistance gene-CMV promoter sequences and the 3'LTR sequence.

The retroviral constructs were transfected into the metastatic M2 clone of K1735, a melanoma cell line, provided by Dr. I. Hellstrom, PNRI, Seattle, WA. Transfected cells were screened to select clones that were expressing scFv-Ig fusion proteins on the cell surface. To demonstrate that the 1D8 scFv IgG-CD80 construct was expressed on the cell surface of the tumor cells, the transfected cells were analyzed by flow immunocytofluorimetry. Transfected cells (K1735-1D8) were incubated for one hour on ice in phycoerythrin-conjugated F(ab')₂ goat anti-human IgG. The unbound conjugate was then removed by washing the cells and flow cytometry analysis was performed using a Coulter Epics XL cell sorter. Results are presented in Figure 41A.

The growth of K1735-1D8 transfected cells was examined *in vivo*. K1735-WT cells grew progressively when transplanted subcutaneously (s.c.) in naive C3H mice. Although the same dose of K1735-1D8 cells initially formed tumors of an approximately 30 mm² surface area, the tumors started to regress around day 7 and had disappeared by day 20 as shown in Figure 41B. Tumor cells that were transfected with a similarly constructed vector encoding a non-binding scFv, a human anti-CD28 scFv construct, grew as well as tumor cells that had not been transfected. The presence of a foreign protein, that is, human IgG1 constant domains or rat variable regions, did not make transfected K1735-WT cells immunogenic; the growth of the K1735-1D8 cells in C3H mice was identical to that of K1735-WT cells (untransfected).

To investigate the roles of CD4⁺ and CD8⁺ T lymphocytes and NK cells in the regression of K1735-1D8 tumors, naive mice were injected intraperitoneally (i.p.) with monoclonal antibodies (monoclonal antibodies, typically 50 µg in a volume 0.1 ml) to remove CD8⁺, CD4⁺ or both CD4⁺ and CD8⁺ T cells, or

were injected with anti-asialo-GMI rabbit antibodies to remove NK cells. Twelve days later, when flow cytometry analysis of spleen cells from identically treated mice showed that the targeted T cell populations were depleted, K1735-1D8 cells were transplanted s.c to each T cell-depleted group. K1735-1D8 had similar growth kinetics in mice that had been injected with the anti-CD8 MAb or control rat IgG while removal of CD4⁺ T cells resulted in the growth of K1735-1D8 with the same kinetics as K1735-WT. This failure to inhibit tumor growth after CD4⁺ T cell removal was observed regardless of the presence or absence of CD8⁺ T cells. K1735-1D8 grew in all NK-depleted mice, although more slowly than in the CD4-depleted group. The results are presented in Figure 41C.

EXAMPLE 26

THERAPEUTIC EFFECT OF TUMOR CELLS EXPRESSING ANTI-4-1BB scFv IgG-CD80 FUSION PROTEIN

This Example examines the ability of K1735-1D8 transfected tumor cells expressing an anti-CD137 scFv on the cell surface to generate a sufficient immune response in mice to mediate rejection of established, untransfected wild type tumors. C3H mice were transplanted with K1735-WT tumors (2×10^6 cells/animal) and grown for approximately six days. Experiments were performed using mice with established K1735-WT tumors of 30 mm² surface area. Mice were vaccinated by s.c. injection of K1735-1D8 or irradiated K1735-WT cells on the contralateral side. Identical injections were repeated at the time points indicated in Figure 42. One group of animals was given four weekly injections of K1735-1D8 cells. According to the same schedule, another group was given irradiated (12,000 rads) K1735-WT cells, and a third group was injected with PBS. The data are plotted in Figure 42. The WT tumors grew progressively in all control mice and in all mice that received irradiated K1735-WT cells. In contrast, the tumors regressed in 4 of the 5 mice treated by immunization with K1735-1D8. The animals remained tumor-free and without signs of toxicity when the experiment was terminated 3 months later. In the fifth mouse, the tumor nodule decreased in size as long as the mouse received K1735-1D8 cells, but the tumor grew back after therapy was terminated.

In another experiment with 5 mice/group, mice were injected intravenously (i.v.) with 3×10^5 K1735-WT cells to initiate lung metastases. Three days later, K1735-1D8 cells were transplanted s.c. This procedure was repeated once

weekly for a month; control mice were injected with PBS. The experiment was terminated when one mouse in the control group died, 37 days after receiving the K1735-WT cells. At that time, lungs of the control mice each had >500 metastatic foci. In contrast, less than 10 metastatic foci were present in the lungs of the immunized mice.

In a third experiment, mixtures of K1735-WT cells and K1735-1D8 cells were injected into immunocompetent syngeneic C3H mice. Mice were injected subcutaneously with K1735-WT cells alone or with a mixture of 2×10^6 K1735-WT cells and 2×10^5 K1735-1D8 cells. Tumor growth was monitored at 5-day intervals.

10

EXAMPLE 27

EXPRESSION OF ANTI-4-1BB scFv IgG-CD80 FUSION PROTEIN ON THE CELL SURFACE OF SARCOMA CELLS

This Example demonstrates expression of an anti-CD137 scFv on the cell surface of a second type of tumor cell by transfecting a murine sarcoma cell line with an anti-CD137 scFv IgG-CD80 construct.

The 1D8 scFv IgG WTH WTCH2CH3-CD80 polynucleotide (SEQ ID NO: 110) was transferred from the pLNCX vector into pcDNA3-hygro vector using restriction enzyme digestion and ligation steps according to standard molecular biology methods. The construct was cut with HindIII+ClaI and the scFv fragment was filled in with Klenow (Roche) and the blunt-ended fragment was ligated into EcoRI site of pcDNA3. Ag104 murine sarcoma tumor cells were transfected with the pcDNA3-hygro vector containing the 1D8 scFv IgG CD80 fusion protein. Hygromycin-resistant clones were screened by flow cytometry using a FITC anti-human IgG antibody to detect expression of the transgene. Only approximately 15% of the resistant clones had detectable fusion protein initially. Positive cells identified by flow cytometry were repeatedly panned on flasks coated with immobilized anti-human IgG (10 μ g/ml) according to standard methods. Panning was performed by incubating cells on the coated plates for 30 min at 37°C; the plates were then washed 2-3x in versene or PBS. After each round, cells were tested for IgG expression by FACS. The histogram in Figure 44 shows the staining pattern after four rounds of panning against anti-human IgG (black). Untransfected cells were stained and are indicated in gray. All of the cells in the population were positive.

EXAMPLE 28

CONSTRUCTION AND CHARACTERIZATION OF A BISPECIFIC scFv Ig FUSION PROTEIN AND scFv Ig FUSION PROTEINS WITH A MUTATION IN THE IGG1 CH2 DOMAIN

5 An anti-CD20 (2H7) scFv IgG fusion protein was constructed that had a mutant hinge (MT (SSS)) and a mutant CH2 domain in which the proline at residue (position number 238 according to Ward *et al.*, *supra*) was substituted with a serine. The 2H7 scFv IgG MTH (S SS) MTCH2WTCH3 encoding polynucleotide (SEQ ID NO: 130) was constructed essentially according to methods described in Examples 1, 10 5, and 13. The IgG mutant hinge-mutant CH2-wild type CH3 domains were also fused to an anti-CD20 (2H7)-- anti-CD40 (40.2.220) bispecific scFv. The anti-CD20-anti-CD40 scFv IgG MTH (SSS) MTCH2WTCH3 encoding polynucleotide sequence is shown in SEQ ID NO: 114 and the encoded polypeptide is shown in SEQ ID NO: 115.

15 COS cells were transiently transfected with vectors comprising the polynucleotide sequences encoding 2H7 scFv IgG MTH (SSS) MTCH2WTCH3 (SEQ ID NO: 131); anti-CD20-anti-CD40 scFv IgG MTH (SSS) MTCH2WTCH3 (SEQ ID NO: 114); 2H7 scFv IgG MTH (SSS) WTCH2CH3 (SEQ ID NO: 58); and 2H7 scFv IgAH IgG WTCH2CH3 (SEQ ID NO: 60) as described in Example 10. 20 Culture supernatants were collected and the fusion proteins were purified by protein A chromatography (see Example 10). The purified polypeptides were fractionated by SDS-PAGE according to the method described in Example 10. Rituximab (anti-CD20 monoclonal antibody), and Bio-Rad prestained molecular weight standards (Bio-Rad, Hercules, CA), and MULTIMARK® molecular weight standards (Invitrogen Life 25 Technologies were also applied to the gel. The results are presented in Figure 45.

 The 2H7 scFv Ig fusion protein that contains a mutation in the CH2 domain was compared to fusion proteins that have the wild type CH2 domain in an ADCC assay. The assays were performed essentially as described in Examples 11 and 19. Fresh resting PBMC (effector cells) were added to ⁵¹Cr-labeled BJAB cells (target 30 cells) at the ratios indicated in Figure 46. Purified 2H7 scFv IgG MTH (SSS) MTCH2WTCH3, 2H7 scFv IgG MTH (SSS) WTCH2CH3, 2H7 scFv IgAH IgG WTCH2CH3, and Rituximab, each at 10 µg/ml were added to the effector/target cell mixtures and incubated for five hours at 37° C Supernatants were harvested and the

amount of chromium released was determined as described in Examples 1 and 19. Percent specific killing by each fusion protein is presented in Figure 46.

EXAMPLE 29

TUMOR CELL SURFACE EXPRESSION OF AN ANTI-HUMAN CD3 scFv IgG FUSION PROTEIN

5

An anti-human CD3 scFv Ig CD80 fusion protein was prepared essentially as described in Examples 1 and 12. The fusion protein comprised an anti-human CD3 scFv fused to wild type IgG1 hinge (SEQ ID NO: 12) and wild type CH2 (SEQ ID NO: 13) and CH3 (SEQ ID NO: 15) domains, fused to CD80 transmembrane and cytoplasmic domains (SEQ ID NO: 29) to enable cell surface expression of the anti-CD3 scFv. The anti-human CD3 scFv IgG WTH WTCH2CH3-CD80 polynucleotide (SEQ ID NO: 110) encoding the polypeptide (SEQ ID NO: 111) was transfected in Reh cells and into T51 cells (lymphoblastoid cell line). Expression of the anti-human CD3 scFv IgG fusion protein was detected by flow cytometry using FITC conjugated goat anti-human IgG (see methods in Examples 4, 10, 16, 18). Figure 47A illustrates expression of the anti-human CD3 fusion protein on the cell surface of Reh cells, and Figure 47B shows expression of the fusion protein on T41 cells.

10

15

ADCC assays were performed with the transfected Reh and T51 cells to determine if expression of the scFv-Ig polypeptides on the cell surface augmented effector cell function. Untransfected and transfected Reh cells and untransfected and transfected T51 cells were pre-labeled with ^{51}Cr (100 μCi) (Amersham) for two hours at 37° C Human PBMC served as effector cells and were added to the target cells (5x 10⁴ cells per well of 96 well plate) at ratios of 20:1, 10:1, 5:1, and 2.5:1. After four hours at 37° C, culture supernatants were harvested and analyzed as described in Examples 11 and 12. Percent specific killing was calculated as described in Example 12. The results are presented in Figure 48.

20

25

EXAMPLE 30

INDUCTION OF CYTOKINE EXPRESSION IN TUMOR CELLS EXPRESSING ANTI-CD28 SCFV ON THE CELL SURFACE

This Example describes the effect of cell surface expressed scFv on
5 cytokine mRNA induction in stimulated lymphocytes co-cultured with tumor cells
transfected with an anti-human CD28 scFv IgG-CD80 fusion protein.

Real time PCR analysis was performed on RNA samples from human
PBMC stimulated with Reh, Reh-anti-CD28 (2e12) (see Example 12 for construction
of 2e12 scFv IgG WTH WHTCH3CH2-CD80 and transfection of Reh cells), and
10 Reh-CD80 (see Example 14) in order to measure the effects of the surface expressed
scFv on cytokine production by the PBMC effector cells. For the real-time PCR
assay, SYBR Green (QIAGEN) (Morrison *et al.*, *Biotechniques* 24: 954-8, 960, 962
(1998)) was used and measured by an ABI PRISM® 7000 Sequence Detection
System (Applied Biosystems, Foster City, CA) that measures the formation of PCR
15 product after each amplification cycle. Cells were harvested from cultures and total
RNA prepared using QIAGEN RNA kits, including a QIA shredder column
purification system to homogenize cell lysates, and RNEASY® mini-columns for
purification of RNA. cDNA was reverse transcribed using equal amounts of RNA
from each cell type and Superscript II Reverse Transcriptase (Life Technologies).
20 SYBR Green real-time PCR analysis was then performed using the prepared cDNA as
template and primer pairs specific for cytokine gene products. The average length of
the PCR products that were amplified ranged from 150-250 base pairs. The cDNA
levels for many activation response molecules including IFN γ , TNF α , GM-CSF, IL-4,
IL-5, IL-6, IL-8, IL-10, IL-12, IL-15, ICOSL, CD80 and CD86 were assayed. Control
25 reference cDNAs for constitutively expressed genes, including GAPDH, β -actin, and
CD3 were measured in each assay. The most significant induction of specific mRNA
was observed for IFN- γ , and more modest induction was observed for CTLA-4 and
ICOS.

EXAMPLE 31

CLONING OF AN ANTI-HUMAN 4-1BB ANTIBODY AND CONSTRUCTION OF AN ANTI-HUMAN 4-1BB SCFV (IG FUSION PROTEIN)

A hybridoma cell line expressing a mouse anti-human monoclonal
antibody (designated 5B9) was obtained from Dr. Robert Mittler, Emory University

Vaccine Center, Atlanta, GA.. The variable heavy and light chain regions were cloned according to known methods for cloning of immunoglobulin genes and as described herein. Cells were grown in IMDM/15% FBS (Invitrogen Life Technologies) media for several days. Cells in logarithmic growth were harvested from cultures and total
5 RNA prepared using QIAGEN RNA kits, including a QIA shredder column purification system to homogenize cell lysates, and RNEASY® mini-columns for purification of RNA according to manufacturer's instructions. cDNA was reverse transcribed using random hexamer primers and Superscript II Reverse Transcriptase (Invitrogen Life Technologies).

10 cDNA was anchor-tailed using terminal transferase and dGTP. PCR was then performed using an anchor-tail complementary primer and a primer that annealed specifically to the antisense strand of the constant region of either mouse Ck (for amplification of VL) or the appropriate isotype mouse CH1 (for amplification of VH). The amplified variable region fragments were TOPO® cloned (Invitrogen Life
15 Technologies), and clones with inserts of the correct size were then sequenced. Consensus sequence for each variable domain was determined from sequence of at least four independent clones. The 5B9 V_L and V_H polynucleotide sequences are shown in SEQ ID NOs: 120 and 116, respectively, and the deduced amino acid sequences are shown in SEQ ID NOs: 121 and 118. The scFv was constructed by a
20 sewing PCR method using overlapping primers containing a synthetic (gly₄ser)₃ linker domain inserted between the light and heavy chain variable regions (see Example 1). The 5B9 scFv polypeptide (SEQ ID NO: 123) is encoded by the polynucleotide sequence comprising SEQ ID NO: 122.

5B9 scFv polynucleotide sequence was fused in frame to the
25 polynucleotide sequence encoding the human IgG1 mutant hinge and wild type CH2 and CH3 (MTH (SSS) WTCH2CH3,) according to methods described in Examples 5, 10, and 13. COS cells were transiently transfected with a vector comprising the 5B9 scFv IgG MTH (SSS) WTCH2CH3 polynucleotide sequence (SEQ ID NO: 132). Supernatant was collected and binding of the 5B9 scFv IgG MTH (SSS) WTCH2CH3
30 polypeptide (SEQ ID NO: 133) was measured by flow immunocytofluorimetry essentially as described in Examples 4, 10, 16, and 18. Culture supernatant from the 5B9 hybridoma cell line was also included in the binding assay. Fresh human PBMC were incubated in the presence of immobilized anti-CD3 for four days prior to the

binding experiment to induce expression of CD137 on the surface of activated T cells. Stimulated PBMC were washed and incubated with COS or hybridoma culture supernatant containing the 5B9 scFv IgG fusion protein or 5B9 murine monoclonal antibody, respectively, for 1 hour on ice. Binding of 5B9 scFv IgG fusion protein or 5B9 murine monoclonal antibody was detected with FITC conjugated anti-human IgG or anti-mouse IgG, respectively. The results are presented in Figure 49.

EXAMPLE 32

CONSTRUCTION OF 2H7 scFv IgG FUSION PROTEINS WITH HINGE MUTATIONS

A 2H7 scFv IgG fusion proteins are constructed with the first cysteine residue and the second cysteine in the IgG1 hinge region substituted with a serine residue to provide MTH (SCC) and MTH (CSC). The template for introduction of the mutations is a polynucleotide encoding 2H7 scFv WTH WTCH2CH3. The oligonucleotide introducing the mutations are 5' PCR primer oligonucleotides HIgGMHcys1 (SEQ ID NO: 140) and HIgGMHcys2 (SEQ ID NO: 141). The constructs are prepared as described previously. The encoding polynucleotides of the mutants are presented in SEQ ID NOs: 163 and 165) and the polypeptide sequences are provided in SEQ ID NOs: 164 and 166).

EXAMPLE 33

CONSTRUCTION OF 2H7 VHL11S scFv (SSS-S)H WCH2 WCH3

A change from leucine to serine at position 11 in the heavy chain variable region (numbering according to Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th ed. Bethesda, MD: Public Health Service, National Institutes of Health (1991)) was introduced into the 2H7 scFv MTH (SSS) WTCH2CH3 fusion protein (SEQ ID NO: 58). The wild type leucine residue was substituted with serine by site-directed mutagenesis using the oligonucleotide Vhser11: 5'-gga ggt ggg agc tct cag gct tat cta cag cag tct ggg got gag tcg gtg agg cc-3' (SEQ ID NO: 577). The 3'-primer for PCR was huIgG1-3' having the sequence 5'-gtc tct aga cta tca ttt acc cgg aga cag-3' (SEQ ID NO: 578) (XbaI site underlined and italicized). After PCR amplification, the fragments were inserted into the TOPO® cloning vector and sequenced to confirm the presence of the V_H11 leucine to serine mutation. The 2H7 scFv-IgG (SSS-S)H WCH2 WCH3 encoding DNA was shuttled into the PSL1180 cloning vector (Pharmacia Biotech, Inc., Piscataway, NJ). The

construct PSL1180-2H7 scFv-IgG (SSS-S)_H WCH2 WCH3 was digested with Sac and XbaI to remove the wild type V_H domain and the connecting region and CH2 and CH3 domains. The PCR product comprising the VH11 mutant was digested with Sac and XbaI and then inserted into the digested PSL1180 construct according to standard molecular biology procedures. The construct was then digested with Hind III and XbaI, and inserted into the mammalian expression vector pD18 (see methods described in Example 1 and Example 10). The mutant is designated 2H7 scFv V_H L11S IgG (SSS-S)_H WCH2 WCH3. The polynucleotide sequence is provided in SEQ ID NO: 369, and the encoded polypeptide sequence is provided in SEQ ID NO: 370.

EXAMPLE 34

EXPRESSION AND OF 2H7 scFv V_H L11S (SSS-S)_H WCH2 WCH3 IN STABLE CHO LINES

CHO DG44 cells were transfected by electroporation with approximately 150 micrograms of linearized expression plasmid encoding the 2H7 V_H L11S scFv (SSS-S)_H WCH2 WCH3. Cultures were plated in selection media containing 100 nM methotrexate, in 96 well, flat bottom tissue culture plates at various numbers of cells/well, ranging from 125 to 2000. Methotrexate resistant clones were selected and culture supernatants were screened for the highest expressors of the fusion protein using a CD20CHO binding assay similar to that described for Figure 1. Clones were amplified after the initial selection in gradually increasing doses of methotrexate. Cells were passaged for two passages in the higher concentration prior to adjusting the concentration to the next higher dose. Clones were amplified to a final concentration of 1 micromolar methotrexate.

Figure 50B illustrates the production levels of 2H7 V_H L11S scFv (SSS-S)_H WCH2 WCH3. Spent supernatants from amplified CHO cells expressing this molecule and growing in stationary T25 flasks were tested for quantitative binding to CD20 CHO cells by flow cytometry. The activity was converted to protein concentration by generation of a standard curve using the same molecule purified from supernatants with Protein A affinity chromatography (Figure 50A). The concentration of the purified protein was determined by A280 using an extinction coefficient provided by the amino acid composition of the recombinant protein, (Vector NTI). Although levels of production varied between clones tested, multiple

clones produced over 1 mg/ml. This level of protein expression is over 10-fold higher than the identical molecule except for the amino acid change in V_H.

Figure 51 illustrates the production levels of 2H7 V_H L11S scFv (SSS-S)_H WCH2 WCH3 by semi-quantitative analysis on SDS-PAGE. Ten microliters of spent supernatant from amplified CHO cells expressing this molecule and growing in stationary T25 flasks were mixed with 10 microliters 2x non-reducing SDS sample buffer, run on SDS-PAGE gels, and stained with coomassie blue.

EXAMPLE 35

CONSTRUCTION AND BINDING CAPACITY OF G28-1 scFv Ig CONSTRUCTS

Construction of the G28-1 (anti-CD37) scFv was performed using total RNA isolated from the G28-1 hybridoma using TRIZOL® (Invitrogen) reagent according to manufacturer's instructions. cDNA was prepared using random primers and the protocol described previously for 2H7 cloning in Example 1. The variable domains of the scFv was cloned using one of two methods: the first method used a family of degenerate 5' oligonucleotides specific for each V region gene family and a single 3' primer specific for the constant region of either the light or heavy chain using methods and primers described in (Ig-Prime Kit Mouse Ig-Primer Set, Novagen). The second approach used the anchor-tailing methods and primers described in (Gilliland, LK et al, Tissue Antigens 47: 1-20 (1995)). In either case, PCR amplified products were cloned into the TOPO® cloning vector (Invitrogen). The clones were digested with EcoRI and screened for inserts of the proper size. Positive clones were sequenced as previously described in Example 1.

Specific primers were then designed for each V region, one with the leader sequence and one without. Primers were also designed to include desired linkers and/or restriction sites at the primer ends. PCR reactions were performed on the TOPO cloned DNA using a 25 cycle program with the following profile: 94 C, 30 sec; 55 C, 30 sec; 72 C, 30 sec, followed by a final extension at 72 C for 8 minutes. PCR products were gel purified and fragments recovered using a QIAQUICK gel extraction kit (QIAGEN, Valencia, CA). Fragments were diluted 1:50 and 1 microliter used for SEWING PCR reactions according to the methods described in Example 1. The following oligonucleotides were used for the secondary PCR reactions of the V_L domain for the G28-1 scFv:

5' primers with SalI site without leader: 5'-
GTTGTTGTGTCGACATCCAGATGACTCAGTC TCCA-3' (SEQ ID NO: 579)

5' primer with HindIII site and leader sequence: 5'-
GTCAAGCTTGCCGCCATGGTATCCA CAGCTCAGTTCCTTGG-3' (SEQ ID
5 NO: 580)

3' primer: 5'-
GCCACCCGACCCACCAACCGCCCGAGCCACCGCCACCTTTGATCTCCA
GTTCGGTG CC-3'(SEQ ID NO: 581)

The primers used for the VH domain are shown below:

10 5' sense primer:
TCGGGCGGTGGTGGGTCGGGTGGCGGCGGATCGTCAGCGGTCCAGCTGCA
GCAGTCTGGA-3' (SEQ ID NO: 582)

3' antisense primer with BclI site: 5'-
TCAGTGCTGATCAGAAGAGACGGTGACTGAGGTTTCCTTG- -3' (SEQ ID NO:
15 583)

A change from leucine to serine at position 11 in the heavy chain variable region (Kabat numbering) was introduced in into the G28-1 scFv by site-directed mutagenesis. The wild type form of the G28-1 scFv was initially constructed by sewing/overlap PCR to insert a (gly4ser)₃ linker between the V_L and V_H domains
20 as described above. However, no Sac I site was introduced as a part of this fusion of the variable domains, so alternative, nearby restriction sites (HaeII and PvuII) near leucine 11 were used to synthesize the VL+mutated V_H domain. Primers were designed to contain one of these sites and the DNA sequence including the L to S change, followed by 12 wild type base pairs. Several attempts at this strategy failed,
25 so an alternative strategy using the Genetailor (Invitrogen) method of site directed mutagenesis was used to introduce the desired mutation. The mutagenesis was carried out according to manufacturer's instructions. Briefly the procedure involves methylation of the plasmid DNA with DNA methylase, amplification of the DNA in a mutagenesis reaction with two overlapping primers, one of which contains the target
30 mutations, trasformation of the plasmid into wild type E. coli which digests all methylated DNA and leaves only the unmethylated, mutated amplification product. Both primers are approximately 30 nucleotides in length (not including the mutation

site on the mutagenic primer, with an overlapping region at the 5' ends of 15-20 nucleotides, for efficient end-joining of the mutagenesis product. The template for the mutagenesis reaction was 100 ng of a plasmid containing the wild type G28-1 scFvIg construct, and the primers used for the G28-1 V_H mutagenesis are as follows:

5 Forward primer:

5'-GCAGCAGTCTGGACCTGAGTCGGAAAAGCCTG-3' (SEQ ID
NO: 584)

Reverse Primer:

5'-CTCAGGTCCAGACTGCTGCAGCTGGAC- CGC-3' (SEQ ID
10 NO: 585)

PCR reactions were performed using the 15 ng methylated template, the primers above, and the usual reaction components as previously described. A 20 cycle program with the following profile was used for amplification: 94 C, 30 sec; 55 C, 30 sec; 68 C, 8 min, followed by a final 68 C extension step for 10 minutes. PCR
15 products were transformed into wild type bacteria, and colonies screened by sequencing. Clones with only the desired mutation were isolated and plasmid prepared as previously described Example 33. The mutant is designated G28-1 scFv VH L11S (SSS-S)H WCH2 WCH3. The polynucleotide sequence is provided in SEQ ID NO: 323, and the encoded polypeptide sequence is provided in SEQ ID NO: 324.

20 The expression level of G28-1 fusion proteins was confirmed using Immunoblot analysis according to the methods described in Example 17. Figure 53 illustrates a large increase in protein expression in the VH L11S mutant G28-1 fusion proteins compared to the G28-1 fusion protein without the mutation.

The G28-1 scFv Ig fusion proteins were transiently transfected and
25 expressed in COS cells according to methods described in Example 10. Figure 52 illustrates the capacity of the G28-1 scFv Ig fusion proteins from the COS supernatants to bind CD37. Ramos and BJAB cells both express human CD37, and were therefore used to screen the G28-1 supernatants for functional activity. Binding of G28-1 scFv (SSS-S)H WCH2 WCH3 and G28-1 scFv VH L11S (SSS-S)H WCH2
30 WCH3 to CD37+Ramos cells was measured by flow cytometry according to methods described in Example 2. Each point on the graph represents the mean of five replicate

transfections. The graph illustrates that the G28-1 scFv V_H L11S (SSS-S)H WCH2 WCH3 is able to bind CD37+Ramos cells.

5 Addition constructs were made with different connecting regions. The pD18 G28-1 scFv VHL11S (SSS-S)H WCH2 WCH3 vector was digested with BclI and XbaI to remove the connecting region, CH2 and CH3. These were replaced with each different connecting region, CH2 and CH3 according to the methods described in Example 13. The new constructs were designated: G28-1 scFv VHL11S (CSS-S)H WH2 WH3 (SEQ ID NO: 325), G28-1 scFv VHL11S (CSC-S)H WH2 WH3 (SEQ ID NO: 327), G28-1 scFv VH L11S (SSC-P)H WH2 WH3 (SEQ ID NO: 329), G28-1
10 scFv VHL11S (SCS-S)H WH2 WH3 (SEQ ID NO: 373), G28-1 scFv VHL11S(CCS-P)H WH2 WH3 (SEQ ID NO: 375), and G28-1 scFv VHL11S (SCC-P)H WH2 WH3 (SEQ ID NO: 377).

The G28-1 scFv was also attached to an IgA connecting region, CH2, CH3 and an IgE CH2, CH3, CH4. The pD18 G28-1 scFv VHL11S (SSS-S) H WCH2
15 WCH3 plasmid was digested using methods above to remove the connecting region CH2, and CH3. The IgA regions were inserted using methods described in Example 13. The construct was designated G28-1 scFv VHL11S IgAH IgACH2 T4CH3 (SEQ ID NO: 381). The IgE CH2 CH3 CH4 region was inserted into the digested pD 18 vector above using methods described in Example 39. The construct was designated
20 G28-1 scFv VHL11S IgECH2 CH3 CH4 (SEQ ID NOs: 379 and 383).

EXAMPLE 36

CHARACTERIZATION OF 2H7 scFv Ig MUTANT FUSION PROTEINS

Figure 54 illustrates the binding capacity of purified 2H7 scFv Ig constructs to CD20+CHO cells. The proteins were transfected into stable CHO cells
25 according to methods described in Example 2. Binding was determined using flow cytometry according to the methods described in Example 2. The graph in Figure 54 illustrates that these proteins retain binding function to CD20 with altered connecting regions. Comparative results were obtained in each of the 2H7 scFv VHL11S mutants with each type of altered connecting region (results omitted).

30 The ability of 2H7scFv-Ig constructs with mutated connecting regions to kill CD20 positive cells in the presence of peripheral blood mononuclear cells (PBMC) through ADCC was tested by measuring the release of ⁵¹Cr from labeled

BJAB cells in a 4 hr. assay using 100:1 ratio of PBMC to BJAB cells. The results shown in Figure 55 indicate that 2H7scFv-Ig mutants can mediate antibody dependent cellular cytotoxicity (ADCC), since the release of ⁵¹Cr was significantly higher in the presence of both PBMC and 2H7scFv-Ig than in the presence of either PBMC or
5 2H7scFv-Ig alone. Comparative results were obtained in each of the 2H7 scFv VHL11S mutants with each type of altered connecting region (results omitted).

The ability of 2H7scFv-Ig mutant fusion proteins to kill CD20 positive cells in the presence of complement was tested using B cell lines Ramos target cells. Rabbit complement was purchased from Pel-Freez (Rogers, AK), and was used in the
10 assay at a final concentration of 1/10. Purified 2H7scFv-Ig was incubated with B cells and complement for 45 minutes at 37° C, followed by counting of live and dead cells by trypan blue exclusion. The results in Figure 56 show that 2H7scFv-Ig mutants in the presence of rabbit complement, lysed B cells expressing CD20.

EXAMPLE 37

15 COMPARATIVE BINDING OF IGA, IGG, AND IGE 2H7 SCFV CONSTRUCTS

Binding capacity of Ig constructs IgA, IgG and IgE were measured using flow cytometry according to the methods described in Example 2, using a commercially available (Caltag) second step specific for each Ig tail. The results in Figure 57 show that all the IgE constructs were able to bind CD20+CHO cells
20 comparable to the binding abilities of IgG and IgA. These results also demonstrate that the IgE constructs were detected with the IgE second step, but not the IgA or IgG second step.

EXAMPLE 38

CONSTRUCTION AND CHARACTERIZATION OF 2H7 VHL11S IGE CONSTRUCTS

IgE tail RNA was isolated from SKO-007 cells(ATCC) using
25 QIAGEN QIAshredder homogenization and RNA minikits. Random-primed cDNA was generated according to the usual protocol, with 4 microliters RNA eluted from the QIAGEN columns. Human IgE from the beginning of CH1 through CH4 (approximately 1.2 kb) was isolated by PCR amplification of 5 microliters cDNA,
30 with an amplification profile of 94 C, 60 sec; 72 C, 2 minutes for 35 cycles, and the following primers:

5' primer: 5'-ggatccacccgctgctgcaaaaacattccctccaatgccacctccgtgac-3'
(SEQ ID NO: 586)

3' primer: 5'-tcattaccgggatttacagacaccgctcgctggacggctgtgaggggctcg-
ctgc-3' (SEQ ID NO: 587)

5 PCR fragments were ligated into PCR 2.1-TOPO vector, and
transformants screened for inserts of the correct size by digestion with EcoRI
according to the methods described in Example 1. One of the clones with the correct
sequence from was used as template to amplify the CH2-CH4 domains with
appropriate restriction sites attached for subcloning as soluble or cell surface (ORF)
10 forms. The following primers were used with an amplification profile of 94 C, 60 sec;
55 C, 60 sec; 72 C, 2 min; for 35 cycles to amplify a fragment of approximately 950
bp:

5' primer: (attaches BclI site to 5' end of CH2 domain of IgE)

5'-gttggtgatcacgtctgctccaggga- cttcacc-3' (SEQ ID NO: 588)

15 3' primer: (attaches, stop codon and XbaI site to 3' end of CH4 of IgE)

5'- gttgtttctagattatcatttaccaggatttacagacaccgctcgctg- 3' (SEQ ID NO:
589)

3' primer: (attaches SfuI and BamHI to 3' end of CH4 without a stop
codon)

20 5'- gttgttttcgaaggatccgctttaccagatttacagacaccgctcgctg- 3' (SEQ ID NO:
590)

The IgE CH2CH3CH4 tail with a stop codon was digested with BclI
and XbaI and inserted into a pD18 vector that contains 2H7 VHL11S scFv. This
construct was designated 2H7 IgECH2CH3CH4. The polynucleotide sequence is
25 provided in SEQ ID NO: 128, and the encoded polypeptide sequence is provided in
SEQ ID NO: 129. The IgE CH2CH3CH4 tail with no stop codon (ORF) (SEQ ID NO:
96) was digested with BclI and SfuI and inserted in into a pD 18 vector that contains
2H7 VHL11S scFv. This construct was designated 2H7 IgECH2CH3CH4(ORF).

30 The human IgE was also amplified as a fragment missing both CH1
and CH2 domains, with only the CH3 and CH4 domains attached to the human IgG1
hinge. Sequential PCR reactions using overlapping 5' oligonucleotides were used to

attach the IgG 1 hinge to the CH3 domain of human IgE. Primers for the first step of the PCR reaction:

5' Primer: 5'- actcacacatccccaccgtccccagcatccaacccgagaggggtgagc- 3'
(SEQ ID NO: 591)

5 Primers for the second step of the PCR reaction:

5' primer: 5'- tctgatcaggagcccaaattcttct- gacaaaactcacacatccccaccg- 3'
(SEQ ID NO: 592)

3' primer: 5'- gttgtttctagattatcatttaccaggatttacaga- caccgctcgctg- 3'
(SEQ ID NO: 593)

10 The PCR product was digested with EcoRI and sequenced according to the methods described in Example 1. Positive clones were inserted into pD18 plasmid containing 2H7 VHL11S scFv (SSS-S)H. The construct was designated 2H7 VHL11S scFv (SSS-S)H IgE CH3CH4. The polynucleotide sequence is provided in SEQ ID NO: 227, and the encoded polypeptide sequence is provided in SEQ ID NO: 228.

15 Binding capacity of 2H7 scFv VH L11S IgECH2 CH3 CH4, was measured using flow cytometry, essentially according to Example 2. The protein was purified using MEP HyperCel, (Cipergen, Catalog # 12035-010, Lot# 200920/0271) chromatography resin and Hydrophobic charge induction chromatography (HCIC). HCIC absorbent is a high capacity, high selectivity, absorbent designed for capture
20 and purification of monoclonal and polyclonal antibodies from various sources including cell culture supernatants. Columns were packed with a 10 ml bead volume of MEP Hypercel, and equilibrated with PBS, pH 7.4 containing 0.1% NaN3. Approximately 1 liter of 2H7 scFv VHL11S IgE CH2CH3CH4 CHO culture supernatant was then run over the column. A series of citrate buffers ranging from pH
25 3-6 were prepared for elution of the fusion protein. The column was washed in PBS. Protein was eluted in fifteen 1 ml fractions at pH6, 5, and 4. A final 15 ml fraction was collected at pH 3.5. Aliquots from each fraction were analyzed for A280 and were also subjected to SDS-PAGE, loading roughly 10 micrograms/well based on the A280 reading. The results of these two analyses indicated that the bulk of the protein
30 did not elute in citrate buffers at the higher pH, but eluted at pH4, and the post elution wash at pH3.5 also contained significant amounts of protein.

The ability of these 2H7 V_H L11S IgE purified proteins to bind CD20+CHO cells was determined using flow cytometry according to the methods described in Example 2 using FITC-conjugated goat-anti-human IgE. Figure 58A illustrates that both purified proteins are able to bind CD20+CHO cells.

5 The ability of these 2H7 V_H L11S IgE purified proteins to mediate ADCC against BJAB target cells with PBMC effectors was measured according to the methods described in Example 2. Figure 58B illustrates that both proteins were able to mediate ADCC at similar levels.

EXAMPLE 39

10 **CONSTRUCTION AND BINDING CAPACITY OF SCFV V_H L11S MUTANTS WITH MOUSE IGA AND IGE TAIL REGIONS**

Murine IgA was cloned from murine spleen RNA using essentially the same methods used to clone the human IgE tails in Example 38. The PCR reactions were performed with a 94 C 60 sec; 52 C 60 sec; 72 C 2 min amplification profile for 15 35 cycles. The PCR primers used to clone CH1-CH4 regions were:

5' primer: 5'-atctgttctctctactactctctctccacct-3' (SEQ ID NO: 594)

3' primer: 5'-tcagtagcagatgccatctccctc- tgacatgatgacagacacgct-3' (SEQ ID NO: 595)

PCR primers used to delete the CH1 region:

20 5' primer: 5'-gttggtgatcacatctgttctctctactactctctctccacct-3' (SEQ ID NO: 596)

3' primer with a stop codon, XbaI site at end of Ig tail, and the T4 mutation in CH3 region:

5'-gttggttctagattatcaatctc- cctctgacatgatgacagacac-3' (SEQ ID NO: 597)

25 3' primer for the ORF, a SfuI and BamHI sites, and T4 mutation in the CH3 region:

5'-gttcttcgaaggatccgcacatctccctctgac- atgatgac-3' (SEQ ID NO: 598)

The mouse IgACH2 T4CH3 tail with a stop codon was digested with BclI and XbaI and inserted into a pD18 vector that contains 2H7 V_HL11S scFv and 30 the IgAH. This construct was designated 2H7 V_HL11 S scFv IgAH mIgACH2

T4CH3. The polynucleotide sequence is provided in SEQ ID NO: 253, and the encoded polypeptide sequence is provided in SEQ ID NO: 254.

The mouse IgACH2 T4CH3 tail with no stop codon (ORF) was digested with BclI and SfuI and inserted in into a pD 18 vector that contains 2H7 VHL11S scFv and IgAH. This construct was designated 2H7 VHL11S scFv IgAH mIgACH2 T4CH3 (ORF). The polynucleotide sequence is provided in SEQ ID NO: 255, and the encoded polypeptide sequence is provided in SEQ ID NO: 256.

Murine IgE was cloned murine IgE La2 (ATCC)RNA essentially according the methods described in Example 38. The PCR reactions were performed with a 94 C 60 sec; 52C 60 sec; 72 C 2 min amplification profile for 35 cycles. Initial PCR primers used to clone the CH1--CH4:

5' primer: 5'-tctatcaggaaccctcagctctacccttgaa- gccctg-3' (SEQ ID NO: 599)

3' primer: 5'-gtgttttagattatcaggatggacggaggagggtgttaccaggct-3' (SEQ ID NO: 600)

PCR primers to remove the CH1 region:

5' primer: 5'-gtgttgatcacgttcgacctgtcaacatcac- tgagcccacc-3' (SEQ ID NO: 601)

3' primer with stop codon and XbaI site:

5'-gtgttttagattatcaggatggacggaggagggtgttaccaggct-3' (SEQ ID NO: 602)

3' primer ORF, SfuI and Bam HI: 5'-gtgttttcgaaggatccgcggatggacggaggagggtgtta-3' (SEQ ID NO: 603)

The mouse IgE CH2CH3CH4 tail with a stop codon was digested with BclI and XbaI and inserted into a pD 18 vector that contains 2H7 VHL11S scFv. This construct was designated 2H7 VHL11S mIgECH2CH3CH4. The polynucleotide sequence is provided in SEQ ID NO: 249, and the encoded polypeptide sequence is provided in SEQ ID NO: 250.

The mouse IgE CH2CH3CH4 tail with no stop codon (ORF) was digested with BclI and SfuI and inserted in into a pD18 vector that contains 2H7

VHL11S scFv. This construct was designated 2H7 VHL11S scFv mIgE_{CH2CH3CH4}(ORF). The polynucleotide sequence is provided in SEQ ID NO: 249, and the encoded polypeptide sequence is provided in SEQ ID NO: 250.

5 Binding capacity of 2H7 V_H L11S mIgE and mIgA (with mouse tail regions) to CD20+CHO cells were also measured by flow cytometry according to the methods described in Example 2 using commercially available IgE or IgA second step reagents (Caltag). Each point in Figure 59 represents the mean of a population with brightness corrected by subtracting the binding of the second step alone. This figure illustrates that these constructs have the ability to bind CD20+CHO cells.

10

EXAMPLE 40

HPLC PROFILES OF 2H7 scFv-Ig MUTANT FUSION PROTEINS

HPLC analysis of purified 2H7 scFv-Ig mutant fusion proteins with altered connecting and CH3 regions. Each protein was purified by Protein A affinity chromatography from supernatants of transfected COS or CHO cells. Twenty-five to
15 fifty micrograms of each sample was run at 1 m/min in PBS on a TSK-GEL G3000S W_{XL} 30 cm column (Tosoh Biosep, Stuttgart, Germany). The arrow near the beginning of each profile represents the sample injection point. Gel filtration standards (Bio-Rad) included thyroglobulin (670 kDa), gamma globulin (158 kDa), ovalbumin (44 kDa), myoglobin (12.5 kDa), and vitamin B-12 (1.35 kDa). Standards
20 were run at the beginning and end of each experiment. Migration positions of standards are shown and did not vary between experiments. (FIGS. 60-62)

EXAMPLE 41

BINDING CAPACITY OF 2H7 V_H L11S MUTANT FUSION PROTEINS

Binding effects of the CH3 mutant were compared to the non-mutated
25 CH3 fusion protein. The constructs were transfected into COS cells and purified from the supernatant using protein A column purification techniques described in Example 2. Figure 63 illustrates the differential effects of CH3 mutations on binding using flow cytometry according to the methods described in Example 2. This figure illustrates that some binding ability is lost when the double point mutation is introduced in the
30 CH3 region.

Binding of fluorescein isothiocyanate (FITC) conjugated 2H7 V_H L11S mutant fusion proteins was determined using flow cytometry. A 1 mg/ml solution of FITC was prepared in DMSO. Fusion proteins were dialyzed in pH 9.3 bicarbonate buffer overnight at 4C in a volume of 2 liters. Concentration of the protein was adjusted to 1-5 mg/ml prior to conjugation. A series of Falcon 5 ml tubes was set up with varying FITC to protein ratios, ranging from 15-60, but minimally ratios of 20 and 40. Conjugation reactions were incubated at 37C for 30 minutes protected from light. Fluorescein labeled protein was separated from free fluorescein on a 2 ml Sephadex G-25 column equilibrated with PBS, 0.5 M NaCl, and 1% NaN₃. The fluorescein labeled protein eluted from the column first and was collected in a 5 ml tube. The degree of labeling was determined by measuring the absorbance of the diluted conjugate at 280 and 494 nm, and utilizing the formulas provided by technical services at Molecular Probes (Eugene, OR). Data has been corrected for FITC: protein ratio. Figure 64 illustrates that these constructs do not lose binding capacity when conjugated to a fluorescent marker.

Purified 2H7 V_H L11S constructs were run on a non-reducing SDS gel according to the methods described in Example 2. The migration patterns are presented in Figure 65.

EXAMPLE 42

CHARACTERIZATION OF 2H7 scFv V_H L11S(CSC-S)H WCH2 WCH3 IN LEC13 CHO CELLS

2H7 scFv V_H L11S (CSC-S)H WCH2 WCH3 were transiently transfected and expressed in Lec13 CHO cells. Lec13CHO cells were used as the mammalian cell hosts for either the 2H7 scFv VHS11 hIgG1 (CSS-S)H WCH2 WCH3 and (CSC-S)H WCH2 WCH3 expressing plasmids in side-by-side transfections. All transfections were performed in 100 mm tissue culture dishes. Cells were transfected when approximately 90% confluent using lipofectamine 2000 (Invitrogen, Catalog #: 11668-027, 0.75 ml), following manufacturer's instructions. Both cell lines were grown in the presence of serum to promote and maintain adherence to the cell culture dishes, simplifying transfection manipulations, washes, and supernatant harvests. DNA: lipofectamine complexes were allowed to form in the absence of serum and antibiotics, following the suggested protocol/conditions recorded in the product insert. Culture supernatants were harvested 72 hours after

transfection, and then again 72 hours after the first harvest. Fusion protein from the two CHO sources was isolated by protein A purification as previously described and used in CD20 binding and ADCC assays.

5 The ability of mutated fusion protein to mediate ADCC in CD20 positive cells was determined using the methods described in Example 2. Constructs expressed in Lec13 CHO cells exhibited better binding to CD20 CHO target cells and also showed significantly improved activity in ADCC assays relative to the CHO DG44 derived proteins at equivalent concentrations as illustrated in Figure 67.

EXAMPLE 43

10 CONSTRUCTION OF HIGH AND LOW AFFINITY CD16 ALLELES

The low(V) and high(F) affinity alleles at position 158 of the human CD16 extracellular domain were cloned from cDNA derived from human PBMC using PCR assay. PCR reactions used random primed cDNA made from PBMC stimulated for 3 days with immobilized anti-CD3 antibody (64.1) prior to harvest.

15 PCR reactions included 2, 4, 6 or 8 microliters of cDNA, each primer at 25 pmol, and an amplification profile of 94 C 60 sec; 55 C 60 sec; 72 C 2 min, for 35 cycles. The PCR primers are listed below:

5' primer - no leader peptide: 5'-
GTTGTTACCGGTGCAATGCGGACTGAAGATCTCCC

20 AAAGGCTGTG-3' (SEQ ID NO: 604)

3' antisense primer: 5'-
GTTGTTTGATCAGCCAAACCTTGAGTGATGGTGATGTTTACA-3' (SEQ ID NO: 605)

Positive clones were sequenced, and inserted into a vector containing

25 an efficient leader peptide and the (SSS-S)H P238SCH2 WCH3 human IgG tail. Two different versions of the CD16 ED fusion proteins were expressed. The first contained the F 158 (high affinity) and the second contained the V158 (low affinity) allele. Constructs were cloned into a (SSS-S)H P238S CH2 WCH3 pD18 plasmid and expressed in COS and CHO cells as previously described in examples 1 and 10. CHO

30 clones were screened for expression using an IgG sandwich ELISA to determine relative expression levels of the fusion proteins in the culture supernatant using the

following protocol: Immulon 1V plates were coated at 4C with 0.4 mg/ml goat anti-human IgG (mouse Adsorbed), (CalTag, Catalog # H10500) in PBS buffer. Plates were then blocked in PBS/1.5% nonfat milk at 4C overnight. Plates were washed three times in PBS/0.1% Tween 20, then incubated with 100 microliters dilution series from CHO clone culture supernatants at room temperature for 3 hours. Four dilutions per clone were added to successive wells, diluting in 5 fold increments from 1:5 to 1:375. In addition a standard curve was derived using CTLA4 hIgG1 (SSS-S)H P238SCH2 WCH3 as a concentration standard. The dilution series utilized 5 fold dilutions starting at 0.5 micrograms/ml; a second set of 8 wells was used to make a 2-fold dilution series starting at 0.34 micrograms/ml. Plates were washed 3 times in PBS/0.1% Tween 20, and incubated with goat anti-human IgG, conjugated to horseradish peroxidase (GAH IgG-HRP) at 1:5000, in PBS/0.5% BSA for 1 hour. Plates were washed four times with PBS/0.1% Tween 20, then TMB chromagen substrate (BD-Pharmingen) was added for 10 minutes, and reactions stopped by addition of 100 microliters IN sulfuric acid. Plates were then read at 415 nm on a SpectraCount plate reader. Concentrations of fusion protein were estimated by comparison of the ODs in the linear range to the CTLA41g standard curve run on each plate.

Proteins were purified using Protein A purification and were directly conjugated to fluorescein isothiocyanate (FITC) as described in Example 42. These proteins were run out on SDS gels under reduced and nonreduced conditions according to the methods described in Example 2. The migration of these proteins is presented in Figure 68.

The ability of the high and low CD 16 alleles to bind 2H7 VH111S(CSC-S)H WCH2 WCH3 or bind 2H7 VH111S (SSS-S)H(P238S)CH2 WCH3 expressed on the cell surface of CD20+CHO target cell is determined using flow cytometry according to the methods described in Example 2. The results in Figure 69 demonstrate both the high and low affinity alleles were able to bind 2H7 VHL11S(CSC-S)H WCH2 WCH3 (SEQ ID NO: 246) and lost some binding capabilities when the P238S mutation was introduced into the CH2 region of the construct (SEQ ID NOs: 417 and 418).

EXAMPLE 44

MAMMALIAN DISPLAY SYSTEM

Figure 70A diagrams how FITC conjugates of FcRIII (CD16) soluble fusion proteins bind to 2H7 scFv-Ig constructs that are attached to CD20 expressed by CHO cells. The CD16 binding to a scFv-Ig provides a screening tool for detecting changes in CD16 binding to an altered scFv-Ig constructs containing targeted or site-specific mutations. Changes in CD 16 binding properties may be changes in binding of either CD 16 high affinity protein (158F) or CD16 low affinity protein (158 V) or both.

A schematic representation of such a screening process is diagrammed in Figure 70B, where scFv-Ig constructs are displayed on the cell surface of mammalian cells. The scFv-Ig molecules in this example are displayed on the cell surface because they contain a transmembrane domain anchor. These molecules may represent a single scFv-Ig construct or may be introduced into a population of mammalian cells as a library of such molecules. Transfected cells with altered binding properties can then be panned, sorted, or otherwise isolated from other cells by altering the stringency of the selection conditions and using CD16 fusion proteins as the binding probe. Cells that express scFv-Ig molecules with altered binding to either CD16 high affinity allele (158F) or CD16 low affinity allele (158V) or both can be isolated. For example, this display system can be used to create a library of mutated Ig tails with short stretches of CH2 sequence replaced with randomized oligonucleotides or possibly randomization of a single residue with all possible amino acid substitutions, including synthetic amino acids. Once such a library is constructed, it can be transfected into COS cells by methods well known in the art. Transfectants can then be bound to the labeled CD 16 constructs, and panned or sorted based on their relative binding properties to multiple allelotypes/isoforms. Panned cells are harvested, and the plasmid DNA is isolated and then transformed into bacteria. This process may be repeated iteratively multiple times until single clones are isolated from the mammalian host cells (see Seed B and Aruffo A, PNAS 84: 3365-3369 (1987) and Aruffo A and Seed B, PNAS 84: 8573-8577 (1987)). One such use of this type of screening system would be to isolate Ig tails which bind equally well to both the high and low affinity alleles of CD16 with the goal of improving effector functions mediated by scFv-Ig constructs in multiple subpopulations of patients. Ig

tails with altered binding properties to other Fc receptors can also be selected using the display system described. Other display systems for example those that use bacteriophage or yeast are not suitable for selection of Ig tails with altered FcR binding properties because of the requirement for glycosylation in the Ig CH2 domain
5 that would not occur in non-mammalian systems.

This system is also useful for selection of altered scFv-Ig molecules that will be produced at higher levels. In this example, mammalian cells such as COS cells can be transfected with a library of scFv-Ig constructs in a plasmid that directs their expression to the cell surface. COS cells that express the highest levels of the
10 scFv-Ig molecules can be selected by techniques well known in the art (for example panning, sterile cell sorting, magnetic bead separation, etc), and plasmid DNA is isolated for transformation into bacteria. After several rounds of selection single clones are isolated that encode scFv-Ig molecules capable of high level expression. When the isolated clones are altered to remove the membrane anchor and then
15 expressed in mammalian cells, the scFv-Ig constructs will be secreted into the culture fluid in high levels. This reflects the common requirement of secreted glycoproteins and cell surface glycoproteins for a signal peptide and processing through the golgi for expression, so that selection for a molecule that illustrates an improvement in expression levels on the cell surface will also select for a molecule that illustrates an
20 improvement in levels of secreted protein.

EXAMPLE 45

CHARACTERIZATION OF G28-1 MABS AND scFVS

Ability of G28-1 mAbs and scFvs to induce apoptosis was measured by binding Annexin V, using the methods described in Example 3. The results in
25 Figure 71 demonstrate that the Annexin V binding of G28-1 antibodies and scFv is increased when treated together with 2H7 antibodies and scFv constructs.

EXAMPLE 46

CONSTRUCTION OF FC2-2 scFV CONSTRUCTS

Contrstruction of the FC2-2 (anti-CD16) scFv was performed using
30 total RNA isolated from the FC2-2 hybridoma and cloned using methods described in Example 35. The polynucleotide sequence is provided in SEQ ID NO: 337, and the encoded polypeptide sequence is provided in SEQ ID NO: 338. The specific primers

for the secondary PCR reaction are listed below. The following are primers for the light chain variable region:

5' primer with HindIII site with no leader: 5'-
GTTGTTAAGCTTGCCGCCATGGATTAC

5 AGGCCCAGGTTCTT-3' (SEQ ID NO: 606)

5' primer with SalI site and leader: 5'-
GTTGTTGTCGACATTGTGATGTCACAGTCTCC

ATCCTCCCTA-3' (SEQ ID NO: 607)

3' primer:

10 5'-TCAGTGCTGATCATGAGGAGACGGTGACTGAGGTTCTT-3'
(SEQ ID NO: 608)

The following are primers for the heavy chain variable region:

5' primer: 5'-

15 TCGGGCGGTGGTGGGTCGGGTGGCGGCGGATCGTCACAGGT
GCAGTTG

AAGGAGTCAGGA-3' (SEQ ID NO: 609)

3' primer: 5'-

ACCCGACCCACCAACCGCCCGAGCCACCGCCACCTTTTATT- TCCAGCTTG
GTGCCACCTCCGAA-3' (SEQ ID NO: 610)

20 A change from leucine to serine at position 11 in the heavy chain
variable region (Kabat numbering) was introduced into the FC2-2 scFv by site-
directed mutagenesis according to the methods described in Example 33. The scFv
was attached to the (SSS-S)H WCH2 WCH3 IgG tail according to methods described
in Example 33. The mutant is designated FC2-2 scFv VH L11S (SSS-S)H WCH2
25 WCH3. The polynucleotide sequence is provided in SEQ ID NO: 345, and the
encoded polypeptide sequence is provided in SEQ ID NO: 346.

EXAMPLE 47

CONSTRUCTION OF 5B9 scFv CONSTRUCTS

Contrstruction of the 5B9 (anti-CD137) scFv was performed using total RNA isolated from the 5B9 hybridoma and cloned using methods described in Example 35. The polynucleotide sequence is provided in SEQ ID NO: 361, and the encoded polypeptide sequence is provided in SEQ ID NO: 362. The specific primers for the secondary PCR reaction are listed bellow. The following are primers for the light chain variable region:

5' primer with HindIII site with no leader: 5'-
10 GTTGTTAAGCTTGCCGCCATGAGGTTCT
CTGCTCAGCTTCTG-3' (SEQ ID NO: 611)
5' primer with SalI site and leader: 5'-
GTTGTTGTCGACATTTGTGATGACGCAGGCTG
CATTCTCCAATT-3' (SEQ ID NO: 612)
15 3' primer: 5'-
TCAGTGCTGATCAGAGGAGGACGGTGACTGAGGTTTCCTTG-3' (SEQ ID NO: 613)

The following are primers for the heavy chain variable region:

5' primer: 5'-
20 CGGGCGGTGGTGGGTCGGGTGGCGGCGGATCGTCAC- AGGTGCAGCTGA
AGCAGTCAGGA-3' (SEQ ID NO: 614)
3' primer: 5'-
CCCGACCCACCAACCGCCCGAGCCACCGCCACCCTTCAGCTCCAGCTTG
GTGCCAGCACC-3' (SEQ ID NO: 615)

25 A change from leucine to serine at position 11 in the heavy chain variable region (Kabat numbering) was introduced in into the 5B9 scFv by site-directed mutagenesis and attached to (SSS-S)H WCH2 WCH3 according to the methods described in Example 33. This construct was designated 5B9 scFv VHL11S(SSS-S)H WCH2 WCH3. The polynucleotide sequence is provided in SEQ
30 ID NO: 365, and the encoded polypeptide sequence is provided in SEQ ID NO: 366.

EXAMPLE 48

CONSTRUCTION OF UCHL1 scFv CONSTRUCTS

Contrstruction of the UCHL1 (anti-CD45RO) scFv was performed using total RNA isolated from the UCHL1 hybridoma and cloned using methods described in Example 35 The polynucleotide sequence is provided in SEQ ID NO: 351, and the encoded polypeptide sequence is provided in SEQ ID NO: 352. The following are primers for the light chain variable region:

5' primer with HindIII site: 5'-
GTTGTTAAGCTTGCCGCCATGAAGTTGCCTGTTAGGCTG
10 TTGGTGCTG-3' (SEQ ID NO: 616)
3' primer with Sac site: 5'-
AGAGCTCCACCTCCTCCAGATCCACCACCGCCCGAGCCAC
CGCCATCTTTGATTTCAGCTTGGT-3' (SEQ ID NO: 617)

The following are primers for the heavy chain variable region:

15 5' primer: 5'-
TTTCAGAGTAATCTGAGAGCTCCACCTCCTCCAGATCCACCACCGC
CCGA-3' (SEQ ID NO: 618)
3' primer: 5'-
TCAGTGCTGATCATGCAGAGAGACAGTGACCAGAGTCCC-3' (SEQ ID NO:
20 619)

A change from leucine to serine at position 11 in the heavy chain variable region (Kabat numbering) was introduced in into the 5B9 scFv by site-directed mutagenesis and connected to a(SSS-S) HWCH2 WCH3 according to the methods described in Example 33. The mutant is designated UCHL1 scFv VH L11S (SSS-S)H WCH2 WCH3. The polynucleotide sequence is provided in SEQ ID NO: 359, and the encoded polypeptide sequence is provided in SEQ ID NO: 360.

EXAMPLE 49

L6 VHL11S scFv (SSS-S)H WCH2 WCH3

A change from leucine to serine at position 11 in the heavy chain variable region (Kabat numbering) was introduced in into the L6 scFv (SSS-S)H

WCH2 WCH3 (constructed according to methods described in Example 106) by site-directed mutagenesis according to the methods described in Example 33. The L6scFvIg (SSS-S)H WCH2 WCH3 pD18 plasmid was used as template. Positive clones were inserted into the pD18 plasmid containing (SSS-S)H WCH2 WCH3 according to methods described in Example 33. The mutant is designated L6 scFv V_H L11S (SSS-S)H WCH2 WCH3. The polynucleotide sequence is provided in SEQ ID NO: 415, and the encoded polypeptide sequence is provided in SEQ ID NO: 416.

PCR primers are listed below:

5' Primer with PstI restriction site: 5'-

10 ggcggtatctctgcagatccagttggtgcagtctggacctgagtcgaagaagcct

ggagag-3' (SEQ ID NO: 620)

3' Primer: 5'-ggacagtgggagtggtgcacc-3' (SEQ ID NO: 621)

EXAMPLE 50

CONSTRUCTION OF HD37 scFv VH11S CONSTRUCT

15 A change from leucine to serine at position II in the heavy chain variable region (Kabat numbering) was introduced in into the HD37 scFv by site-directed mutagenesis according to the methods described in Example. 33. The HD37 scFv (SSS-S)H WCH2 WCH3 pD 18 plasmid was used as a template. Positive clones were inserted into the pD18 plasmid containing (SSS-S)H WCH2 WCH3 according to methods described in Example 33. The mutant is designated HD37 scFv V_H L11S (SSS-S)H WCH2 WCH3. The polynucleotide sequence is provided in SEQ ID NO: 401, and the encoded polypeptide sequence is provided in SEQ ID NO: 402. PCR primers are listed below:

25 5' primer: 5'-caggttcagctgcagcagctctggggctgagtc- ggtgaggcctgg-3' (SEQ ID NO: 622)

3' primer: 5'-ggaggattcgtctgcagtcagagtggc-3' (SEQ ID NO: 623)

EXAMPLE 51

2H7 scFv VH11S CONSTRUCTS

30 Additional 2H7 VH L11S constructs were made with different connecting regions. The pD18 2H7 scFv VHL11S (SSS-S)H WCH2 WCH3 vector

was digested with BclI and XbaI to remove the connecting region, CH2 and CH3. These were replaced with each different connecting region, CH2 and CH3 according to the methods described in Example 13. The new constructs were designated: 2H7scFv VHL11S (CSS-S)H WH2 WH3 (SEQ ID NO: 371), 2H7scFv VHL11S (CSC-S)H WH2 WH3 (SEQ ID NO: 245). 2H7 scFv VHL IS was also attached to an IgA connecting region, CH2, CH3 and an IgE CH2, CH3, CH4. The pD18 2H7 scFv VH L11S (SSS-S)H WCH2 WCH3 plasmid was digested using methods above to remove the connecting region CH2, and CH3. The IgA regions were inserted using methods described in Example 13. The construct was designated 2H7 scFv VHL11S IgAH IgACH2 T4CH3 (SEQ ID NOs: 251 and 253). The IgE CH2 CH3 CH4 region was inserted into the digested pD18 vector above using methods described in Example 39. The construct was designated 2H7 scFv VHL 1 S IgECH2 CH3 CH4 (SEQ ID NO: 247).

EXAMPLE 52

2H7 scFv VH L11S(CSC-S)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is connected to a human IgG1 connecting region, CH2 and CH3 region, where in the second cysteine and the proline in the connecting region have been changed to serines (SSS-S) as described in Example 32. The polynucleotide sequence is provided in SEQ ID NO: 245, and the encoded polypeptide sequence is provided in SEQ ID NO: 246.

EXAMPLE 53

2H7 scFv VH L11S IgE CH2 CH3 CH4

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is attached to a human IgE constant region containing CH2, CH3 and CH4 as described in Example 38. The polynucleotide sequence is provided in SEQ ID NO: 247, and the encoded polypeptide sequence is provided in SEQ ID NO: 248.

EXAMPLE 54

2H7 scFv VH L11S mIgE CH2 CH3 CH4

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is attached to a mouse IgE constant region containing CH2, CH3 and CH4 as described in Example 39. The polynucleotide sequence is provided in SEQ ID NO: 249, and the encoded polypeptide sequence is provided in SEQ ID NO: 250.

EXAMPLE 55

2H7 scFv VH L11S mIgAH WIGACH2 T4CH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a mouse IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon as described in Example 39. The polynucleotide sequence is provided in SEQ ID NO: 253, and the encoded polypeptide sequence is provided in SEQ ID NO: 254.

EXAMPLE 56

2H7 scFv VH L11S(SSS-S)H K322S CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is connected to a mutated human IgG connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The K322S mutation in the CH2 region is at a residue 322, where a Lysine has been changed to a serine using overlapping PCR assay. An (SSS-S)H WCH2 WCH3 IgG1 template in the pD18 vector was used for PCR amplification, to create (SSS-S)H derivatives containing these CH2

mutations. PCR reactions used a cycling profile of 94 C, 30 sec; 55 C, 30 sec; 72 C, 30 sec. for 37 cycles to complete the reactions. This IgG1 derivative was constructed by using sequential PCR reactions with overlapping oligonucleotides in the primary and secondary reactions. The primary amplification primers introduced the mutation(s), but deleted one end of the Fc domain. Secondary reaction primers reattached these ends using overlapping primers. The first overlapping primer was added at the beginning of the PCR, the reactions allowed to proceed for 12 cycles, paused and then the second overlapping primer added to the reactions followed by 25 more cycles to complete the overlap extension PCR reactions.

Primers for the first PCR reaction:

5' Primer: 5'-

ggagatggtttctcgcgatgggggctgggagggcttggagaccgagcactgtactcc-3' (SEQ ID NO 624)

3' primer: 5'-ggacagtgggagtggcacc-3' (SEQ ID NO 625)

PCR products were cloned into TOPO vector and sequenced for verification. Positive vectors were used as templates for the second overlapping PCR reaction.

5' primer:

5'ccgtctctgatcaggagcccaaattctctgacaaaactcacatccccaccgtccccagc-3' (SEQ ID NO 626)

5' primer overlapping primer: 5'-

tccccaccgtccccagcacctgaactcctgggggatcgtcagtcttctcttcccc
caaaacc-3' (SEQ ID NO 627)

3' primer: 5'-caggaaacagctatgac-3' (SEQ ID NO 628)

PCR product was cloned into TOPO vector and sequenced. The polynucleotide sequence is provided in SEQ ID NO: 263, and the encoded polypeptide sequence is provided in SEQ ID NO: 264.

EXAMPLE 57

2H7 scFv VH L11S(CSS-S)H K322S CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region,

where the leucine has been changed to a serine, as described in Example 33. This binding region is attached to a mutant IgG connecting region, where the second and third cysteines have been changed to serines and the proline has been changed to serine (CSS-S), according to methods described in Example 13. The connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The K322S mutation in the CH2 region is at a residue 322, where a Lysine has been changed to a serine using overlapping PCR assay. region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The mutation in the CH2 region was added by overlapping PCR reaction essentially according to Example 57, with (CSS-S)H WCH2 WHC3 IgG1 pD18 vector as a template in the first PCR reaction and different primers in the second PCR reaction, which are listed below.

5' primer: 5'-

ccgtctctgatcaggaccccaaatcttgacaaaactcacacatccccaccgtccccagc-3' (SEQ ID NO: 629)

5' overlapping primer: 5'-

15 tccccaccgtccccagcacctgaactcctggggg- gatcgtagcttctctcccccaaaacc-3'
(SEQ ID NO 630)

3' primer: 5'-caggaaacagctatgac-3' (SEQ ID NO 628)

PCR products were cloned into the TOPO vector and sequenced. The polynucleotide sequence is provided in SEQ ID NO: 267, and the encoded polypeptide sequence is provided in SEQ ID NO: 268.

EXAMPLE 58

2H7 scFv VH L11S(SSS-S)H P331S CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated IgG1 CH2 region and a wild type IgG1 CH3 region. The mutation P331S mutation in the CH2 region, where the proline at residue 331 has been changed to a serine, was incorporated using a single PCR reaction, using a (SSS-S) H WCH2 WCH3 pD18

template and a cycling profile of 94 C, 30 sec; 55 C, 30 sec; 72 C, 30 sec, for 37 cycles. The specific primers for reaction are listed below.

5' primer:

5'ccgtctctgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtccccagc-3' (SEQ ID NO
5 631)

3' Primers: 5'-

gcagggtgtacacctgtggttctcggggctgccctttggctttggagatggtttctcgatggaggctgggagg-3' (SEQ
ID NO 632)

The polynucleotide sequence is provided in SEQ ID NO: 273, and the
10 encoded polypeptide sequence is provided in SEQ ID NO: 274.

EXAMPLE 59

2H7 scFv VH L11S(CSS-S)H P331S CH2 WCH3

This binding region is attached to a mutant IgG connecting region,
where the second and third cysteines have been changed to serines and the proline has
15 been changed to serine (CSS-S), according to methods described in Example 13. The
connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3
region. The mutation P331S mutation in the CH2 region, where the proline at residue
331 has been changed to a serine, was incorporated using a single PCR reaction, using
a (CSS-S)H WCH2 WCH3 pD18 template and a cycling profile of 94 C, 30 sec; 55 C,
20 30 sec; 72 C, 30 sec, for 37 cycles. The specific primers for reaction are listed below.

5' primer: 5'-

ccgtctctgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtccccagc-3' (SEQ ID NO 633)

3' Primer: 5'-

gcagggtgtacacctgtggttctcggggctgccctttggctttggagatggtttctcgatggaggctgggagg-3' (SEQ
25 ID NO 634)

The polynucleotide sequence is provided in SEQ ID NO: 275, and the
encoded polypeptide sequence is provided in SEQ ID NO: 276:

EXAMPLE 60

2H7 scFv VH L11S (SSS-S)H T256N CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is connected to a mutated human IgG connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The T256N mutation in the CH2 region, the threonine at residue 256 has been changed to an asparagine, using the overlapping PCR methods described in Example 56. The specific primers are listed below.

Primers for the first PCR reaction:

5' primer:

5'ttctcttccccccaaaaccaaggacaccctcatgatctcccgaaccctgaggtcac-3' (SEQ ID NO 635)

3' primer: 5'-ggacagtgggagtggcacc-3' (SEQ ID NO 636)

PCR product cloned into TOPO vector and sequenced. This product was used as the template in the second PCR reaction. Primers for the second PCR reaction:

5' primer:

5'ccgtctctgatcaggagcccaaattctctgacaaaactcacacatccccaccgtccccagc-3' (SEQ ID NO 638)

5' overlapping primer: 5'-

tccccaccgtccccagcacctgaactctggggg- gatcgtcagtcttctcttccccccaaaacc-3' (SEQ ID NO) 3' primer: 5'-caggaaacagctatgac-3' (SEQ ID NO 628)

The polynucleotide sequence is provided in SEQ ID NO: 281, and the encoded polypeptide sequence is provided in SEQ ID NO: 282.

EXAMPLE 61

2H7 scFv VH L11S(SSS-S)H RTPE/QNAK (255-258) CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region,

where the leucine has been changed to a serine, as described in Example 33. This binding region is connected to a mutated human IgG connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The RTPE/QNAK mutation in the CH2 region, where residues 255-258 have been mutated from arginine, threonine, proline, glutamic acid to glutamine, asparagines, alanine and lysine, respectively, using the overlapping PCR reactions described in Example 56. The specific primers are listed below.

10 PCR primers for the first PCR reaction:

5' primer: 5'-

ttcctcttccccccaaaaccaaggacaccctcatgatctcccagaacgctaaggtcacatgc-3' (SEQ ID NO: 641)

3' primer: 5'-ggacagtgggagtggcacc-3' (SEQ ID NO 636)

15 The PCR product was cloned into TOPO vector, sequenced and used as a template for the second PCR reaction. The primers for the second PCR reaction:

5 primer:

5'ccgtctctgatcaggagcccaaattctctgacaaaactcacatccccaccgtccccagc-3' (SEQ ID NO 637)

20 5' overlapping primer: 5'-

tccccaccgtccccagcacctgaactcctggggggatcgtcagtct- tctcttccccccaaaacc-3' (SEQ ID NO 638)

3' primer: 5'-caggaaacagctatgac-3' (SEQ ID NO 628)

25 The polynucleotide sequence is provided in SEQ ID NO: 289, and the encoded polypeptide sequence is provided in SEQ ID NO: 290.

EXAMPLE 62

2H7 scFv VH L11S (SSS-S)H K2900 CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This

30

binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The K290Q mutation in the CH2 region, where the Lysine at residue 290 has been changed to a Glutamine, using a single PCR reaction according to the methods described in Example 58. The specific primers used in this reaction are listed below.

5' primer:

5'ccgtctctgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtccccagc-3' (SEQ ID NO 642)

3' primer: 5'-gctccccgggctgtgtcttggc-3' (SEQ ID NO 645)

PCR products were cloned into TOPO and sequenced. The polynucleotide sequence is provided in SEQ ID NO: 297, and the encoded polypeptide sequence is provided in SEQ ID NO: 298.

EXAMPLE 63

2H7 scFv VH L11S (SSS-S)H A339P CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region; where the leucine has been changed to a serine, as described in Example 33. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The A339P mutation in the CH2 region, where the alanine at residue 339 has been changed to a proline, using a single PCR reaction according to the methods described in Example 58. The specific primers used in this reaction are listed below.

5' primer:

5'ccgtctctgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtccccagc-3' (SEQ ID NO 644)

3' Primer: 5'-

ggaggtgggcagggtgtacacctgtggtctcggggctgcccttgggttggagatgg-3' (SEQ ID NO 647)

PCR products were cloned into TOPO and sequenced. The polynucleotide sequence is provided in SEQ ID NO: 305, and the encoded polypeptide sequence is provided in SEQ ID NO: 306.

EXAMPLE 64

5 **G28-1 scFv (SSS-S)H WCH2 WCH3**

This construct has a G28-1 (anti-CD37) single chain Fv binding region made according to methods described in Example 35. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in
10 Example 5. This connecting region is connected to a wild type human IgG1 CH2 and CH3 region as described in Example 1. This construct has previously been referred to as G28-1-MHWTG1C and G28-1 scFv Ig, both have the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 319, and the encoded polypeptide sequence is provided in SEQ ID NO: 320.

15 EXAMPLE 65

G28-1 scFv IgAH WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region made according to methods described in Example 35. This binding region is connected to a human IgA connecting region and wild type human IgG CH2 and CH3
20 constant regions as described in Example 5. This construct has previously been referred to as: G28-1-IgAHWTG1C. The polynucleotide sequence is provided in SEQ ID NO: 321, and the encoded polypeptide sequence is provided in SEQ ID NO: 322.

EXAMPLE 66

G28-1 scFv VH L11S(SSS-S)H WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to
25 methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide
30

sequence is provided in SEQ ID NO: 323, and the encoded polypeptide sequence is provided in SEQ ID NO: 324.

EXAMPLE 67

G28-1 scFv VH L11S (CSS-S)H WCH2 WCH3

5 This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a mutant human IgG1 connecting region, where the second and third cysteines have been changed to serines and the proline has been
10 changed to serine (CSS-S), according to methods described in Example 13. This connecting region is attached to wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 325, and the encoded polypeptide sequence is provided in SEQ ID NO: 326.

EXAMPLE 68

G28-1 scFv VH L11S(CSC-S)H WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a mutant human IgG1 connecting region, where the
20 second cysteine and the proline has been changed to serine (CSC-S), according to methods described in Example 32. This connecting region is attached to wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 327, and the encoded polypeptide sequence is provided in SEQ ID NO: 328.

25 EXAMPLE 69

G28-1 scFv VH L11S(SSC-P)H WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This
30 binding region is attached to a mutant human IgG1 connecting region, where the first and second cysteines have been changed to serines (SSC-P), according to methods

described in Example 13. This connecting region is connected to a wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 329, and the encoded polypeptide sequence is provided in SEQ ID NO: 330.

5

EXAMPLE 70

CTLA4 (SSS-S)H P238SCH2 WCH3

This construct has the extra cellular CTLA-4 binding region as described in Example 14. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced using a PCR assay. PCR reactions were performed using random primed cDNA prepared from human tonsil B cell RNA. PCR amplifications used an amplification profile of 94 C 4 min; [94 C 1 min; 55 C 1 min; 72 C min; for 30 cycles followed by a final extension step for 6 minutes at 72 C. PCR fragments were TOPO cloned and clones with EcoRI inserts of approximately 800 bp were sequenced as described in Example 1. The primers used for the PCR are listed below:

5' primer: 5'-
20 gttgttgatcaggagcccaaatcttctgacaaaactcacacatctccaccgtccccagcacctgaactcctgggtgg
accgtcagttctcc-3' (SEQ ID NO 648)

3' primer: 5'gttgttcttagattatcattaccg- gagacag-3' (SEQ ID NO 649)

This construct has previously been referred to as CTLA-4 IgG MTH (SSS) MTCH2WTCH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 331, and the encoded polypeptide sequence is provided in SEQ ID NO: 332.

25

EXAMPLE 71

CTLA4 (CCC-P) WH WCH2 WCH3

This construct has a CTLA-4 binding region as described in Example 14. This binding region is attached to a wild type human IgG1 connecting region (CCC-P) as described in Example 1. This connecting region is connected to a wild

30

type human IgG1 CH2 and CH3 region as described in Example 1. This construct has previously been referred to as CTLA-4 IgG WTH (CCC) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 47, and the encoded polypeptide sequence is provided in SEQ ID NO:

5 78.

EXAMPLE 72

FC2-2 scFv (SSS-S)H WCH2 WCH3

This construct has a FC₂-2 (anti-CD16) single chain Fv made according to methods described in Example 46. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is connected to a wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 343, and the encoded polypeptide sequence is provided in SEQ ID NO: 344.

15

EXAMPLE 73

FC2-2 scFv VHL11S(SSS-S)H WCH2 WCH3

This construct has a FC₂-2 (anti-CD 16) single chain Fv with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 46. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is connected to a wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 345, and the encoded polypeptide sequence is provided in SEQ ID NO: 346.

25

EXAMPLE 74

UCHL-1 scFv (SSS-S)H WCH2 WCH3

This construct has a UCHL-1 (anti-CD45RO) single chain Fv made according to methods described in Example 48. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5.

30

This connecting region is connected to a wild type human IgG1 CH2 and CH3 region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 357, and the encoded polypeptide sequence is provided in SEQ ID NO: 358.

EXAMPLE 75

5

UCHL-1 scFv VHL11S(SSS-S)H WCH2 WCH3

This construct has a UCHL-1 (anti-CD45RO) single chain Fv with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 48. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is connected to a wild type human IgG1 CH2 and CH3 constant region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 359, and the encoded polypeptide sequence is provided in SEQ ID NO: 360.

15

EXAMPLE 76

5B9 scFv (SSS-S)H WCH2 WCH3

This construct has a 5B9 (anti-CD137) single chain Fv made according to methods described in Example 47. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is connected to a wild type human IgG1 CH2 and CH3 constant region as described in Example 1. This construct has previously been referred to as 5B9 scFv IgG MTH (SSS) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 132, and the encoded polypeptide sequence is provided in SEQ ID NO: 133.

25

EXAMPLE 77

5B9 scFv VHL11S(SSS-S)H WCH2 WCH3

This construct has a 5B9 (anti-CD137) single chain Fv with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 47. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines

30

and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is connected to a wild type human IgG1 CH2 and CH3 constant region as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 365, and the encoded polypeptide sequence is provided in
5 SEQ ID NO: 366.

EXAMPLE 78

2H7 scFv (SSS-S)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1
10 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is connected to a wild type IgG1 CH2 and CH3 constant region as described in Example 1. This construct has previously been referred to as 2H7-MHWTG1C, CytosB-(MHWTG1C)--Ig, anti-CD20 scFv IgG MTH (SSS) WTCH2CH3, CytosB-
15 MHWTG1C, 2H7 scFv-human IgG1 wild type hinge-CH2--CH3, and 2H7 scFv IgG MTH (SSS) WTCH2CH3, which all have the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 57, and the encoded polypeptide sequence is provided in SEQ ID NO: 58.

EXAMPLE 79

2H7 scFv (SSS-S)H P238SCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This binding region is
25 connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in
30 Example 70. This construct has previously been referred to as 2H7 scFv IgG MTH (SSS) MTCH2WTCH3, anti-CD20 scFv IgG MTH (SSS) MTCH2CH3, and CytosB-MHMG1C which all have the same sequence as the above construct. The

polynucleotide sequence is provided in SEQ ID NO: 419, and the encoded polypeptide sequence is provided in SEQ ID NO: 420.

EXAMPLE 80

2H7 scFv IgAH WCH2 WCH3

5 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a human IgA connecting region and wild type human IgG1 CH2 and CH3 constant regions as described in Example 5. This construct has previously been referred to as 2H7 scFv IgAH IgG WTCH2CH3, 2H7 scFv IgA hinge-IgG1 CH2--CH3, and Cytob-
10 IgAHWTHG1C, which all have the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 59, and the encoded polypeptide sequence is provided in SEQ ID NO: 60.

EXAMPLE 81

2H7 scFv IgAH WIGACH2 T4CH3

15 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a human IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon
20 as described in Example 13. This construct has previously been referred to as 2H7 scFv IgAH IgAT4, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 70, and the encoded polypeptide sequence is provided in SEQ ID NO: 71.

EXAMPLE 82

2H7 scFv IgAH WIGACH2 WCH3+JCHAIN

25 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a wild type human IgA connecting region as described in Example 5. This connecting region is attached to a wild type human IgA CH2 and CH3 constant region according to methods described
30 in Example 13. This constant region is attached to a J-chain region as described in

Example 13. The polynucleotide sequence is provided in SEQ ID NO: 61, and the encoded polypeptide sequence is provided in SEQ ID NO: 62.

EXAMPLE 83

2H7 scFv (CCC-P) WH WCH2 WCH3

5 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a wild type human IgG1 connecting region (CCC-P) as described in Example 1. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv Ig WTH
10 (CCC) WTCH2CH3, 2H7 scFv IgG WTH WTCH2CH3, and 2H7 scFv-Ig, which both have the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 27, and the encoded polypeptide sequence is provided in SEQ ID NO: 28.

EXAMPLE 84

2H7 scFv (SSS-S)H WCH2 F405YCH3

15 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region
20 is attached to a wild type human IgG1 CH2 and a mutated human IgG1 CH3 region. The F405Y mutation, where the phenylalanine at residue 405 has been changed to a tyrosine, was introduced according to methods described in Example 21. This construct has previously been referred to as 2H7 scFv MTH WTCH2 MTCH3 Y405, which has the same sequence as the above construct. The polynucleotide sequence is
25 provided in SEQ ID NO: 153, and the encoded polypeptide sequence is provided in SEQ ID NO: 154.

EXAMPLE 85

2H7 scFv (SSS-S)H WCH2 F405ACH3

30 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to

serines (SSS-S) according to methods described in Example 5. This connecting region is attached to a wild type human IgG1 CH2 and a mutated human IgG1 CH3 region. The F405a mutation, where the phenylalanine at residue 405 has been changed to an alanine, was introduced according to methods described in Example 21. This construct has previously been referred to as 2H7 scFv MTH WTCH2 MTCH3 A405, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 155, and the encoded polypeptide sequence is provided in SEQ ID NO: 156.

EXAMPLE 86

2H7 scFv (SSS-S)H WCH2 Y407ACH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to a wild type human IgG1 CH2 and a mutated human IgG1 CH3 region. The Y407A mutation, where the tyrosine at residue 407 has been changed to an alanine, was introduced according to methods described in Example 21. This construct has previously been referred to as scFv MTH WTCH2 MTCH3 A407, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 157, and the encoded polypeptide sequence is provided in SEQ ID NO: 158.

EXAMPLE 87

2H7 scFv (SSS-S) HWCH2 F405A, Y407ACH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S), according to methods described in Example 5. This connecting region is attached to a wild type human IgG1 CH2 and a mutated human IgG1 CH3 region. The F405a and Y407A mutation, where the phenylalanine at residue 405 has been changed to an alanine and the tyrosine at residue 407 has been changed to an alanine, was introduced according to methods described in Example 21. This construct has previously been referred to as scFv MTH WTCH2 MTCH3 A405A407,

which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 161, and the encoded polypeptide sequence is provided in SEQ ID NO: 162.

EXAMPLE 88

5

2H7 scFv (CSS-S)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a mutant human IgG 1 connecting region, where the second and third cysteines have been changed to serines and the proline has been changed to serine (CSS-S), according to methods described in Example 13. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv MTH (CSS) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 134, and the encoded polypeptide sequence is provided in SEQ ID NO: 135.

15

EXAMPLE 89

2H7 scFv (SCS-S)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a mutant human IgG1 connecting region, where the first and third cysteines have been changed to serines and the proline has been changed to serine (SCS-S), according to methods described in Example 13. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv IgG MTH (SCS) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 136, and the encoded polypeptide sequence is provided in SEQ ID NO: 137.

25

EXAMPLE 90

2H7 scFv (SSC-P)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a mutant human IgG1 connecting region, where the first and second cysteines have been changed to serines (SSC-P), according to methods described in Example 13. This connecting region is

30

attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv MTH (SSC) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 138, and the encoded polypeptide sequence is provided in SEQ ID NO: 139.

EXAMPLE 91

2H7 scFv (CSC-S)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a mutant human IgG 1 connecting region, where the second cysteine and the proline has been changed to serine (CSC-S), according to methods described in Example 32. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv MTH (CSC) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 165, and the encoded polypeptide sequence is provided in SEQ ID NO: 166.

EXAMPLE 92

2H7 scFv (CCS-P)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a mutant human IgG1 connecting region, where third cysteine has been changed to a serine (CCS-P), according to methods described in Example 22. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv MTH (CCS) WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 167, and the encoded polypeptide sequence is provided in SEQ ID NO: 168.

EXAMPLE 93

2H7 scFv (SCC-P)H WCH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a mutant human IgG1

connecting region, where first cysteine has been changed to a serine (SCC-P), according to methods described in Example 32. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1.

This construct has previously been referred to as 2H7 scFv MTH (SCC)

- 5 WTCH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 163, and the encoded polypeptide sequence is provided in SEQ ID NO: 164.

EXAMPLE 94

2H7 scFv VH L11S(SSS-S)H WCH2 WCH3

- 10 This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 33. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to
- 15 methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as 2H7 scFv VH11SER IgG MTH (SSS) WTCH2CH3 and 2H7 scFv VH11SER WTH WTCH2CH3, which both have the same sequence as the above construct. The polynucleotide sequence is provided in
- 20 SEQ ID NO: 369, and the encoded polypeptide sequence is provided in SEQ ID NO: 370.

EXAMPLE 95

2H7 scFv VH L11S(CSS-S)H WCH2 WCH3

- 25 This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 33. This binding region is attached to a mutant human IgG1 connecting region, where the second and third cysteines have been changed to serines and the proline has been changed to serine (CSS-S), according to methods described in Example 13. This
- 30 connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 371, and the encoded polypeptide sequence is provided in SEQ ID NO: 372.

EXAMPLE 96

G28-1 scFv VH L11S(SCS-S)H WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a mutant human IgG1 connecting region, where the first and third cysteines have been changed to serines and the proline has been changed to serine (SCS-S), according to methods described in Example 13. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 373, and the encoded polypeptide sequence is provided in SEQ ID NO: 374.

EXAMPLE 97

G28-1 scFv VH L11S(CCS-P)H WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a mutant human IgG1 connecting region, where third cysteine has been changed to a serine (CCS-P), according to methods described in Example 22. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 375, and the encoded polypeptide sequence is provided in SEQ ID NO: 376.

EXAMPLE 98

G28-1 scFv VH L11S(SCC-P)H WCH2 WCH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a mutant human IgG1 connecting region, where first cysteine has been changed to a serine (SCC-P), according to methods described in Example 32. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The polynucleotide sequence is

provided in SEQ ID NO: 377, and the encoded polypeptide sequence is provided in SEQ ID NO: 378.

EXAMPLE 99

G28-1 scFv VH L11S mIgE CH2 CH3 CH4

5 This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a wild type mouse IgE CH2 CH3 and CH4 region using methods described in Example 39. The polynucleotide sequence is provided in SEQ
10 ID NO: 379, and the encoded polypeptide sequence is provided in SEQ ID NO: 380.

EXAMPLE 100

G28-1 scFv VH L11S mIgAH WIGACH2 T4CH3

 This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region,
15 where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a connecting region from mouse IgA as described in Example 39. This connecting region is attached to a mouse IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a 4 amino acid truncation at residues as described in Example 39. The polynucleotide
20 sequence is provided in SEQ ID NO: 381, and the encoded polypeptide sequence is provided in SEQ ID NO: 382.

EXAMPLE 101

G28-1 scFv VH L11S hIgE CH2 CH3 CH4

 This construct has a G28-1 (anti-CD37) single chain Fv binding region
25 with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a wild type human IgE constant region containing CH2, CH3 and CH4 as described in Example 38. The polynucleotide sequence is provided in SEQ ID NO: 383, and the encoded polypeptide sequence is provided in SEQ ID
30 NO: 384.

EXAMPLE 102

G28-1 scFv VH L11S H1GAH W1GACH2 T4CH3

This construct has a G28-1 (anti-CD37) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 35. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a human IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon as described in Example 13. The polynucleotide sequence is provided in SEQ ID NO: 385, and the encoded polypeptide sequence is provided in SEQ ID NO: 386.

EXAMPLE 103

HD37 scFv IgAH WCH2 WCH3

The HD37 scFv was cloned from the HD37 hybridoma using the Novagen-Ig family primer sets, TOPO cloning and sequencing and sewing PCR assay. For the initial PCR reactions prior to sewing, the TOPO clone templates HD37 VH C-1 and HD37 KVL B-9 were used at 1:100 with an amplification profile of 94 C, 30 sec; 55 C, 30 sec; 72 C, 30 seconds for 25 cycles. To provide templates for the secondary SEWING PCR reactions, primary reaction products were gel purified, QIAQUICK purified, and the eluates diluted 50 fold. One microliter each V_L and VH overlapping templates were added to PCR reactions with the following amplification profile: 94 C, 60 sec; 55 C, 60 sec; 72 C, 60 sec; for 30 cycles. After two cycles, the machine was paused, and the flanking 5'V_L and 3'VH primers were added to the reactions at 25 pmol each, and the PCR reactions resumed. PCR products were checked on a gel for the presence of an 750-800 bp fragment, and the reactions products QIAQUICK purified and digested with the appropriate restriction enzymes for insertion into pD 18 Ig expression vectors.

PCR of V_L domain with native leader peptide and part of glyser linker:

5' primer: 5'-gttgtaagcttgccgccatggagacagacacactctgctatgg-3' (SEQ ID NO 652)

3' primer: 5'

gccacccgacccaccaccgcccagccaccgccaccttgattccagcttggtgcctcc-3' (SEQ ID NO 651)

PCR of VL domain without leader peptide (SalI site) and part of glyser linker:

5 5' primer: 5'-gtgtgtgtcgacattgtgtgacccaatctcca-3' (SEQ ID NO 652)

3' priemr: 5'-

gccacccgacccaccaccgcccagccaccgccaccttgattccagcttggtgcctcc-3' (SEQ ID NO 651)

PCR of VH domain with part of glyser linker and BclI site for fusion to -Ig tails.

10 5' primer: 5'-

tcgggcgggtggtgggtcgggtggcggcgatcgacaggttcagctgcagcagctctgg-3' (SEQ ID NO 653)

3' primer: 5'-tcagtgtgatcagaggagacgggtgactgaggttccttg-3' (SEQ ID NO 654)

15 This binding region is connected to a wild type human IgA connecting region and wild type human IgG CH2 and CH3 constant regions as described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as HD37 scFv-IgAHWGTG1C and HD37-IgAHWGTG1C, which both have the same sequence as the above construct. The polynucleotide sequence is provided in
20 SEQ ID NO: 397, and the encoded polypeptide sequence is provided in SEQ ID NO: 398.

EXAMPLE 104

HD37 scFv (SSS-S)H WCH2 WCH3

25 This construct has a HD 37 single chain Fv as described in Example 103. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as HD37-MHWGTG1C and HD37 scFv-
30 IgMHWGTG1C, which both have the same sequence as the above construct. The

polynucleotide sequence is provided in SEQ ID NO: 399, and the encoded polypeptide sequence is provided in SEQ ID NO: 400.

EXAMPLE 105

HD37 scFv VH L11S(SSS-S)H WCH2 WCH3

5 This construct has a HD 37 single chain Fv with a mutation in the heavy chain variable region at amino acid residue 11, where leucine has been changed to serine according to the methods described in Example 50. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in
10 Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 401, and the encoded polypeptide sequence is provided in SEQ ID NO: 402.

EXAMPLE 106

L6 scFv IgAH WCH2 WCH3

15 The L6 scFv was cloned from the L6 hybridoma (1Hellstrom) using the anchor-tailing method described in the Tissue Antigens Paper from 1996. The PCR profile was 94 C, 1 min; 50 C, 2 min; 72 C, 2 min; for 35 cycles. Once consensus sequence was obtained for VL and VH regions from at least 4 TOPO
20 clones, primers were ordered for PCR reactions prior to SEWING PCR reactions as follows: For the initial PCR reactions prior to sewing, the TOPO cloned templates L6VK and L6V_H were used at 1:100 with an amplification profile of 94 C 30 sec; 55 C, 30 sec; 72 C, 30 seconds for 25 cycles. To provide templates for the secondary SEWING PCR reactions, primary reaction products were gel purified, QIAQUICK
25 purified, and the eluates diluted 50 fold. One microliter each V_L and V_H overlapping templates were added to PCR reactions with the following amplification profile: 94 C, 60 sec; 55 C, 60 sec; 72 C, 60 sec; for 30 cycles. After two cycles, the machine was paused, and the flanking 5'V_L and 3'V_H primers were added to the reactions at 25 pmol each, and the PCR reactions resumed. PCR products were checked on a gel for
30 the presence of an 750-800 bp fragment, and the reactions products QIAQUICK purified and digested with the appropriate restriction enzymes for insertion into pD18 Ig expression vectors.

PCR of V_L domain with native leader peptide and part of glyser linker:

L6VLHindIII: 5'-gttgtaagcttgccgccatggatttcaagtgcagatttcagcttc-3'
(SEQ ID NO 655)

L6VLLK3: 5'-

5 gccacccgacccaccaccgcccagccaccgccaccagagagctctttcagctccagcttggt-3' (SEQ ID NO 656)

PCR of VL domain without leader peptide (Sall site) and part of glyser linker:

5' primer: 5'-gttgtgtcgcacattgttctctcccagctctccagcaatcctgtctg-3' (SEQ ID NO 657)

3' primer: 5'-

gccacccgacccaccaccgcccagccaccgccaccagagagctctttcagctccagcttggt-3' (SEQ ID NO 656)

15 PCR of V_H domain with part of glyser linker and BclI site for fusion to -Ig tails.

5': 5'-tcgggcgggtggtgggtcgggtggcggcgatctctgcagatccagttggtgcagtct-3'
(SEQ ID NO 658)

3'Bcl: 5'-tcagtgctgatcagaggagactgtgagagtgggtgccttg-3' (SEQ ID NO 659)

20 This binding region is connected to a human IgA connecting region and wild type human IgG1 CH2 and CH3 constant regions as described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as L6 scFv-IgAHWTG1C, which HAS the same sequence as the above construct. The
25 polynucleotide sequence is provided in SEQ ID NO: 413, and the encoded polypeptide sequence is provided in SEQ ID NO: 414.

EXAMPLE 107

L6 scFv VHL11S(SSS-S)H WCH2 WCH3

30 This construct has a L6 single chain Fv with a mutation in the heavy chain variable region at amino acid residue 11, where leucine has been changed to

serine according to the methods described in Example 49. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The polynucleotide sequence is provided in SEQ ID NO: 415, and the encoded polypeptide sequence is provided in SEQ ID NO: 416:

EXAMPLE 108

2H7 scFv-LLAMA IgG1

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a llama IgG1 hinge, CH2 and CH3 regions according to the methods described in Example 10. The polynucleotide sequence is provided in SEQ ID NO: 21, and the encoded polypeptide sequence is provided in SEQ ID NO: 22.

EXAMPLE 109

2H7 scFv-LLAMA IgG2

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a llama IgG2 hinge, CH2 and CH3 regions according to the methods described in Example 10. The polynucleotide sequence is provided in SEQ ID NO: 23, and the encoded polypeptide sequence is provided in SEQ ID NO: 24.

EXAMPLE 110

2H7 scFv-LLAMA IgG3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a llama IgG3 hinge, CH2 and CH3 regions according to the methods described in Example 10. The polynucleotide sequence is provided in SEQ ID NO: 25, and the encoded polypeptide sequence is provided in SEQ ID NO: 26.

EXAMPLE 111

CD16 Low (ED)(SSS-S)H P238SCH2 WCH3

This construct has the extra cellular, CD16 low affinity allele binding domain as described in Example 43. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The polynucleotide sequence is provided in SEQ ID NO: 421, and the encoded polypeptide sequence is provided in SEQ ID NO: 422.

EXAMPLE 112

CD16-9 HIGH (ED)(SSS-S)H P238SCH2 WCH3

This construct has the extra cellular, CD16 high affinity allele binding domain as described in Example 43. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The polynucleotide sequence is provided in SEQ ID NO: 425, and the encoded polypeptide sequence is provided in SEQ ID NO: 426.

EXAMPLE 113

2E12 scFv (SSS-S)H P238SCH2 WCH3-HCD80TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is

attached to a human CD80 transmembrane and cytoplasmic tail region (hCD80 TM/CT). The hCD80 TM/CT was cloned using random primed cDNA derived from the BJAB cell line according to methods described in Example 12. This TM/CT region was attached to a Ig CH3 region with an open reading frame (ORF). Open reading frame versions of scFvIg constructs of interest were created by replacement of the soluble versions of each -Ig tail with ORF (open reading frame versions) of these tails. PCR primers were designed for the existing clones of soluble -Ig tails which delete the stop codon and add one or more restriction sites to the 3' end of the new -Ig cassettes. The desired transmembrane and cytoplasmic tail sequences can then be subcloned downstream of these new -Ig cassettes. Each construct utilized the existing available 5' BCLI oligonucleotide used in amplifying the soluble version of the tails for the PCR reactions. The 3' oligonucleotides replace the stop codon with out of frame restriction sites fused to the coding region for protein domains involved in regulation of apoptosis.

The PCR amplifications were carried out with 25 pmol of each primer, standard PCR reagents, and varying volumes of either cloned domains or cDNA obtained from PBMC, spleen, or thymus RNA. The reactions used a cycling profile of 94 C, 60sec; 55 C, 60 sec; 72 C, 2 min, for 35 cycles. The primers for the IgG ORF are listed below.

5' primer: 5'-
gtttagatcaggagcccaaatctctgacaaaactcacacatctccaccgtcccagcacctgaactcctgggggaccgt
cagtctcc-3' (SEQ ID NO 660)

3' primer: 5'-
gttgtttcgaaggatccgctttacccgggagcagggagaggctcttctgcgtgtagtg-3' (SEQ ID NO 661)

The polynucleotide sequence is provided in SEQ ID NO: 437, and the encoded polypeptide sequence is provided in SEQ ID NO: 438.

EXAMPLE 114

10A8 scFv (SSS-S)H P238SCH2 WCH3--hCD80TM/CT

This construct has a 10A8 (anti-CD2152) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is

attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. This CH3 region is attached to a hCD80 TM/CT according to methods described in Example 113. The polynucleotide sequence is provided in SEQ ID NO: 439, and the encoded polypeptide sequence is provided in SEQ ID NO: 440.

EXAMPLE 115

40.2.220 scFv (SSS-S)H P238SCH2 WCH3--HCD80TM/CT

This construct has a 40.2.220 (anti-CD40) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. This CH3 region is attached to a hCD80 TM/CT according to methods described in Example 113.

EXAMPLE 116

2H7 scFv (SSS-S)H P238SCH2 WCH3--HCD80TM/CT

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S), according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. This CH3 region is attached to a hCD80 TM/CT according to methods described in Example 113. This construct has previously been referred to as: The polynucleotide sequence is provided in SEQ ID NO: 441, and the encoded polypeptide sequence is provided in SEQ ID NO: 442.

EXAMPLE 117

G19-4 scFv (SSS-S)H P238SCH2 WCH3--hCD80TM/CT

This construct has a GI 9 (anti-CD3) single chain Fv binding region described in Example 29. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. This CH3 region is attached to a hCD80 TM/CT according to methods described in Example 113. The polynucleotide sequence is provided in SEQ ID NO: 443, and the encoded polypeptide sequence is provided in SEQ ID NO: 444.

EXAMPLE 118

2E12 scFv (SSS-S)H WCH2 WCH3--hCD80TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This CH3 region is attached to a hCD80 TM/CT according to methods described in Example 113. This construct has previously been referred to as 2e12 scFv IgG WTH WHTCH3CH2-CD80, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 126, and the encoded polypeptide sequence is provided in SEQ ID NO: 127.

EXAMPLE 119

2E12 scFv IGAH IGACH2 T4CH3--hCD80TM/CT

This construct has a 2I12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a human IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon as

described in example 13. This CH3 region is attached to a hCD80 TM/CT according to methods described in Example 113. The specific primers used to create an IgA ORF are listed below.

5' primer: 5'-gttggtgatcagccagttccctcaactccacctacc-3' (SEQ ID NO: 662)

5 3' primer: 5'-gttgttttcgaaggatccgcgtccacctccgcatgacaacaga (SEQ ID NO: 663)

The polynucleotide sequence is provided in SEQ ID NO: 445, and the encoded polypeptide sequence is provided in SEQ ID NO: 446.

EXAMPLE 120

10 2E12 scFv IgE CH2CH3CH4--hCD80TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is attached to a human IgE constant region containing CH2, CH3 and CH4 as described in Example 38. This CH4 region is attached to a hCD80 TM/CT essentially according to methods described in
15 Example 113. The specific primers used to create an IgE ORF are listed below.

5' primer: 5'-gttggtgatcacgtctgtccagggacttcacc-3' (SEQ ID NO: 664)

3' primer: 5'-gttgttttcgaaggatccgctttaccagatttacagaccgcgcgtg-3'
(SEQ ID NO: 665)

The polynucleotide sequence is provided in SEQ ID NO: 183, and the
20 encoded polypeptide sequence is provided in SEQ ID NO: 184.

EXAMPLE 121

2E12 scFv (SSS-S)H P238SCH2 WCH3--mFADD-TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1
25 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is
30 attached to a mouse FADD transmembrane and cytoplasmic tail region (mFADD

TM/CT). This region is cloned using essentially the same methods described in Example 113. The domain was PCR amplified from randomly primed cDNA from mouse spleen RNA. The specific primers are listed below.

5' primer: 5'- gttgtggatccttcgaacccattcctggtgctgctgcactcgctg-3' (SEQ ID NO: 666)

3' primer: 5'- gttgttatcgatctcgagtcagggtgtttctgaggaagacacagt-3' (SEQ ID NO: 667)

The specific primers used to create a mouse IgG ORF are listed below.

5'primer: 5'-
10 gttgtagatctggagcccagagggcccacaatcaagccctctcctccaagcaaaagccca-3' (SEQ ID NO: 668)

3'primer: 5'-gttgttttcgaaggatccgctttacccggagtcggggagaag-3' (SEQ ID NO: 669)

The polynucleotide sequence is provided in SEQ ID NO: 447, and the
15 encoded polypeptide sequence is provided in SEQ ID NO: 448.

EXAMPLE 122

2E12 scFv (SSS-S)H WCH2 WCH3--mFADD-TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1
20 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region was attached to a mFADD TM/TM region according to methods described in Example 113 and 121. The polynucleotide sequence is provided
25 in SEQ ID NO: 449, and the encoded polypeptide sequence is provided in SEQ ID NO: 450.

EXAMPLE 123

2E12 scFv (SSS-S)H WCH2 WCH3--mCASP3-TM/CT

This construct has a 2e 12 (anti-CD28) single chain Fv binding region
30 described in Example 12. This binding region is connected to a mutated human IgG1

connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region is attached to a mouse casp3 transmembrane and
5 cytoplasmic tail region according to methods described in Examples 113 and 121. The specific primers used to isolate the mcasp3 TM/CT region are listed below:

5'primer: 5'- gttgttgatccttcgaacatggagaacaacaaaacctcagtggattca- 3'
(SEQ ID NO: 670)

3'primer: 5'-gttgttatcgatctcgagctagtgataaaagtacagttcttcgt- 3' (SEQ ID
10 NO: 671)

The polynucleotide sequence is provided in SEQ ID NO: 453, and the encoded polypeptide sequence is provided in SEQ ID NO: 454.

EXAMPLE 124

2E12 scFv (SSS-S)H P238SCH2 WCH3--MCASP3-TM/CT

15 This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3
20 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is connected to a mcasp3 TM/CT region according to methods described in Examples 113, 121 and 123. The polynucleotide sequence is provided in SEQ ID NO: 455, and the encoded polypeptide sequence is provided in SEQ ID NO: 456.

25 EXAMPLE 125

2E12 scFv (SSS-S)H WCH2 WCH3--MCASP8-TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to
30 serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in

Example 1. The CH3 region is attached to a mouse casp8 transmembrane and cytoplasmic tail region (mcase8 TM/CT) essentially according to methods described in Example 113 and 121. The specific primers used to clone the mcase8 TM/CT region are listed below.

5 5' primer: 5'-gtgtttcgaacatggattccagagttgtctttatgctattgctg- 3' (SEQ ID NO: 672)

3' primer: 5'- gttgttatcgatctcgagtcattagggagggaagaagagcttcttccg- 3' (SEQ ID NO: 673)

10 The polynucleotide sequence is provided in SEQ ID NO: 459, and the encoded polypeptide sequence is provided in SEQ ID NO: 460.

EXAMPLE 126

2E12 scFv (SSS-S)H P238SCH2 WCH3--MCASP8-TM/CT

15 This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region was
20 attached to a mcase8 TM/CT region according to methods described in Examples 113, 121 and 125. The polynucleotide sequence is provided in SEQ ID NO: 461, and the encoded polypeptide sequence is provided in SEQ ID NO: 462.

EXAMPLE 127

2E12 scFv (SSS-S)H WCH2 WCH3--HCASP3-TM/CT

25 This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in
30 Example 1. The CH3 region was attached to a human casp3 transmembrane and cytoplasmic tail region (hcase3 TM/CT) essentially according to methods describe in

Examples 113. The specific primers used to clone the hcas3 TM/CT region are listed below.

5' Primer: 5'-gttggtggtccttcgaacatggagaacactgaaaactcagtggat- 3' (SEQ ID NO: 674)

5 3' Primer: 5'- gttgttatcgatctcgagttagtgataaaaatagagttctttt- gtgag- 3' (SEQ ID NO: 675)

The polynucleotide sequence is provided in SEQ ID NO: 465, and the encoded polypeptide sequence is provided in SEQ ID NO: 466.

EXAMPLE 128

10 2E12 scFv (SSS-S)H P238SCH2 WCH3--HCASP3-TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is attached to a hcas3 TM/CT region according to methods described in Examples 113 and 127. The polynucleotide sequence is provided in SEQ ID NO: 467, and the encoded polypeptide sequence is provided in SEQ ID NO: 468.

EXAMPLE 129

2E12 scFv (SSS-S)H WCH2 WCH3--HCASP8-TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region is attached to a human casp8 transmembrane and cytoplasmic tail region (hcas8 TM/CT) according to methods described in Example 133. The specific primers used to clone the hcas8 TM/CT region are listed below.

5' Primer: 5'-gttggtggatccttcgaacatg- gacttcagcagaaatctttatgat- 3' (SEQ ID NO: 676)

3' Primer: 5'- gttgttatcgatgcatgctcaatcagaagggaagacaagttttttct- 3' (SEQ ID NO: 677)

5 The polynucleotide sequence is provided in SEQ ID NO: 473, and the encoded polypeptide sequence is provided in SEQ ID NO: 474.

EXAMPLE 130

2E12 scFv (SSS-S)H P238SCH2 WCH3--hCASP8-TM/CT

This construct has a 2e12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is attached to a hcasp8 TM/CT according to methods described in Examples 113 and 129. The polynucleotide sequence is provided in SEQ ID NO: 475, and the encoded polypeptide sequence is provided in SEQ ID NO: 476.

EXAMPLE 131

1D8 scFv (SSS-S)H P238SCH2 WCH3--hCD80TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is attached to a hCD80 TM/CT region according to methods described in Example 113. The polynucleotide sequence is provided in SEQ ID NO: 108, and the encoded polypeptide sequence is provided in SEQ ID NO: 109.

EXAMPLE 132

1D8 scFv (SSS-S)H WCH2 WCH3--hCD80TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region is attached to a hCD80 TM/CT region according to methods described in Example 113. This construct has previously been referred to as 1D8 scFv IgG WTH WTCH2CH3-CD80, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 481, and the encoded polypeptide sequence is provided in SEQ ID NO: 482.

EXAMPLE 133

1D8 scFv-mIgAH WIGA CH2 T4CH3--hCD80TM/CT

This construct has a 1 D8 (anti-4-1 BB) single chain Fv binding region described in Example 25. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a mouse IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon as described in Example 39. The CH3 region can be attached to a hCD80 TM/CT region according to methods described in Examples 113 using primers that create an IgA ORF (SEQ ID NOs: 483 and 484).

EXAMPLE 134

1D8 scFv IgE CH2CH3CH4--hCD80TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is attached to a human IgE constant region containing CH2, CH3 and CH4 as described in Example 38. The CH4 region is attached to a hCD80 TM/CT according to methods described in Examples 113 and 120. The polynucleotide sequence is provided in SEQ ID NO: 175, and the encoded polypeptide sequence is provided in SEQ ID NO: 176.

EXAMPLE 135

1D8 scFv (SSS-S)H P238SCH2 WCH3--mFADD-TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region was attached to a mFADD TM/TM region according to methods described in Example 113 and 121. The polynucleotide sequence is provided in SEQ ID NO: 487, and the encoded polypeptide sequence is provided in SEQ ID NO: 488.

EXAMPLE 136

1DS scFv (SSS-S)H WCH2 WCH3--mFADD-TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region was attached to a mFADD TM/TM region according to methods described in Example 113 and 121. The polynucleotide sequence is provided in SEQ ID NO: 485, and the encoded polypeptide sequence is provided in SEQ ID NO: 486.

EXAMPLE 137

1D8 scFv (SSS-S)H WCH2 WCH3--mCASP3-TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region was attached to a mcasp3 TM/TM region according to

methods described in Example 113, 121 and 123. The polynucleotide sequence is provided in SEQ ID NO: 489, and the encoded polypeptide sequence is provided in SEQ ID NO: 490.

EXAMPLE 138

5 **1D8 scFv (SSS-S)H P238SCH2 WCH3--MCASP3-TM/CT**

 This construct has a 1 D8 (anti-4-1 BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is
10 attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region was attached to a mcasp3 TM/TM region according to methods described in Example 113, 121 and 123. The polynucleotide sequence is provided in SEQ ID NO: 491, and the
15 encoded polypeptide sequence is provided in SEQ ID NO: 492.

EXAMPLE 139

1D8 scFv (SSS-S)H WCH2 WCH3--MCASP8-TM/CT

 This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1
20 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region was attached to a mcasp8 TM/TM region according to methods described in Example 113, 121 and 125. The polynucleotide sequence is
25 provided in SEQ ID NO: 493, and the encoded polypeptide sequence is provided in SEQ ID NO: 494.

EXAMPLE 140

1D8 scFv (SSS-S)H P238SCH2 WCH3--MCASP8-TM/CT

 This construct has a 1 D8 (anti-4-1 BB) single chain Fv binding region
30 described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to

serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region was attached to a mcasp8 TM/TM region according to methods described in Example 113, 121 and 125. The polynucleotide sequence is provided in SEQ ID NO: 495, and the encoded polypeptide sequence is provided in SEQ ID NO: 496.

EXAMPLE 141

1D8 scFv (SSS-S)H WCH2 WCH3--HCASP3-TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region is attached to a hcasp3 TM/CT according to methods described in Examples 113 and 127. The polynucleotide sequence is provided in SEQ ID NO: 497, and the encoded polypeptide sequence is provided in SEQ ID NO: 498.

EXAMPLE 142

1D8 scFv (SSS-S)H P238SCH2 WCH3--HCASP3-TM/CT

This construct has a 1 D8 (anti-4-1 BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is attached to a hcasp3 TM/CT according to methods described in Examples 113 and 127. The polynucleotide sequence is provided in SEQ ID NO: 499, and the encoded polypeptide sequence is provided in SEQ ID NO: 500.

EXAMPLE 143

1D8 scFv (SSS-S)H WCH2 WCH3--HCASP8-TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. The CH3 region is attached to a hcas8 TM/CT according to methods described in Examples 113 and 129. The polynucleotide sequence is provided in SEQ ID NO: 501, and the encoded polypeptide sequence is provided in SEQ ID NO: 502.

EXAMPLE 144

1D8 scFv (SSS-s)H P238SCH2 WCH3--HCASP8-TM/CT

This construct has a 1D8 (anti-4-1BB) single chain Fv binding region described in Example 25. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This hinge region is attached to a mutated human IgG1 CH2 region and a wild type human IgG1 CH3 region. The P238S mutation, where a proline at residue 238 was changed to a serine, was introduced according to methods described in Example 70. The CH3 region is attached to a hcas8 TM/CT according to methods described in Examples 113 and 129. The polynucleotide sequence is provided in SEQ ID NO: 503, and the encoded polypeptide sequence is provided in SEQ ID NO: 504.

EXAMPLE 145

L6 scFv (SSS-S)H WCH2 WCH3

This construct has a L6 scFv binding domain as described in Example 105. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This construct has previously been referred to as L6 scFv-IgMHWGTG1C, which has the same sequence as the above construct. The polynucleotide sequence is provided in

SEQ ID NO: 415, and the encoded polypeptide sequence is provided in SEQ ID NO: 416.

EXAMPLE 146

2H7 scFv CD154 (L2)

5 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region has been attached to CD154 extracellular domain according to methods described in Example 4. The polynucleotide sequence is provided in SEQ ID NO: 690, and the encoded polypeptide sequence is provided in SEQ ID NO: 691.

EXAMPLE 147

2H7 scFv CD154 (S4)

10 This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region has been attached to CD154 extracellular domain according to methods described in Example 4, such that methods
15 of attachment resulted in a truncated version compared to the construct describe in Example 146. The polynucleotide sequence is provided in SEQ ID NO: 692, and the encoded polypeptide sequence is provided in SEQ ID NO: 693.

EXAMPLE 148

CTLA4 IgAH IgACH2CH3

20 This construct has the extra cellular CTLA-4 binding region as described in Example 14. This binding region is attached to a wild type human IgA connecting region as described in Example 5. This connecting region is attached to a wild type human IgA CH2 and CH3 constant region according to methods described in Example 13. This constant region is attached to a J-chain region as described in
25 Example 13. This construct has previously been refered to as CTLA-4 IgAH IgACH2CH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 505, and the encoded polypeptide sequence is provided in SEQ ID NO: 506.

EXAMPLE 149

CTLA4 IgAH IgACH2 T4CH3

This construct has the extra cellular CTLA-4 binding region as described in Example 14. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a human IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon as described in Example 13. This construct has previously been referred to as CTLA-4 IgAH IgA-T4, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 507, and the encoded polypeptide sequence is provided in SEQ ID NO: 508.

EXAMPLE 150

2H7 scFv IgAH IgACH2CH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a wild type human IgA connecting region as described in Example 5. This connecting region is attached to a wild type human IgA CH2 and CH3 constant region according to methods described in Example 13. The polynucleotide sequence is provided in SEQ ID NO: 61, and the encoded polypeptide sequence is provided in SEQ ID NO: 62.

EXAMPLE 151

2H7 scFv IgAH IgAHCH2 T18CH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a human IgA constant region consisting of a wild type CH2 region and a mutated CH3 region where there is a truncation of 18 amino acid residues prior to the 3' stop codon as described in Example 13. The polynucleotide sequence is provided in SEQ ID NO: 509, and the encoded polypeptide sequence is provided in SEQ ID NO: 510.

EXAMPLE 152

2H7-40.2.220 scFv (SSS-S)H WCH2 WCH3

A bispecific fusion protein was constructed between 2H7 scFv (anti-CD20) and 40.2.220 (anti-CD40), which both target B cell receptors. The 2H7 scFv hIgG1 (SSS-S)H WCH2 WCH3 construct in the expression vector pD18 was passaged through dam⁻ bacteria in order to permit cleavage at the BclI site. Cleaved plasmid was treated with alkaline phosphatase prior to ligation to a BclI cut linker-CD40 scFv fragment. This fragment was synthesized from the existing 40.2.220 scFv by successive PCR reactions with overlapping primers. The linker attached is a previously patented (BMS patent issued) helical type linker with a high number of lysine and glutamic acid residues. The scFv for CD40 was PCR amplified without the leader peptide as a Sall-BclI fragment, but with the hinge type linker substituted at the amino terminus as a BclI-Sall fragment, to mediate insertion of the linker-scFv cassette as a BclI fragment between the scFv and -Ig tail included in an existing scFvIg construct for CD20 (2H7 scFv hIgG1 constructs). The 3' end was similar to the other scFv V_H molecules with an out of frame BclI site fused to the VTVSS type sequence at the end of the V_H domain. PCR oligos:

40.2.220 scFv:

5' primer--40.2.220S5: 5'-gttggtgtcgcacattgttctgactcagtcctccagccaccctgtc-
3' (SEQ ID NO: 678)

3' primer--40.2.220Bcl3: 5'-gttggtgatcagagacagtgaccagtgtcccttgg-3'
(SEQ ID NO: 679)

Linker Primers:

BclI-Sall fragment created by annealing complementary oligonucleotides. This fragment is then ligated into BclI digested vector along with a Sall-BclI scFv to create the (linker-scFv) BclI fragment desired for shuttling.

5' primer: 5'-
gatcaatccaactctgaagaagcaaagaaaggagggccaaaaaggaggaagccaagaaatctaacagcg-3' (SEQ ID NO: 560)

3' primer: 5'-

tcgacgctgtagatttcttggttcctccttttggcctcctcttcttcttgcttcttcagagttggatt-3' (SEQ ID NO: 563)

5 This BclI fragment was then ligated downstream of the 2H7 scFv in pD 18-Ig. Transformants were screened for the presence of a 2.4 kb HindIII-Xba insert and positive clones sequenced prior to further studies. COS cell transient transfections were performed with this construct and culture supernatants screened for the presence of protein of the predicted size and for binding to CD20 and to CD40 transfected CHO cells.

10 New bispecific constructs can be created by designing hinge type linkers which incorporate one or preferably two restriction sites at either end of the linker, facilitating asymmetric digests and transfer of (linker-scFv) or (scFv-linker) cassettes between different constructs. These constructs will also incorporate the V_H L11S and other V region substitutions, which presumably facilitate proper folding and
15 result in increased expression of the molecules in which they are inserted.

This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1.
20 This construct has previously been referred to as anti-CD20-anti-CD40 scFv IgG MTH (SSS) MTCH2WTCH3, which has the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NO: 114, and the encoded polypeptide sequence is provided in SEQ ID NO: 115.

EXAMPLE 153

25 **2H7 scFv IgAH IgACH2 T4CH3--hCD80 TM/CT**

This construct has a 2H7 (anti-CD20) single chain Fv binding region as described in Example 1. This binding region is attached to a connecting region from human IgA as described in Example 5. This connecting region is attached to a human IgA constant region consisting of a wild type CH2 region and a mutated CH3
30 region where there is a truncation of 4 amino acid residues prior to the 3' stop codon as described in Example 13. This CH3 region is attached to a hCD80 TM/CT according to the methods described in Examples 113 and 119. This construct has

previously been referred to as 2H7 scFv IgA hinge IgA-T4-CD80 and 2H7 scFv IgAH IgA-T4-CD80, which both have the same sequence as the above construct. The polynucleotide sequence is provided in SEQ ID NOs: 70 and 29, and the encoded polypeptide sequence is provided in SEQ ID NOs: 71 and 30.

5

EXAMPLE 154

G19-4 scFv (CCC-P) WH WCH2 WCH3--hCD80 TM/CT

This construct has a G19 (anti-CD3) single chain Fv binding region described in Example 29. This binding region is attached to a wild type human IgG1 connecting region (CCC-P) as described in Example 1. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This CH3 region is attached to a hCD80 TM/CT according to the methods described in Examples 113. The polynucleotide sequence is provided in SEQ ID NOs: 112 and 29, and the encoded polypeptide sequence is provided in SEQ ID NOs: 113 and 30.

15

EXAMPLE 155

2E12 scFv (CCC-P) WH WCH2 WCH3--hCD80 TM/CT

This construct has a 2E12 (anti-CD28) single chain Fv binding region described in Example 12. This binding region is attached to a wild type human IgG1 connecting region (CCC-P) as described in Example 1. This connecting region is attached to wild type human IgG1 CH2 and CH3 constant regions as described in Example 1. This CH3 region is attached to a hCD80 TM/CT according to the methods described in Examples 113. The polynucleotide sequence is provided in SEQ ID NO: 126, and the encoded polypeptide sequence is provided in SEQ ID NO: 127.

25

EXAMPLE 156

2H7 VHL11S scFv (SSS-S) IgECH3CH4

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 33. This binding region is connected to a mutated human IgG1 connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. This connecting region is attached to human IgE

30

CH3 and CH4 constant region. This truncated constant region was created according to the methods described in Example 38. The polynucleotide sequence is provided in SEQ ID NO: 227, and the encoded polypeptide sequence is provided in SEQ ID NO: 228

5

EXAMPLE 157

IGD HINGE

An alternative hinge region can be isolated from human IgD immunoglobulin hinge region by using PCR assay to isolate the desired region. The PCR reaction is the same used in Example 1. This hinge was truncated by 6 amino acid residues at the 3' end. The primers used in this PCR reaction are listed below

5" Primer: 5'-
GTGGATCCAGGTTCGAAGTCTCCAAAGGCACAGGCC- 3' (SEQ ID NO: 534)
3' primer: 5'-
GTTGTCGACTGCACCGGTCTTTGTCTCTCTCTCTTC- 3'(SEQ ID NO: 517)

15 The polynucleotide sequence is provided in SEQ ID NO: 237, and the encoded polypeptide sequence is provided in SEQ ID NO: 238.

EXAMPLE 158

hCD28 TM/CT

For some of the cell surface ORF constructs, the transmembrane domain of CD80 was substituted with the transmembrane domain of human CD28 because it forms a dimer on the cell surface rather than a monomer as the CD80 does. Several of the molecules which drive the apoptotic program require oligomerization/trimerization to form a signaling complex; therefore, it is important to be able to control initiation of signaling by controlling the degree of oligomerization of these recombinant receptors on the cell surface. The primers used in PCR amplification of the CD28 tail are given below:

5' Primer: 5'- gttgtggatccttcgaacccttttgggtgctggtggtggtggtgga- 3'
(SEQ ID NO: 518)
3' primer: 5'-gttggtatcgatctcgagtcaggagcgataggctgcgaagtc-3'(SEQ ID
30 NO: 519)

EXAMPLE 159

2H7 scFv VH L11S (SSS-S)H K322L CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine as described in Example 33. This binding region is connected to a mutated human IgG connecting region where all of the cysteines and one proline have been changed to serines (SSS-S) according to methods described in Example 5. The connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The K322L mutation in the CH2 region is at a residue 322, where a Lysine has been changed to a leucine using overlapping PCR described in Example 56, but with different primers for the first PCR reaction, which are listed below.

5' primer:

5'ttctcttcccccaaaaccaaggacaccctcatgatctcccgaaccctgaggtcac-3' (SEQ ID NO: 635)

3' primer: 5'-ggacagtgggagtggcacc-3' (SEQ ID NO: 621)

PCR product was cloned into TOPO vector and sequenced. The polynucleotide sequence is provided in SEQ ID NO: 265, and the encoded polypeptide sequence is provided in SEQ ID NO: 266.

EXAMPLE 160

2H7 scFv VH L11S (CSS-S)H K322L CH2 WCH3

This construct has a 2H7 (anti-CD20) single chain Fv binding region with a point mutation at amino acid residue 11 in the heavy chain variable region, where the leucine has been changed to a serine, as described in Example 33. This binding region is attached to a mutant IgG connecting region, where the second and third cysteines have been changed to serines and the proline has been changed to serine (CSS-S), according to methods described in Example 23. The connecting region is attached to a mutated IgG CH2 region and a wild type IgG CH3 region. The K322L mutation in the CH2 region is at a residue 322, where a Lysine has been changed to a leucine using overlapping PCR described in Example 56, using primers from Example 159 in the first PCR reaction and primers from Example 57 for the second PCR reaction.

PCR products were cloned into the TOPO vector and sequenced. The polynucleotide sequence is provided in SEQ ID NO: 261, and the encoded polypeptide sequence is provided in SEQ ID NO: 262.

Additional representative constructs or sequences within various of
5 examples of the present invention are as follows:

NT

HuIgG1 wild type hinge, CH2, CH3 (nucleotidesequence) (SEQ ID NO: 1)

tctgatcaggagcccaaatcttgacaaaactcacacatgccaccgtgccagcacctgaactcctggggggaccgtc
10 agtcttctcttcccccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgt
gagccacgaagaccctgaggtcaagttcaactgtacgtggacggcgtggaggtgcataatgccaagacaaagccgcg
ggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggag
tacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacaatctcaaagccaaagggcagccccgag
aaccacaggtgtacaccctgcccccatcccggtgagctgaccaagaaccaggtcagcctgacctgcctggtcaaagg
15 cttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccggtg
ctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctca
tgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

20 **HuIgG1 wild type hinge, CH2, CH3** (amino acid sequence) (SEQ ID NO: 2)

SDQEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV
SHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNG
KEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLV
KGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN
25 VFSCSVMHEALHNHYTQKSLSLSPGK

NT

Llama IgG1 hinge, CH2, CH3 (nucleotide sequence) (SEQ ID NO: 3)

tgatcaagaaccacatggaggatgcacgtgccncagtgccncaatgccngcncngaactnccaggaggcccttct
30 gtctttgtcttcccccgaaaccaaggacgtcctctccatttttgagggccgagtcacgtgcgttgtagtggacgtcggaaa

gaaagaccccgaggtcaatttcaactggtatattgatggcgttgaggtgcgaacggccaatacgaagccaaaagaggaac
agttcaacagcacgtaccgcgtgggtcagcgtcctgccccatccagcaccaggactggctgacggggaaggaattcaagt
caagggtcaacaacaaagctctcccgcccccatcgagaggaccatctccaaggccaaagggcagacccgggagccgc
aggtgtacaccctggccccacaccgggaagaactggccaaggacaccgtgagcgtaacatgcctggtcaaaggcttcta
5 cccagctgacatcaacgttgagtggcagaggaacggtcagccggagtcagagggcacctacgccaacacgccgccaca
gctggacaacgacgggacctacttctctacagcaagctctcggtgggaaagaacacgtggcagcggggagaaacctta
acctgtgtggtgatgcatgaggccctgcacaaccactacaccagaaatccatcacccagtcttcgggtaaatagtaatcta
ga

10 AA

Llama IgG1 hinge, CH2, CH3 (in figure 23 as Llama IgG1) (amino acid sequence) (SEQ ID NO: 4)

EPHGGCTCPQCPAPELPGGPSVFVFPKPKDVLSISGRPEVTCVVVDVGKEDP
EVNFNWYIDGVEVRTANTKPKEEQFNSTYRVVSVLPIQHQDWLTGKEFKCK
15 VNNKALPAPIERTISKAKGQTREPQVYTLAPHREELAKDTVSVTCLVKGFYP
DINVEWQRNGQPESEGTYANTPPQLDNDGTYFLYSRLSVGKNTWQRGETLT
GVVMHEALHNHYTQKSITQSSGK

NT

20 **Llama IgG2 (nucleotide sequence) (SEQ ID NO: 5)**

tgatcaagaaccaagacacaaaaccacaaccacaaccacaaccaatcctacaacagaatccaagtgctcc
aatgtccagcccctgagtcctgggagggccctcagttctcatcttcccccgaaaccaaggacgtcctctccattctg
ggaggccccgaggtcacgtgcgttggttagacgtgggccaggaagaccccgaggtcagtttcaactggtacattgatggc
gctgaggtgcgaacggccaacacgaggccaaaagaggaacagttcaacagcacgtaccgcgtggtcagcgtcctgccc
25 atccagcaccaggactggctgacggggaaggaattcaagtgaaggtcaacaacaaagctctcccgcccccatcgaga
agaccatctccaaggccaaagggcagacccgggagccgcaggtgtacaccctggccccacaccgggaagagctggcc
aaggacaccgtgagcgtaacatgcctggtcaaaggcttctaccacctgatataacgttgagtggcagaggaatgggca
gccggagtcagagggcacgtacgccaccacgccacccagctggacaacgacgggacctacttctctacagcaagct
ctcggtgggaaagaacacgtggcagcagggagaaaccttcacctgtgtggtgatgcacgaggccctgcacaaccactac
30 acccagaaatccatcacccagtcttcgggtaaatagtaatctaga

AA

Llama IgG2 (amino acid sequence) (SEQ ID NO: 6)

DQEPKTPKPQPQPQPQNPTTESKCPKCPAPELLGGPSVFIFPPKPKDVLSISGR
PEVTCVVVDVVGQEDPEVSFNWYIDGAEVRTANTRPKEEQFNSTYRVVSVLPI
5 QHQDWLTGKEFKCKVNNKALPAPIEKTISKAKGQTREPQVYTLAPHREELAK
DTVSVTCLVKGFYPPDINVEWQRNGQPESEGTYATTPPQLDNDGTYFLYSKL
SVGKNTWQQGETFTCVVMHEALHNHYTQKSITQSSGK

NT

10 **Llama IgG3 Fc (nucleotide sequence)** (SEQ ID NO: 7)

tgatcaagcgcaccacagcgaagaccccagctccaagtgtcccaaatgccaggccctgaactccttgaggggcccacg
gtcttcatcttcccccgaaagccaaggacgtcctctccatcacccgaaaacctgaggtcacgtgcttggtggacgtggg
taaagaagaccctgagatcgagttcaagctggtccgtggatgacacagaggtacacacggctgagacaaagccaaagga
ggaacagttcaacagcacgtaccgcgtggtcagcgtcctgccatccagcaccaggactggctgacggggaaggaattc
15 aagtgaaggtcaacaacaaagctctcccagccccatcgagaggaccatctccaaggccaaagggcagacccgggag
ccgcaggtgtacacctggccccacaccgggaagagctggccaaggacaccgtgagcgtaacctgcctggtcaaaggc
ttctcccagctgacatcaacgttgagtggcagaggaatgggcagccggagtcagagggcacctacgccaacacgccgc
cacagctggacaacgacgggacctacttctctacagaaaactctccgtgggaaagaacacgtggcagcagggagaagt
cttcacctgtgtggtgatgcacgaggctctacacaatcactccaccagaaatccatcacccagttctcgggtaaataagta
20 ctagagggcc

AA

Llama IgG3 Fc (amino acid sequence) (SEQ ID NO: 8)

DQAHHSEDPSSKCPKCPGPELLGGPTVFIFPPKAKDVLSITRKPEVTCLWWTW
25 VKKTLRSSSSWSVDDTEVHTAETKPKEEQFNSTYRVVSVLPIQHQDWLTGKE
FKCKVNNKALPAPIERTISKAKGQTREPQVYTLAPHREELAKDTVSVTCLVKG
FFPADINVEWQRNGQPESEGTYANTPPQLDNDGTYFLYSKLSVGKNTWQQG
EVFTCVVMHEALHNHSTQKSITQSSGK

30 NT

HuIgG1 wild type hinge (nucleotide sequence) (SEQ ID NO: 9)

gatcaggagcccaaattctgtgacaaaactcacacatgccaccgtgccagca

AA

5 **HuIgG1 wild type hinge (amino acid sequence)** (SEQ ID NO: 10)

DQEPKSCDKTHTCPPCPA

NT

HuIgG1 H2, wild type hinge with leu at second position (nucleotide sequence)

10 gatctggagcccaaattctgtgacaaaactcacacatgccaccgtgccagca (SEQ ID NO: 11)

AA

HuIgG1 H2, wild type hinge with leu at second position (amino acid sequence)

DLEPKSCDKTHTCPPCPA (SEQ ID NO: 12)

15

NTHuIgG1 wild type CH2 (nucleotide sequence) (SEQ ID NO: 13)

cctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacaccctcatgatctccggaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcacca
20 ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaaccatctcc
aaagccaaa

AA

HuIgG1 wild type CH2 (amino acid sequence) (SEQ ID NO: 14)

25 PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE
VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK
TISKAK

NT

HuIgG1 wild type CH3 (nucleotide sequence) (SEQ ID NO: 15)

gggcagccccgagaaccacaggtgtacaccctgcccccatccggaggagatgaccaagaaccaggtcagcctgac
ctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaag
5 accacgcctcccgtgctggactccgacggctccttctctctatagcaagtcaccgtggacaagagcaggtggcagca
ggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccggg
taaata

AA

10 **HuIgG1 wild type CH3 (amino acid sequence)** (SEQ ID NO: 16)

GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSVSMHEALHNHYTQKSLSLS
PGK

15 NT

HuIgG1 mutated hinge (C-C-C→S-S-S) (nucleotide sequence) (SEQ ID NO: 17)

gatcaggagcccaaattcttctgacaaaactcacacatccccaccgtccccagca

AA

20 **HuIgG1 mutated hinge (C-C-C→S-S-S) (amino acid sequence)** (SEQ ID NO: 18)

DQEPKSSDKTHTSPPSPA

NT

25 **HIgG1MTH WTCH2CH3 (mutant hinge with wild type CH2 and CH3 (reads from the hinge+Ig tail) (nucleotide sequence)** (SEQ ID NO: 19):

tgatcaccccaaattcttctgacaaaactcacacatctccaccgtcctcagcacctgaactcctgggtggaccgtcagtcttcc
tcttccccccaaaaccaaggacaccctcatgatctccggaccctgaggtcacatgcgtgggtggacgtgagccac
gaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggag

cagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtg
caaggtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccaca
gggtgtacacctgccccatcccgggatgagctgaccaagaaccaggctcagcctgacctgcctggctcaaaggcttctatcc
cagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactc
5 cgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgt
gatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgataatctaga

AA

Mutant hinge, but wild type CH2 and CH3 (amino acid sequence) (SEQ ID NO:
10 20)

DHPKSSDKTHTSPPSSAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSH
EDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKG
FYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVF
15 SCSVMHEALHNHYTQKSLSLSPGK

NT

LLG1-5'bgl **35 mer** **Llama IgG1 5'** (SEQ ID NO: 542)

5'-gtt gtt gat caa gaa cca cat gga gga tgc acg tg-3'

20

NT

LLG2-5'bgl **32 mer, Llama IgG2-5'** (SEQ ID NO: 543)

5'-gtt gtt gat caa gaa ccc aag aca cca aaa cc-3'

25 NT

LLG3-5'bgl **33 mer, Llama IgG3-5'** (SEQ ID NO: 544)

5'-gtt gtt gat caa gcg cac cac agc gaa gac ccc-3'

NT

LLseqsense 19mer, llama sequencing primer (SEQ ID NO: 450)

5'-ctg aga tcg agt tca gct g-3'

5 **LLseqAS 19 mer (SEQ ID NO: 451)**

5'-cct cct ttg gct ttg tct c-3'

NT

2H7 scFv llama IgG1 (nucleotide sequence) (SEQ ID NO: 21)

10 aagcttcccgcctggaatttcaagtgcagatttcagcttctgtaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctcccccaccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgaggatttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg
15 ggcgggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgtgtactatagtaacttactgtgtacttcgagtctggggcacagggaccacggtcaccgtctctctgtatcaagaa
20 ccacatggaggatgcacgtgccncagtgcccncaatgccngcncngaaactnccaggaggcccttctgtctttgtctt
ccccccgaaaccaaggacgtcctctccattttggaggccgagtcacgtgcgttgtagtggtgacgtcggaaagaaagacc
ccgaggtcaatttcaactggtatattgatggcgttgaggtgcgaacggccaatacgaagccaaaagaggaacagttcaaca
gcacgtaccgcgtggtcagcgtcctgcccacccagcaccaggactggctgacggggaaggaattcaagtgaagggtcaa
caacaaagctctccccggcccccatcgagaggaccatctccaaggccaaagggcagaccgggagccgcaggtgtaca
25 ccctggccccacaccgggaagaactggccaaggacaccgtgagcgtaacatgcctggtcaaaggcttctaccagctga
catcaacgttgagtggcagaggaacggtcagccggagtcagagggcacctacgccaacacgccgccacagctggaca
acgacgggaccttactctctacagcaagctctcggtgggaaagaacacgtggcagcggggagaaaccttaacctgtgtg
gtgatgcatgaggccctgcacaaccactacaccagaaatccatcaccagttctcggttaaatagtaaatctaga

30 AA

2H7 scFv llama IgG1 (amino acid sequence) (SEQ ID NO: 22)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
5 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSSDQEPHGGCTCPQCPAPELPGGPSVFVFPPKPKDVLSIFGGRVTCVVVD
VGKKDPEVNFNWDYIDGVEVRTANTKPKEEQFNSTYRVVSVLPIQHGDWLTG
KEFKCKVNNKALPAPIERTISKAKGQTREPQVYTLAPHREELAKDTVSVTCLV
10 KGFYPADINVEWQRNGQPESEGTYANTPPQLDNDGTYFLYSKLSVGKNTWQ
RGETLTCVVMHEALHNHYTQKSITQSSGK

NT

2H7 scFv llama IgG2 (nucleotide sequence) (SEQ ID NO: 23)

15 aagcttgccgcatggatttcaagtcagattttcagcttctgctaactcagtcgttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtcgagtcgggtctgggaccttactctctcacaatcagcagagtcggaggtgaagatgctgccactt
attactgccagcagtcggagtttaacccaccacgcttcggtgctgggaccaagtcggagctgaaagatggcgggtggctcg
20 ggcgggtggatctggaggaggtgggagctctcaggctatctacagcagtcctggggctgagctggtgagccctgggg
cctcagtcgaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtcggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcaagaa
25 cccaagacacaaaaccacaaccacaaccacaaccacaaccacaaccacaacccaatcctacaacagaatccaagtccccaatgtccag
ccctgagctcctgggagggccctcagtccttctctcccccgaaaccaaggacgtcctctccatttctgggagggcccg
aggtcacgtgcgttggttagacgtgggccaggaagaccccgaggtcagttcaactggtacattgatggcgtgaggtg
cgaacggccaacacgaggccaaaagaggaacgttcaacagcacgtaccgctggtcagcgtcctgccatccagcac
caggactggctgacggggaaggaattcaagtgaaggtcaacaacaagctctccggcccccatcgagaagaccatct
30 ccaaggccaaagggcagaccgggagccgcaggtgtacacctggccccacaccgggaagagctggccaaggacac
cgtgagcgtaacatgcctggtcaaaggcttctacccacctgatataacgttgagtcgagaggaatgggcagccggagt
cagagggcacgtacgccaccacgccacccacagctggacaacgacgggaccttctctacagcaagctctcggtggg

aaagaacacgtggcagcagggagaaaccttcacctgtgtggtgatgcacgaggccctgcacaaccactacaccagaaa
tccatcaccagctcttcgggtaaataagtaatctaga

AA2H7 scFv llama IgG2 (amino acid sequence) (SEQ ID NO: 24)

5 MDFQVQIFSFLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPFTGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
10 VTVSSDQEPKTPKPQPQPQPQNPTTESKCPKCPAPELLGGPSVFIFPPKPKDVL
SISGRPEVTCVVVDVGQEDPEVSFNWYIDGAEVRTANTRPKEEQFNSTYRVV
SVLPIQHQDWLTGKEFKCKVNNKALPAPIEKTISKAKGQTREPQVYTLAPHRE
ELAKDTVSVTCLVKGFYPPDINVEWQRNGQPESEGTYATTPPQLDNDGTYFL
YSKLSVGKNTWQQGETFTCVVMHEALHNHYTQKSITQSSGK

15

NT2H7 scFv llama IgG3 (nucleotide sequence) (SEQ ID NO: 25)

aagcttgccgcatggatttcaagtcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattcaggggagaaggtcacatgacttcagggccagctcaagtgaag
ttacatgcactggtagcagagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
20 cctgctcgttcagtggcagtggtctgggaccttactcttcacaatcagcagagtgaggagctgaagatgctgccactt
attactgccagcagtgaggatttaacccaccacgcttcggtgctgggaccaagctggagctgaaagatggcggtggtcgc
ggcggtggtgagctggaggaggtgggagctctcaggcttactacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgctcgaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
25 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaacttacttggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaagcg
aaccacagcgaagacccagctccaagtgcccaaatgccaggccctgaactccttgaggggcccacggcttctcatcttc
ccccgaaagccaaggacgtcctctccatcacccgaaaacctgaggtcacgtgcttggtggacgtgggtaaagaagac
cctgagatcgagttcaagctggtccgtggatgacacagaggtacacacggctgagacaaagccaaaggaggaacagttc
30 aacagcacgtaccgctggtcagcgtcctgccatccagcaccaggactggctgacggggaaggaattcaagtgaag
gtcaacaacaaagctctcccagcccccatcgagaggaccatctccaaggccaaagggcagacccgggagccgcaggt
gtacaccctggccccacaccgggaagagctggccaaggacaccgtgagcgtaacctgcctggtcaaaggcttcttccca

gctgacatcaacggtgagtgaggaggaatgggcagccggagtcagagggcacctacgccaacacgccgccacagctg
gacaacgacgggacctacttctctacagaaactctccgtgggaaagaacacgtggcagcagggagaagtcttcacctg
tgtggtgatgcacgaggctctacacaatcactccacccagaaatccatcaccagtccttcgggtaaataagtaatctagagg
ccc

5

AA

2H7 scFv llama IgG3 (amino acid sequence) (SEQ ID NO: 26)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
10 CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSSDQAHSHSEDPSSKCPKCPGPELLGGPTVFIFPPKAKDVLSITRKPEVTC
LWWTWVKKTLRSSSSWSVDDTEVHTAETKPKEEQFNSTYRVVSVLPIQHGD
15 WLTGKEFKCKVNNKALPAPIERTISKAKGQTREPQVYTLAPHREELAKDTS
VTCLVKGFPPADINVEWQRNGQPESEGTYANTPPQLDNDGTYFLYSKLSVGK
NTWQQGEVFTCVVMHEALHNHSTQKSITQSSGK

NT

20 **2H7+Completely WT IgG tail** (nucleotide sequence)

2H7 scFv WTH WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 27)

aagcttgccgcatggatttcaagtcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
25 cctgctcgcttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtcgc
ggcggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
30 actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcaggag

cccaaatcttgtgacaaaactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttctcttcc
 ccccaaaaccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga
 ccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
 caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
 5 gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
 acaccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagc
 gacatgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
 ggctccttctctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
 catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccggtaaatgatctaga

10

AA

2H7 scFv WTH WTCH2CH3 (amino acid sequence) (SEQ ID NO: 28)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
 WYQQKPGSSPKPWYIAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
 15 CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
 VTVSSDQEPKCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV
 VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
 20 WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSL
 TCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRW
 QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

NTCD80 transmembrane domain and cytoplasmic tail (+restriction sites)

25 **(nucleotide sequence) (SEQ ID NO: 29)**

gcggatccttgaacctgctcccatcctgggccattaccttaatctcagtaaagggaattttgtgatatgctgcctgacctact
 gctttgcccccaagatgcagagagagaaggaggaatgagagattgagaagggaaagtgtacgccctgtataaatcatg

AA

CD80 transmembrane domain and cytoplasmic tail (amino acid sequence)

adpsnllpswaitlisvngifviccltycfaprcrerrmerlrresvrpv (SEQ ID NO: 30)

5 NT

40.2.220 VL (anti-human CD40 scFv #1--VL) (nucleotide sequence) (SEQ ID NO: 31)

aagcttatggattttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgttct
gactcagtcctccagccaccctgtctgtgactccaggagatagagtcctctcttctgcagggccagccagagtattagcgac
10 tacttacactgggatcaacaaaaatcacatgagtcctcaaggcttctcatcaaatatgcttccattccatctctgggatccctc
caggttcagtggcagtgatcagggtcagatttcactctcagtatcaacagtggtgaacctgaagatgttggaatttattactg
tcaacatggtcacagctttccgtggacgttcggtggaggcaccaagctggaaatcaaacgg

AA

15 **40.2.220 VL (anti-human CD40 scFv #1--VL) (amino acid sequence)** (SEQ ID NO: 32)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPATLSVTPGDRVSLSCRASQSISDY
LHWYQQKSHESPRLLIKYASHSISGIPSRFSGSGSGSDFTLSINSVEPEDVGIYY
CQHGHSPWTFGGGTKLEIKR

20

NT

40.2.220 VH (for anti-human CD40 scFv #1--VH) (nucleotide sequence) (SEQ ID NO: 33)

cagatccagttggtgcaatctggacctgagctgaagaagcctggagagacagtcaggatctctgcaaggcttctgggtat
25 gccttcacaactactggaatgcagtggggtgcaagagatgccaggaaagggttgaaaggattggttgataaacacccc
actctggagtgccaaaatatgtagaagacttcaaggacggttgccttctcttggaaacctctgccaacactgcatattaca
gataagcaacctcaaagatgaggacacggctacgtattctgtgtgagatccgggaatggttaactatgacctggcctactttg
cttactggggccaaggagactggtcactgtctctgatca

AA

40.2.220 VH (for anti-human CD40 scFv #1--VH) (amino acid sequence) (SEQ ID NO: 34)

QIQLVQSGPELKKPGETVRISCKASGYAFTTTGMQWVQEMPGKGLKWIGWIN
5 TPLWSAKICRRLQGRFAFSLETSANTAYLQISNLKDEDTATYFCVRSNGNYD
LAYFAYWGQGLVTVS

NT

40.2.220 scFv (anti-human CD40 scFv #1) (nucleotide sequence) (SEQ ID NO: 35)

10 aagcttatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgttct
gactcagctctccagccaccctgtctgtgactccaggagatagagtctctcttctcctgcagggccagccagagtattagcgac
tacttacactggatcaacaaaaatcacatgagctctccaaggcttctcatcaaatatgcttccattccatctctgggatccctc
caggttcagtgaggcagtgatcagggtcagatttactctcagtatcaacagtggtgaacctgaagatgttgaatttattactg
tcaacatggtcacagcttccgtggacgttcggtggaggcaccaagctggaaatcaaacggggtggcgggtggctcgggc
15 ggaggtgggtcgggtggcggtgatctcagatccagttggtgcaatctggacctgagctgaagaagcctggagagacag
tcaggatctcctgcaaggcttctgggtatgccttcacaactactggaatgcagtggtgcaagagatgccaggaaagggtt
gaagtggattggctggataaacaccccactctggagtgccaaaatatgtagaagacttcaaggacggttgccttctcttgg
aaacctctgccaaactgcatatttacagataagcaacctcaaagatgaggacacggctacgtatttctgtgtgagatccgg
gaatggtaactatgacctggcctactttgcttactggggccaaggacactggtcactgtctctgatca

20

AA

40.2.220 scFv (anti-human CD40 scFv #1) (amino acid sequence) (SEQ ID NO: 36)

25 MDFQVQIFSLLISASVIMSRGVDIVLTQSPATLSVTPGDRVSLSCRASQSISDY
LHWYQQKSHESPRLLIKYASHSISGIPSRFSGSGSGSDFTLSINSVEPEDVGIYY
CQHGHSFPWTFGGGTKLEIKRGGGGSGGGGSGGGGSIQLVQSGPELKKPGE
TVRISCKASGYAFTTTGMQWVQEMPGKGLKWIGWINTPLWSAKICRRLQGR
FAFSLETSANTAYLQISNLKDEDTATYFCVRSNGNYDLAYFAYWGQGLVTVS

30

NT

2e12 VL (with L6 VK leader peptide) (nucleotide sequence) (SEQ ID NO: 37)

atggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgtcaccc
aatctccagcttctttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaagtggtgaatattatgtca
5 caagttaatgcagtggtaccaacagaaaccaggacagccaccaaaactcctcatctctgtgcatccaacgtagaatctgg
ggtcctgccaggttagtggcagtggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgatattgca
atgtatttctgtcagcaaagtaggaaggttccttgacgttcggtggaggcaccaagctggaaatcaaacgg

AA

10 **2e12 VL (with L6 VK leader peptide) (amino acid sequence) (SEQ ID NO: 38)**

MDFQVQIFSLLISASVMSRGVDIVLTQSPASLAVSLGQRATISCRASESVEYY
VTSLMQWYQQKPGQPPKLLISAASNVESGVPARFSGSGSGTDFSLNIHPVEED
DIAMYFCQQSRKVPWTFGGGTKLEIKR

15 NT

2e12 VH (no leader peptide) (nucleotide sequence) (SEQ ID NO: 39)

caggtgcagctgaaggagtcaggacctggcctggtggcgccctcacagagcctgtccatcacatgcaccgtctcagggtt
ctcattaaccggctatggtgtaaactgggttcgccagcctccaggaaaggtctggagtggctgggaatgatatgggtga
tggaaacacagactataattcagctctcaaaccagactgagcatcaccaaggacaactccaagagccaagttttctaaaa
20 atgaacagtctgcaaactgatgacacagccagatactactgtgccagagatggttatagtaactttcattactatgttatggact
actggggtcaaggaacctcagtcaccgtctcctca(gatctg)

AA

2e12 VH (amino acid sequence) (SEQ ID NO: 40)

25 QVQLKESGPGLVAPSQSL SITCTVSGFSLTGYGVNWVRQPPGKGLEWLGMIW
GDGSTDYNSALKSRLSITKDNSKSQVFLKMNSLQTDDTARYYCARGYSNFH
YYVMDYWGQGTSVTVSS

NT

2e12scFv(+Restriction sites) (nucleotide sequence) (SEQ ID NO: 41)

aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttctttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgtgaatatta
tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
5 tctgggggtccctgccaggtttagtggcagtgggctctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtattttctgtcagcaaagtaggaaggttccttggacgttcgggtggaggcaccaagctggaaatcaaacggggtgg
cgggtggctcgggcgagggtgggtcgggtggcgcggtatcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatgggtgaaactgggttcgccagcct
ccaggaaagggctgtgagtggtgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactg
10 agcatcaccaaggacaactccaagagccaagttttcttaaaaatgaacagctctgcaaactgatgacacagccagataactact
gtgccagagatggttatagtaactttcattactatgttatggactactgggggtcaaggaacctcagtcaccgtctcctct(gatc
ag)

AA

15 **2e12scFv (amino acid sequence) (SEQ ID NO: 42)**

MDFQVQIFSLLISASVIMSRGVDIVLTQSPASLAVSLGQRATISCRASESVEYY
VTSLMQWYQQKPGQPPKLLISAASNVESGVPARFSGSGSGTDFSLNIHPVEED
DIAMYFCQQSRKVPWTFGGGKLEIKRGGGGSGGGGSGGGGSQVQLKESGP
GLVAPSQSLITCTVSGFSLTGYGVNWVRQPPGKGLEWLGMWGDGSTDYNS
20 ALKSRLSITKDNSKSQVFLKMNSLQTTDDTARYYCARDGYSNFHYVMDYW
GQGTSVTVSS

NT

10A8 is anti-CD152 (CTLA-4)

25 **10A8 VL (with L6 VK leader peptide) (nucleotide sequence) (SEQ ID NO: 43)**

atggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacatccagatgacac
agtcctcatcctcactgtctgcatctctgggaggcaagtcaccatcacttgcaaggcaagccaagacattaagaagtatata
ggttgggtaccaacacaagcctggaaaagggtccaggctgctcatatattacacatctacattacagccaggcatcccatcaa
ggttcagtggaagtgggtctgggagagattatccctcagcatcagaaacctggagcctgaagatattgcaacttattattgtc
30 aacagtatgataatcttcattgacgttcggctcggggacaaagttggaaataaaacgg

AA

10A8 VL (amino acid sequence) (SEQ ID NO: 44)

MDFQVQIFSFLISASVIMSRGVDIQMTQSPSSLSASLGGKVTITCKASQDIKK
5 YIGWYQHKPGKGPRLLIYYTSTLQPGIPSRFSGSGSGRDYSLIRNLEPEDIATY
YCQQYDNLPLTFGSGTKLEIKR

NT

10A8 VH (no leader peptide) (nucleotide sequence) (SEQ ID NO: 45)

10 gatgtacagcttcaggagtcaggacctggcctcgtgaaaccttctcagtcctctgtctctcacctgctctgtcactggctactcc
atcaccagtggtttctactggaactggatccgacagtttccgggaaacaaactggaatggatgggccacataagccacgac
ggtaggaataactacaacccatctctcataaatcgaatctccatcactcgtgacacatctaagaaccagttttcctgaagttga
gttctgtgactactgaggacacagctacatatattctgtgcaagacactacggtagtagcggagctatggactactggggtca
aggaacctcagtcaccgtctcctctgatca

15

AA

10A8 VH (amino acid sequence) (SEQ ID NO: 46)

DVQLQESGPGLVKPSQSLTCSVTGYSITSGFYWNWIRQFPGNKLEWMGHIS
HDGRNNYNPSLINRISITRDTSKNQFFLKLSSVTTEDTATYFCARHYGSSGAM
20 DYWGQGTSVTVSS

NT

10A8 scFv (nucleotide sequence) (SEQ ID NO: 47)

aagcttatggatttcaagtcagatttccagcttctgctaatacagtcgttcagtcataatgtccagaggagtcgacatccaga
25 tgacacagtcctccatcctcactgtctgcattctctgggaggcaaaagtcaccatcacttgcaaggcaagccaagacattaagaa
gtatatagggttggtaccaacacaagcctggaaaagggtcccagggtgctcatatattacacatctacattacagccaggcatcc
catcaagggttcagtggaagtgggtctgggagagattattccctcagcatcagaaacctggagcctgaagatattgcaacttat
tattgtcaacagtatgataatcttcattgacgttcggctcggggacaaagtggaaataaaacggggtggcgggtggctcgg
gcgggtgggtgggtcgggtggcggcgatctgatgtacagcttcaggagtcaggacctggcctcgtgaaaccttctcagtcctc

tgtctctcacctgctctgtcactggctactccatcaccagtgggttctactggaactggatccgacagttccgggaacaaac
tggaatggatgggccacataagccacgacggtaggaataactacaacccatctctcataaatcgaatctccatcactcgtga
cacatctaagaaccagtttttctgaagttgagttctgtgactactgaggacacagctacatatttctgtgaagacactacggg
agtagcggagctatggactactgggggtcaaggaacctcagtcaccgtctcctctgatca

5

AA

10A8 scFv (amino acid sequence) (SEQ ID NO: 48)

MDFQVQIFSLLISASVIMSRGVDIQMTQSPSSLSASLGGKVTITCKASQDIKK
YIGWYQHKPGKGPRLLIYYTSTLQPGIPSRFSGSGSGRDYSLIRNLEPEDIATY
10 YCQQYDNLPLTFGSGTKLEIKRGGGGSGGGGSGGGGSDVQLQESGPGLVKPS
QSLSLTCSVTGYSITSGFYWNWIRQFPGNKLEWMGHISHDGRNNYNPSLINRI
SITRDTSKNQFFLKLSSVTTEDTATYFCARHYGSSGAMDYWGQGTSVTVSSD

NT

15 **40.2.220-hmIgG1-hCD80 (nucleotide sequence)** (SEQ ID NO: 49)

aagcttatggatttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgttct
gactcagctctccagccaccctgtctgtgactccaggagatagagtctctcttctgcagggccagccagagtattagcgac
tacttacactggatatcaaaaaatcacatgagtcctcaaggcttctcatcaaatatgcttccattccatctctgggatccctc
caggttcagtggcagtggtatcagggtcagatttcactctcagtatcaacagtggtgaacctgaagatgttggaatttattactg
20 tcaacatggtcacagctttccgtggacgttcggtggaggcaccaagctggaaatcaaacggggtggcgggtggctcgggc
ggaggtgggtcgggtggcggcgatctcagatccagttggtgcaatctggacctgagctgaagaagcctggagagacag
tcaggatctcctgcaaggcttctgggtatgccttcacaactactggaatgcagtggtgcaagagatgccaggaaagggtt
gaagtggattggctggataaacacccactctggagtgcataaatgtagaagacttcaaggacggtttgccttctctttgg
aaacctctgccaacactgcatatttacagataagcaacctcaaagatgaggacacggctacgtatttctgtgtgagatccgg
25 gaatggtaactatgacctggcctactttgcttactggggccaaggacactggcactgtctctgatctggagcccaaatctt
ctgacaaaactcacatccccaccgtccccagcacctgaactcctggggggatcgtcagttctctcttccccccaaaacc
caaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacgaagacctgaggtc
aagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacg
taccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaa
30 agccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcc
cccatccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatcccagcgacatcgccg

tggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctt
cctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctc
tgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacctgctcccatcctgggcca
ttaccttaatctcagtaaattggaatTTTTGTGATATGCTGCCTGACCTACTGCTTTGCCCAAGATGCAGAGAGAGAAGGAGGA
5 atgagagattgagaagggaaagtgtacgccctgtataaatcgat

AA

40.2.220-hmtIgG1-hCD80 (amino acid sequence) (SEQ ID NO: 50)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPATLSVTPGDRVSLSCRASQSISDY
10 LHWYQQKSHESPRLLIKYASHSISGIPSRFSGSGSGSDFTLSINSVEPEDVGIYY
CQHGHSPWTFGGGTKLEIKRGGGGSGGGGSGGGGSQIQLVQSGPELKKPGE
TVRISCKASGYAFTTTGMQWVQEMPGKGLKWIGWINTPLWSAKICRRLQGR
FAFSLETSANTAYLQISNLKDEDTATYFCVRSNGNGNYDLAYFAYWGQGTLLV
VSDLEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVVVD
15 VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLN
GKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL
VKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQG
NVFSCSVMHEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVNGIFVICCLT
YCFAPRCRERRRNERLRRESVRPV

20

NT

2e12scFv- hmtIgG1-CD80 fusion protein (nucleotide sequence) (SEQ ID NO: 51)

aagcttatggattttcaagtcagattttcagcttctgtaaatcagtgcttcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttctttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaagtggtgaatatta
25 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
tctggggtccctgccaggttagtggtcagtggtgctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaagtaggaaggttccttgacgttcggtggaggcaccaagctggaaatcaaacggggtgg
cgggtggctcgggagggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcct
30 ccaggaaagggtctggagtggctgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactg
agcatcaccaaggacaactccaagagccaagttttcttaaaaatgaacagctctgcaaactgatgacacagccagatactact

gtgccagagatgggtatagtaactttcattactatgttatggactactgggggtcaaggaacctcagtcaccgtctcctcagatct
ggagcccaaatcttctgacaaaactcacatccccaccgtccccagcacctgaactcctggggggatcgtagcttctcctc
tccccccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacga
agaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
5 gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga
aggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
tgtacaccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccca
gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
10 tgcagtaggctctgcacaaccactacacgcagaagagcctctcctgtctccggtaaagcggatccttgaacctgctcc
catcctggggccattaccttaatctcagtaaatggaattttgtgatatgctgcctgacctactgctttgccccaaagatgcagaga
gagaaggaggaatgagagattgagaagggaagtgtacgccctgtataaatcgat

AA

15 **2e12scFv- hmtIgG1-CD80 fusion protein (amino acid sequence)** (SEQ ID NO: 52)
MDFQVQIFSLLISASVIMSRGVDIVLTQSPASLAVSLGQRATISCRASESVEYY
VTSLMQWYQQKPGQPPKLLISAASNVESGVPARFSGSGSGTDFSLNIHPVEED
DIAMYFCQQRKVPWTFGGGTKLEIKRGGGGSGGGGSGGGGSQVQLKESGP
GLVAPSQSLITCTVSGFSLTGYGVNWVRQPPGKGLEWLGMWGDGSTDYNS
20 ALKSRLSITKDNSKSQVFLKMNSLQTDDTARYYCARDGYSNFHYVMDYW
GQGTSVTVSSDLEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPE
VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVD
25 KSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVN
GIFVICCLTYCFAPRCRERRRNERLRRESVRPV

NT

10A8 scFv-hmtIgG1-CD80 (nucleotide sequence) (SEQ ID NO: 53)
30 aagcttatggatttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacatccaga
tgacacagttccatctcactgtctgcatctctgggaggcgaagtcaccatcacttgaaggcaagccaagacattaagaa

gtatataggttggtaccaacacaagcctggaaaagggtcccaggctgctcatatattacacatctacattacagccaggcatcc
 catcaaggttcagtggaagtgggtctgggagagattattccctcagcatcagaaacctggagcctgaagatattgcaacttat
 lattgtcaacagtatgataatcttccattgacgttcggctcggggacaaagttggaaataaaacggggtggcggtggctcgg
 gcggtgggtgggtcgggtggcggcggtatctgatgtacagcttcaggagtcaggacctggcctcgtgaaaccttctcagtctc
 5 tgtctctcacctgctctgtcactggctactccatcaccagtggtttctactggaactggatccgacagttccgggaaacaaac
 tggaatggatgggccacataagccacgacggttaggaataactacaacctctctcataaatgaatctccatcactcgtga
 cacatctaagaaccagtttttctgaagttgagttctgtgactactgaggacacagctacataatttctgtgcaagacactacggt
 agtagcggagctatggactactggggtcaaggaacctcagtcaccgtctcctctgatctggagcccaaatctcttgacaaaa
 ctcacacatccccaccgtccccagcacctgaactcctggggggatcgtcagttctcttccccccaaaacccaaggaca
 10 ccctcatgatctcccgaccctgaggtcacatgcgtgggtggacgtgagccacgaagacctgaggtcaagttcaact
 ggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtg
 gtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccc
 agccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatcccg
 ggatgagctgaccaagaaccaggtcagcctgacctggtcaaaggcttctatcccagcgacatcgccgtggagtgagg
 15 agagcaatgggcagccggagaacaactacaagaccacgcctcccgtgtggactccgacggctccttcttctctacagc
 aagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaacca
 ctacacgcagaagagcctctccctgtctccgggtaaagcggatccttgaacctgctcccatcctgggccattaccttaatct
 cagtaaatggaattttgtgatatgctgcctgacctactgctttgccccaaagatgcagagagagaaggaggaatgagagatt
 gagaagggaagtgtagccctgtataaatcgat

20

AA

10A8 scFv-hm1gG1-CD80 (amino acid sequence) (SEQ ID NO: 54)

MDFQVQIFSLLISASVIMSRGVDIQMTQSPSSLSASLGGKVTITCKASQDIKK
 YIGWYQHKPGKGPRLLIYYTSTLQPGIPSRFSGSGSGR DYSL SIRNLEPEDIATY
 25 YCQQYDNLPLTFGSGTKLEIKRGGGSGGGGSGGGGSDVQLQESGPGLVKPS
 QSLSLTCSVTGYSITSGFYWNWIRQFPGNKLEWMGHISHDGRNNYNPSLINRI
 SITRDTSKNQFFLKLSSVTTEDTATYFCARHYGSSGAMDYWGQGTSVTVSSD
 LEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVVVDVSH
 EDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
 30 YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKG
 FYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF

SCSVMHEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVNGIFVICCLTYCF
APRCRERRRNERLRRESVRPV

NT

5 **500A2-hmtIgG1-CD80 (nucleotide sequence)** (SEQ ID NO: 55)

atgttgatacatctcagctccttgggctttactcttctggatttcagcctccagaagtacatagtctgactcagactccagc
cactctgtctctaattcctggagaaagagtcacaatgacctgtaagaccagtcagaatattggcacaatcttactggtatca
ccaaaaaccaaaggaggctccaagggtctcatcaagtatgcttcgcagtccattcctgggatccccctccagattcagtggc
agtgggttcggaacagatttcactctcagcatcaataacctggagcctgatgatcggaaattattactgtcaacaaagtaga
10 agctggcctgtcacgttcggctcctggcaccaagctggagataaaacggggtggcgggtggctcggcgagggtgggtcg
gggtggcggcgatctcaggtcaagctgcagcagtcgggttctgaactagggaaacctggggcctcagtgaactgtcctg
caagacttcaggctacatattcacagatcactatatttctgggtgaaacagaagcctggagaaagcctgcagtggatagga
aatgtttatggtgaaatgggtgtacaagctacaatcaaaaattccagggaaggccacactgactgtagataaaatctctag
cacagcctacatggaactcagcagcctgacatctgaggattctgccatctattactgtgaagaaggccggtagcgacggg
15 ccatgctatggactactggggtcaggggatccaagttaccgtctcctctgatctggagcccaaatcttctgacaaaactcaca
catccccaccgtccccagcacctgaactcctggggggatcgtcagttcttcttcccccaaaacccaaggacaccctca
tgatctcccgacccttgaggtcacatgcgtggtgggtggacgtgagccacgaagaccctgaggtcaagttcaactggtac
gtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagc
gtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtcaaggtctccaacaaagccctcccagcccc
20 catcgagaaaaccatctccaaagccaaaggcgagccccgagaaccacaggtgtacaccctgccccatcccgggatga
gctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagc
aatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagcaagct
caccgtggacaagagcaggtggcagcaggggaacgttctcatgctccgtgatgcatgaggctctgcacaaccactaca
cgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacctgtccccatcctgggccattaccttaatctcagta
25 aatggaattttgtgatatgctgcctgacctactgcttggcccaagatgcagagagagaaggaggaatgagagattgagaa
gggaaagtgtacgccctgtataaatcgat

AA

30 **500A2-hmtIgG1-CD80 (amino acid sequence)** (SEQ ID NO: 56)

MLYTSQLLGLLLFWISASRSDIVLTQTPATLSLIPGERVTMTCKTSQNIGTILH
WYHQKPKEAPRALIKYASQSIPGIPSRFSGSGSETDFTLSINNLEPDDIGIYYCQ

QSRSWPVTFPGGTKLEIKRGGGGSGGGGSGGGGSQVKLQQSGSELGKPGASV
KLSCKTSGYIFTDHYISWVKQKPGESLQWIGNVYGGNGGTSYNQKFQ GKATL
TVDKISSTAYMELSSLTSEDSAIYYCARRPVATGHAMDYWGQGIQVTVSSDL
EPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVVVDVSHE
5 DPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEY
KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGF
YPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFS
CSVMHEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVNGIFVICCLTYCFA
PRCRERRRNERLRRESVRPV

10

NT

2H7 scFv MTH(SSS)WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 57)

aagcttgcgcccatggatttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatctgtctgcatctccaggggagaaggtcacaatgactgcagggccagctcaagtgaag
15 ttacatgcactggtaccagcagaagccaggatcctcccccaccctggatttatcccccattcaacctggcttctggagtc
cctgctcgttcagtggtcagtggtctgggaccttactctctcacaatcagcagagtggaggctgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctc
ggcggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
20 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgacaggag
cccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagcttctcttcc
cccaaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga
25 ccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagc
gacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgac
30 ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

2H7 scFv MTH(SSS)WTCH2CH3 (amino acid sequence) (SEQ ID NO: 58):

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
5 CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV
VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
10 LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

15 **HuIgGMHncs1 (oligo for CSS) (nucleotide sequence)** (SEQ ID NO: 550)

gtt gtt gat cag gag ccc aaa tct tgt gac aaa act cac

NT

HuIgGMHncs2 (oligo for SCS=ncs2) (nucleotide sequence) (SEQ ID NO: 549)

20 gtt gtt gat cag gag ccc aaa tct tct gac aaa act cac aca tgc cca ccg

NT

HuIgGMHncs3 (oligo for SSC=ncs3) (nucleotide sequence) (SEQ ID NO: 548)

gtt gtt gat cag gag ccc aaa tct tct gac aaa act cac aca tct cca ccg tgc cca gca cct g

25

NT

hIgGWT3xba (3' oligo for above mutation introduction) (nucleotide sequence)
(SEQ ID NO: 551)

gtt gtt tct aga tca ttt acc cgg aga cag gga gag gct ctt ctg cgt gta g

NT

Vhser11: (oligo for Leu to Ser at VH11) (nucleotide sequence) (SEQ ID NO: 552)

5 gga ggt ggg agc tct cag gct tat cta cag cag tct ggg gct gag tcg gtg agg cc

NT

huIgG1-3' (3' oligo to amplify IgG1 C regions, 3' end of CH3) (nucleotide sequence) (SEQ ID NO: 553)

10 gtc tct aga cta tca ttt acc cgg aga cag

NT

huIgA/Gchim5 (oligo for pcr#1) (nucleotide sequence) (SEQ ID NO: 554)

cca tct ccc tca act cca cct acc cca tct ccc tca tgc gca cct gaa ctc ctg
15

NT

huIgAhg-5' (oligo for pcr#2) (nucleotide sequence) (SEQ ID NO: 555)

gtt gtt gat cag cca gtt ccc tca act cca cct acc cca tct ccc caa ct

20 NT

huIgA3' (nucleotide sequence) (SEQ ID NO: 557)

gtt gtt tct aga tta tca gta gca ggt gcc gtc cac ctc cgc cat gac aac

NT

2H7 scFv IgAH IGG WT CH2CH3 (2H7 scFv with IgA hinge and WT CH2 and CH3) (nucleotide sequence) (SEQ ID NO: 59)

aagcttgccgcatggattttcaagtgcagattttcagcttcctgctaatacagtgcttcagtcataattgccagaggacaaattg
5 ttctctcccagctctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagttaag
ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgtgccactt
attactgccagcagtgagggtttaaccacccacgttcgggtgctgggaccaagctggagctgaaagatggcgggtggctcg
ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
10 cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagtcaagggcaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcgggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactgtgtacttcgatgtctggggcacagggaccacggtcaccgtctctgatcagccagtt
ccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgcgcacctgaactcctggggggaccgt
15 cagtcttctcttccccccaaaacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtgggtgggtggacgt
gagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcg
ggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactgggtgaatggcaaggag
tacaagtgaaggtctccaacaaagccctccagcccccacagagaaacaatctccaaagccaaagggcagccccgag
aaccacaggtgtacacctgcccccatcccggtatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaagg
20 cttctatcccagcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccggtg
ctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctca
tgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

25 **2H7 scFv IgAH IGG WT CH2CH3** (amino acid sequence) (SEQ ID NO: 60)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
30 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSDQVPVSTPPTPSPSTPPTPSPSCAPELLGGPSVFLFPPKPKDTLMISRTPEV
TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL

HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDK
SRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

5 NT

2H7 scFv IgAH IgACH2CH3 (2H7 scFv IgAhinge and IgA CH2 and CH3)

(nucleotide sequence) (SEQ ID NO: 61)

aagcttgccgccatggattttcaagtgcagattttcagcttcctgtaatcagtgccttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
10 ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgccagtggtctgggacctcttactctctcacaatcagcagagtggaggctgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
ggcggtgggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaagatgctcgaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
15 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcagcca
gttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtcactgca
ccgaccggccctcgaggacctgctcttaggttcagaagcgtacgtgcacactgaccggcctgagagatgcctcag
20 gtgtcaccttcacctggacgccctcaagtgggaagagcgtgttcaaggaccacctgaccgtgacctctgtggctgtaca
gcgtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttcagctgctgacctaccccgagt
ccaagacccccgtaaccgccaccctctcaaaatccggaacacattccggcccagaggtccacctgtgcgcgccgctc
ggaggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagccccaaggatgtgctggttcgct
ggctgcaggggtcacaggagctgccccgcgagaagtacgtgacttgggcatcccggcaggagcccagccagggcacc
25 accaccttcgtgtgaccagcactgctgcgtggcagccgaggactggaagaagggggacaccttctctgcatggtgg
gccacgagggcctgccgctggccttcacagaagaccatcgaccgcttggcgggtaaacccacctgtcaatgtgtct
gttgctcatggcgagggtggacggcacctgctactgataatctaga

AA

30 **2H7 scFv IgAH IgACH2CH3 (2H7 scFv IgA hinge and IgA CH2 and CH3)**

(amino acid sequence) (SEQ ID NO: 62)

MDFQVQIFSFLNISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
 CQQWSFNPPFTGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
 5 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
 VTVSSDQVPSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLT
 GLRDASGVTFWTWPSSGKSAVQGPPDRDLCGCYSVSSVLPGCAEPWNHGKTF
 TCTAAYPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFS
 PKDVLVRWLQGSQELPREKYLTWASRQEPSQGTTTTFAVTSILRVA AEDWKK
 10 GDTFSCMVGHEALPLAFTQKTIDRLAGKPTHVNVSVVMAEVDGTCY

NT

IgA hinge-CH2-CH3 (Human IgA tail, full length) (nucleotide sequence) (SEQ ID NO: 63)

15 tgatcagccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgac
 tgctactgcaccgaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgaga
 gatgcctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccacctgaccgtgacctctgt
 ggctgctacagcgtgtccagtgctcctgccgggctgtgccgagccatggaacctgggaagaccttcacttgactgctgcc
 taccctgagtgccaagacccccgtaaccgccaccctctcaaaatccggaaacacattccggccccgaggtccacctgctgcc
 20 gccgccgtcggaggagctggccctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagcccaaggatgtg
 ctgggtcgtggctgcaggggtcacaggagctccccgcgagaagtacctgacttgggcatcccggcaggagcccagc
 cagggcaccaccaccttcgctgtgaccagcatactgcgcgtggcagccgaggactggaagaagggggacaccttctct
 gcatggtggggcacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgt
 caatgtgtctgtgtcatggcgagggtggacggcacctgctactgataatctaga

25

AA

IgA hinge-CH2-CH3 Protein sequence, (Human IgA tail, full length) (amino acid sequence) (SEQ ID NO: 64)

DQPVPSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLTGLRDA
 30 SGVTFTWTPSSGKSAVQGPPDRDLCGCYSVSSVLPGCAEPWNHGKTFTCTAA
 YPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFSPKDVL

VRWLQGSQELPREKYLTWASRQEPSQGTTFVAVTSILRVAEDWKKGDTFSC
MVGHEALPLAFTQKTIDRLAGKPTHVNVSVVMAEVDGTCY

NT

5 **Human J Chain (nucleotide sequence)** (SEQ ID NO: 65)

agatctcaagaagatgaaaggattgttcttggtgacaacaaatgtaagtgtgcccggattacttccaggatcatccgttctcc
gaagatcctaatgaggacattgtggagagaaacatccgaattattgttcctctgaacaacaggagagaatatctctgatccac
ctcaccattgagaaccagatttgtgtaccattgtctgacctcagctgtaaaaaatgtgacacctacagaagtggagctggataa
tcagatagttactgctaccagagcaatatctgtgatgaagacagtgtacagagacctgctacacttatgacagaaacaag
10 tctacacagctgtggtccactcgtatatggtggtgagacaaaatggtggaacagccttaaccccagatgcctgctatc
ctgactaatctaga

AA

Human J Chain (amino acid sequence) ((SEQ ID NO: 66)

15 RSQEDERIVLVDNKCKCARITSRIIRSSDPNEDIVERNIRIIVPLNNRENISDPTS
PLRTRFVYHLSDLSCKKCDPTEVELDNQIVTATQSNICDEDSATETCYTYDRN
KCYTAVVPLVYGGETKMOVETALTPDACYP

NT

20 **HUJCH5nl (J chain 5' primer)** (nucleotide sequence) (SEQ ID NO: 558)

gtt gtt aga tct caa gaa gat gaa agg att gtt ctt

NT

HUJCH3 (J chain 3' primer-antisense) (nucleotide sequence) (SEQ ID NO: 559)

25 gtt gtt tct aga tta gtc agg ata gca ggc atc tgg

AA

4 carboxy terminal amino acids deleted from IgA CH3 (amino acid sequence)

(SEQ ID NO: 67)

GTCY

5

NT

IgAH IgAT4 Human IgA tail, truncated (3T1)-(missing last 4 amino acids from carboxy terminus) (nucleotide sequence) (SEQ ID NO: 68)

tgatcagccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgac
10 tgtcactgcaccgaccggccctcgaggacgtgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgaga
gatgcctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccacctgaccgtgacctgt
ggctgctacagcgtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttgactgctgcc
taccgagtcgaagacccccgtaaccgccacccctctcaaatccggaacacattccggcccgagggtccacctgctgcc
gcccgcgtcggaggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtg
15 ctggttcgctggctgcaggggtcacaggagctccccgcgagaagtacctgacttgggcatcccggcaggagcccagc
cagggcaccaccaccttcgctgtgaccagcatactgcgcgtggcagccgaggactggaagaagggggacaccttctct
gcatggtgggccacgagggcctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgt
caatgtgtctgtgtcatggcggagggtggactgataatctaga

20 AA

IgAH IgAT4 (amino acid sequence) (SEQ ID NO: 69)

DQVPVSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLTGLRDA
SGVTFTWTPSSGKSAVQGPPDRDLGCGYSVSSVLPGCAEPWNHGKTFTCTAA
YPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFSPKDVL
25 VRWLQGSQELPREKYLTWASRQEPSQGTTFVAVTSILRVAAEDWKKGDTFSC
MVGHEALPLAFTQKTIDRLAGKPTHVNVSVVMAEVD

NT

HUIGA3T1 (oligo 3': to delete 4 amino acids at carboxy end of IgA CH3) (SEQ
30 ID NO: 562)

gtt gtt tct aga tta tca gtc cac ctc cgc cat gac aac aga cac

NT

HUIGA3T2 (oligo to delete 14 aa at end of IgA –T4) (SEQ ID NO: 565)

5 gtt gtt tct aga tta tca ttt acc cgc caa gcg gtc gat ggt ctt

NT

2H7 scFv IgAH IgAT4 (2H7 scFv IgA 3T1 construct)--truncates the CH3 domain at the 3'end (nucleotide sequence) (SEQ ID NO: 70)

10 aagcttgccgcatggattttcaagtcagattttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattctccaggggagaaggtcacaatgactgcagggccagctcaagtgaag
ttacatgcactgggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggacaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
15 ggcggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacatttaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtaacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcagcca
20 gttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtcactgca
ccgaccggccctcaggacctgctcttaggttcagaagcgtacgtgcacactgaccggcctgagagatgcctcag
gtgtcaccttcacctggacgccctcaagtgggaagagcgtgttcaaggaccacctgaccgtgacctctgtggctgtaca
gcgtgtccagtgtcctgccgggtgtgccgagccatggaacctgggaagaccttcacttgactgctgcctaccccgagt
ccaagacccccgtaaccgccaccctctcaaaatccggaacacattccggcccgagggtccacctgctgccgccgctc
25 ggaggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagccccaaagatgtgctggttcgct
ggctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccgagcaggagcccagccagggcacc
accaccttcgctgtgaccagcactgctgcgtggcagccgaggactggaagaagggggacaccttctcctgcatggtgg
gccacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgtcaatgtgtct
gttgtcatggcgagggtgactgataatctaga

30

AA

2H7 scFv IgAH-T4 (amino acid sequence) (SEQ ID NO: 71)

MDFQVQIFSFLNISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
5 CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSSDQVPSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLT
GLRDASGVTFWTWPSSGKSAVQGPDRDLCGCYSVSSVLPGCAEPWNHGKTF
10 TCTAAYPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFS
PKDVLVRWLQGSQELPREKYLTWASRQEPSQGTTTTFAVTSILRVAAEDWKK
GDTFSCMVGHEALPLAFTQKTIDRLAGKPTHVNVSVVMAEVD

AA

- 15 **14 amino acids deleted from IgAH-T4 (so that total of 18 amino acids deleted from wild type IgA CH3 (amino acid sequence) (SEQ ID NO: 72)**

PTHVNVSVVMAEVD

NT

- 20 **IgAH IgA-T18 (human IgA Tail truncated, 3T2) (nucleotide sequence) (SEQ ID NO: 73)**

Tgatcagccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccga
ctgtcactgcaccgaccggccctcgaggacgtgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgag
agatgcctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccctgaccgtgacctctg
25 tggctgctacagcgtgtccagtgtcctgccgggctgtgccgagccatggaacatgggaagaccttcacttgactgctgc
ctaccccgagtccaagaccccgtaaccgccaccctctcaaatccggaaacacattccggcccgaggtccacctgctgc
cgccgccgtcgaggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagccccaaggatgt
gctggttcgctggctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccggcaggagcccag
ccagggcaccaccaccttcgctgtgaccagcactgctgcgtggcagccgaggactggaagaagggggacaccttctcc
30 tgcattggtgggcacagggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaa

AA

IgAH IgA-T18 (amino acid sequence) (SEQ ID NO: 74)

DQVPVSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLTGLRDA
5 SGVTFTWTPSSGKSAVQGPPDRDLCGCYSVSSVLPGCAEPWNHGKTFTCTAA
YPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFSPKDV
VRWLQGSQELPREKYLTWASRQEPSQGTTFFAVTSILRVAEDWKKGDTFSC
MVGHEALPLAFTQKTIDRLAGK

10 NT

2H7 scFv IgAH IgAT18 (human IgA Tail truncated, 3T2) (nucleotide sequence)
(SEQ ID NO: 75)

aagcttggcccatggattttcaagtcagattttcagcttctgctaatacagtcgttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttcagggccagctcaagtgaag
15 ttacatgcactggtaccagcagaagccaggatcctccccaaaccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtcggcagtcgggtctgggaccttactctctcacaatcagcagagtcggaggctgaagatgctgccactt
attactgccagcagtcggagtttaacccaccacgttcggtgctgggaccaagtcggagctgaaagatggcggctggctc
ggcgggtggatctggaggagtcgggagctctcaggcttatctacagcagtcctggggctgagctggtgaggcctgggg
cctcagtcgaagatgtcctgcaaggcttctggctacacattaccaggttacaatatgcactgggtaaagcacacactagaca
20 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagtcagcagcctgacatctgaagactctgcggctctatttctgtgcaa
gagtcggtgtactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcagcca
gttcctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtcactgca
ccgaccggccctcaggagcctgctcttaggttcagaagcgtacacgtgcacactgaccggcctgagagatgcctcag
25 gtgtcaccttcacctggacgccctcaagtcgggaagagcgtgttcaaggaccacctgaccgtgacctctgtggctgtaca
gcgtgtccagtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttcactgctgcctaccccagtc
ccaagacccccgctaaccgccacctctcaaaatccggaaacacattccggcccagggtccacctgctgccgccgccgtc
ggaggagctggccctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagcccaaggatgtgctggttcgct
ggctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccgagcaggagcccagccagggcacc
30 accaccttcgctgtgaccagcactgctgcgctggcagccgaggactggaagaagggggacaccttctcctgcatggtgg
gccacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaa

AA

2H7 scFv IgAH IgAT18 (amino acid sequence) (SEQ ID NO: 76)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
5 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSSDQVPSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLT
10 GLRDASGVTFWTWPSSGKSAVQGPPDRDLCGCYSVSSVLPGCAEPWNHGKTF
TCTAAYPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFS
PKDVLVRWLQGSQELPREKYLTWASRQEPSQGTTTFAVTSILRVAAEDWKK
GDTFSCMVGHEALPLAFTQKTIDRLAGK

15 NT

CTLA-4 IgG WTH WTCH2CH3 (human-oncoMLP-CTLA4EC-hIgGWT)
(nucleotide sequence) (SEQ ID NO: 77)

gcaacctacatgatggggaatgagttgaccttctagatgattccatctgcacgggcacctccagtggaaatcaagtgaacc
tcactatccaaggactgagggccatggacacgggactctacatctgcaaggtggagctcatgtaccaccgccatactacc
20 tgggcataggcaacggaacccagatttatgtaattgatccagaaccgtgcccagattctgatcaacccaaatcttgacaa
aactcacacatgcccaccgtgcccagcacctgaactcctggggggaccgtcagttctctcttcccccaaaaccaagg
acacctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttc
aactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgt
gtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccct
25 cccagcccccatcgagaaaacaatctcaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatc
ccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagt
gggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttcttctctac
agcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaa
ccactacacgcagaagagcctctccctgtctccgggtaaatga

30

AA

CTLA-4 IgG WTH WTCH2CH3 (amino acid sequence) (SEQ ID NO: 78)

MGVLLTQRTLLSLVLALLFPSMASMAMHVAQPAVVLASSRGIASFVCEYASP
GKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDDSICTGTSSGNQVNL
5 TIQGLRAMDTGLYICKVELMYPPPYLIGINGTQIYVIDPEPCPDSQPKSCD
KTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKF
NWXVVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN
KALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAV
EWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHE
10 ALHNHYTQKSLSLSPGK

NT

Human OncoM leader Peptide+CTLA4 EC (BclI) (nucleotide sequence) (SEQ ID NO: 79)

15 Atgggggtactgctcacacagaggacgtgctcagctcgttcctgcactcctgttccaagcatggcgagcatggcaatg
cacgtggcccagcctgctgtggtactggccagcagccgaggcatgccagcttgtgtgtgagtatgcattccaggcaaa
gccactgagggtccgggtgacagtgttcggcaggctgacagccaggtgactgaagtctgtcggcaacctacatgatgg
ggaatgagttgaccttctagatgattccatctgcacgggcacctccagtggaaatcaagtgaacctcactatccaaggact
gagggccatggacacgggactctacatctgcaaggtggagctcatgtaccaccgccatactacctgggcataggcaac
20 ggaaccagatttatgtaattgatccagaaccgtgccagattctgatcaa

AA

Human OncoM leader Peptide+CTLA4 EC (amino acid sequence) (SEQ ID NO: 80)

25 MGVLLTQRTLLSLVLALLFPSMASMAMHVAQPAVVLASSRGIASFVCEYASP
GKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDDSICTGTSSGNQVNL
TIQGLRAMDTGLYICKVELMYPPPYLIGINGTQIYVIDPEPCPDSQ

NT

Human OncoM leader (nucleotide sequence) (SEQ ID NO: 81)

atgggggtactgctcacacagaggacgctgctcagctctggccttgactcctgtttccaagcatggcgagcatg

5 AA

Human OncoM leader (amino acid sequence) (SEQ ID NO: 82):

MGVLLTQRTL LSLVLALLFPSM

NT

10 **Human CTLA4 EC (no LP) (nucleotide sequence)** (SEQ ID NO: 83)

Gcaatgcacgtggcccagcctgctgtgtgtactggccagcagccgaggcatcggcagctttgtgtgtgagtatgcattcca
ggcaaagccactgaggtccgggtgacagtgttcggcaggctgacagccaggtgactgaagtctgtcggcaacctaca
tgacggggaatgagttgaccttctagatgattccatctgcacgggcacctccagtggaaatcaagtgaacctcactatcca
aggactgagggccatggacacgggactctacatctgcaaggtggagctcatgtaccaccgccatactacctgggcatag
gcaacggaaccagatttatgtaattgatccagaaccgtgccagattct

AA

Human CTLA4 EC (no LP) (amino acid sequence) (SEQ ID NO: 84)

AMHVAQPAVV LASSRG IASFVCEYASPGKATEVRVTVLRQADSQVTEVCAA
20 TYMTGNELTFLDDSICTGTSSGNQVNLTIQGLRAMDTGLYICKVELMYPPPPYY
LGIGNGTQIYVIDPEPCPDS

NT

25 **Human CTLA4 IgG MTH (SSS) MTCH2CH3 (nucleotide sequence)** (SEQ ID NO: 85)

Atgggggtactgctcacacagaggacgctgctcagctctggccttgactcctgtttccaagcatggcgagcatggcaatg
cacgtggcccagcctgctgtgtgtactggccagcagccgaggcatcggcagctttgtgtgtgagtatgcattccaggcaaa
gccactgaggtccgggtgacagtgttcggcaggctgacagccaggtgactgaagtctgtcggcaacctacatgatgg

ggaatgagttgaccttctagatgattccatctgcacgggcacctccagtggaaatcaagtgaacctcactatccaaggact
gagggccatggacacgggactctacatctgcaaggtggagctcatgtaccacccgccatactacctgggcataggcaac
ggaaccagatttatgtaattgatccagaaccgtgcccagattctgatcaacccaaatcttctgacaaaactcacacatccc
accgtccccagcacctgaactcctggggggatcgtcagtcttctcttccccccaaaaccaaggacacctcatgatctcc
5 cggaccctgaggtcacatgcgtgggtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacg
gcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctca
ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcga
gaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccggtgatgagctgac
caagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatggg
10 cagccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttcttctctacagcaagctcaccgtg
gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcaga
agagcctctccctgtctccgggtaaatga

AA

15 **Human CTLA4 IgG MTH (SSS) MTCH2CH3** (SEQ ID NO: 86)

MGVLLTQRTL LSLVLALLFPSMASMAMHVAQPAVVLASSRGIASFVCEYASP
GKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDDSICTGTSSGNQVNL
TIQGLRAMDTGLYICKVELMYPPPYLIGINGTQIYVIDPEPCPDSQPKSSDK
THTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN
20 WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNK
ALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVE
WESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEA
LHNHYTQKSLSLSPGK

25 NT

CTLA-4 IgAH IgACH2CH3 (human-oncoMLP-CTLA4EC-IgA) (nucleotide
sequence) (SEQ ID NO: 87)

atgggggtactgctcacacagaggacgctgctcagtctggctcctgcactcctgtttccaagcatggcgagcatggcaatgc
acgtggcccagcctgctgtggtactggccagcagccgagggcatgccagcttgtgtgtgagtatgcatctccaggcaaa
30 ccactgaggtccgggtgacagtgttcggcagggtgacagccaggtgactgaagtctgtcggcaacctacatgatggg
gaatgagttgaccttctagatgattccatctgcacgggcacctccagtggaaatcaagtgaacctcactatccaaggactg

gaaccagatttatgtaattgatccagaaccgtgccagattctgatcagccagttccctcaactccacctaccccatctccct
 caactccacctaccccatctccctcatgctgccacccccgactgtcactgcaccgaccggccctcgaggacctgctcttag
 gttcagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcagggtgcacctcacctggacgccctcaagt
 ggaagagcgctgttcaaggaccacctgaccgtgacctctgtggctgctacagcgtgtccagtgctctgccgggctgtgcc
 5 gagccatggaacctgggaagaccttcacttgcaactgctgcctaccccgagtccaagaccccgctaaccgccaccctctc
 aaaatccggaaacacattccggcccaggtccacctgctgccgccgccgtcggaggagctggccctgaacgagctggt
 gacgctgacgtgcctggcacgtggcttcagccccaaggatgtgctggttcgctggctgcaggggtcacaggagctgcc
 cgcgagaagtacctgacttgggcatcccggcaggagcccagggcaccaccaccttcgtgtgaccagcatactgc
 gcgtggcagccgaggactggaagaagggggacaccttctcctgcatggtgggccacgaggccctgccgtggccttca
 10 cacagaagaccatcgaccgcttggcgggtaaaccacccatgtcaatgtgtctgttgcattggcggaggtggactgataat
 ctaga

AA

CTLA-4 IgAH IgA-T4 (amino acid sequence) (SEQ ID NO: 90)

15 Mgvlitqrllslvlallfpsmasmamhvaqpavvlassrgiasfvceyaspgkatevrvtvlrquadsqvtevcaatym
 mgneltfddsiactgtssgnqvnltiqglramdtglyickvelmypppyylgigngtqiyyidpepcpsdqpvpstpp
 tpspstptpspscchprlslhrpaledlllgseailtctltglrdasgvtftwtpssgksavqgppdrdlcgcysvssvlpgc
 aepwnhgtftictaaypesktpltatlsksgntfrpevhllpppseelalnelvtltclargfspkdvlvrlqgsqelprek
 yltwasrqepsqgtttfavtsilrvaaedwkkgdtfscmvghaelplafktidrlagkpthvnnsvvmaevd

20

NT

human IgG1 CH2 with 238 mutation pro→ser (nucleotide sequence) (SEQ ID NO: 91)

cctgaactcctggggggatcgtcagttctctcttcccccaaaaccaaggacacctcatgatctcccggacccctgag
 25 gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactgtacgtggacggcgtggaggtgc
 ataatgccaagacaaagccgaggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
 ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacctctcc
 aaagccaaag

AA

human IgG1 CH2 with 238 mutation pro→ser (amino acid sequence) (SEQ ID NO: 92)

PELLGGSSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE
5 VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK
TISKAK

AA

Amino acids surrounding Pro to Ser in CH2 (amino acid sequence) (SEQ ID NO:
10 93)

PAPELLGGPS

AA

Amino acids surrounding Pro to Ser in CH2 (amino acid sequence) (SEQ ID NO:
15 94)

PAPELLGGSS

NT

hIgE5Bcl (nucleotide sequence) (SEQ ID NO: 575)

20 gtt gtt gat cac gtc tgc tcc agg gac ttc acc cc

NT

hIgE3stop (nucleotide sequence) (SEQ ID NO: 576)

gtt gtt tct aga tta act ttt acc ggg att tac aga cac cgc tcg ctg g

25

NT

hIgE3BB (leaves an open reading frame at end of gene to read into transmembrane and cytoplasmic tail domain attached at either the BamHI or SfuI sites) (nucleotide sequence) (SEQ ID NO: 95)

5 gtt gtt ttc gaa gga tcc gct tta ccg gga ttt aca gac acc gct cgc tgg

NT

human IgE Fc (CH2-CH3-CH4) ORF (SEQ ID NO: 96):

tgatcacgtctgctccagggacttcaccccgcccaccgtgaagatcttacagtcgtcctgcgacggcggcgggcacttccc
10 cccgaccatccagctcctgtgcctcgtctctgggtacacccagggactatcaacatcacctggctggaggacgggcagg
tcatggacgtggacttgtccaccgcctctaccacgcaggaggggtgagctggcctccacacaaagcgagctcacctcagc
cagaagcactggctgtcagaccgcacctacacctgccaggtcacctatcaagggtcacacctttgaggacagcaccaagaa
gtgtgcagattccaacccgagaggggtgagcgcctacctaagccggcccagcccgttcgacctgttcacccaagtcgc
ccacgatcacctgtctggtggtggacctggcaccagcaaggggaccgtgaacctgacctggtcccgggccagtgggaa
15 gcctgtgaaccactccaccagaaaggaggagaagcagcgcgaatggcacgttaaccgtcacgtccacctgccggtggg
caccggagactggatcgaggggggagacctaccagtgcagggtgacccaccccccacctgccagggccctcatgcggtc
cacgaccaagaccagcggcccgctgctgccccggaagtctatgcgtttgcgacgccggagtggccggggagccggg
acaagcgcacctcgcctgcctgatccagaacttcagcctgaggacatctcgggtgcagtggctgcacaacgaggtgcag
ctcccgagcggccggcacagcacgacgcagccccgcaagaccaagggctccggcttcttcgtcttcagccgcctggag
20 gtgaccagggccgaatgggagcagaaagatgagttcatctgccgtgcagtcctgaggcagcagccccctcacagacc
gtccagcgagcgggtgtctgtaaattcccggtaaagcggatccttcgaa

AA

human IgE Fc (CH2-CH3-CH4) ORF (amino acid sequence) (SEQ ID NO: 97)

25 DHVCSRDFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTINITWLEDGQV
MDVDLSTASTTQEGELASTQSELTLSQKHWLSDRITYTCQVITYQGHTFEDSTK
KCADSNPRGVSAYLSPSPFDLFIRKSPTITCLVVDLAPSKGTVNLTWSRASGK
PVNHSTRKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTHPHLPRALMRS
TTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDISVQWLHNEVQ
30 LPDARHSTTQPRKTKSGGFFVFSRLEVTRAWEQKDEFICRAVHEAASPSQTV
QRAVSVNPGKADPS

NT

IFhIgGwtBcl5 (nucleotide sequence) (SEQ ID NO:98)

5 gtt gtt tga tca gga gcc caa atc ttg tga caa aac tca cac atg ccc acc gtg ccc agc acc (63
mer)

NT

hIgGWT3xba (nucleotide sequence) (SEQ ID NO: 551)

10 gtt gtt tct aga tca ttt acc cgg aga cag gga gag gct ctt ctg cgt gta g

NT

**HulgGMHWC (sense, 5' primer for mutating wild type hinge CCC to mutant
SSS) (nucleotide sequence) (SEQ ID NO:99)**

15 gtt gtt gat cag gag ccc aaa tct tct gac aaa act cac aca tct cca ccg tcc cca gca cct gaa ctc
ctg ggt gga ccg tca gtc ttc c

NT

1D8 VH (nucleotide sequence) (SEQ ID NO:100)

20 caggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacagaccctgtccctcacatgcactgtctctgggtt
ctcattaaccagcgatgggtgtacactggattcgacagcctccaggaaagggtctggaatggatgggaataatatattatgat
ggaggcacagattataattcagcaattaaatccagactgagcatcagcagggacacctccaagagccaagtttcttaaaaa
tcaacagtctgcaaactgatgacacagccatgtattactgtgccagaatccactttgattactggggccaaggagtcatggtc
acagtctcctct

25 AA

1D8 VH (no leader) (amino acid sequence) (SEQ ID NO:101)

qvqlkeagpglvqptqtlsltctvsgfsltsdgvhwirpppgkglewmgiiyydgtdynsaiksrlsisrdtsksqvflk
inslqtdtdamyycarihfdywgqgvmvtvss

NT

1D8 VL (no leader) (nucleotide sequence) (SEQ ID NO:102)

gacattgtgctcactcagtcctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctcc
5 agtgaagttacatgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttc
tggagttccaaatcgcttcagtggtcagtggtctgggacctcttattctctcgcaatcaacaccatggagactgaagatgctg
ccacttattactgtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgg

AA

10 **1D8 VL (amino acid sequence) (SEQ ID NO:103)**

divltqspttiaaspgekvitcrasssvsymywyqqksgasplwiydtsklasgvpnrfsgsgsgtsyslaintmete
daatyycqqwsstpltfsgskleikr

NT

15 **1D8 scFv (nucleotide sequence) (SEQ ID NO: 104)**

aagcttatggattttcaagtcagattttcagcttctgtaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagtcctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctccagtgaagtta
catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
atcgcttcagtggtcagtggtctgggacctcttattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
20 gtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcgggtggtcggg
cgggtgggtgggtcgggtggcggcgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
ccctgtccctcatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagttttcttaaaaatcaacagtcctgcaaactgatgacacagccatgtattactgtgccagaatccac
25 ttgattactggggccaaggagtcaggtcacagtcctctgatca

AA

1D8 scFv (amino acid sequence) (SEQ ID NO: 105)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPTTIAASPGKEKVTITCRASSSVSYM
YWYQQKSGASPKLWIYDTSKLGVPNRFSGSGSGTSYSLAINTMETEDAAT
5 YYCQQWSSTPLTFGSGTKLEIKRGGGGSGGGGSGGGGSQVQLKEAGPGLVQP
TQTLSTCTVSGFSLTSDGVHWIRQPPGKGLEWMGIYYDGGTDYNSAIKSRL
SISRDTSKSQVFLKINSLQTDDTAMYYCARIHFDYWGQGVMVTVSS

NT

10 **1D8 scFv IgG WTH WTCH2CH3 (nucleotide sequence)** (SEQ ID NO: 106)

aagcttatggattttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaaccatagctgcatctccaggggagaaggcaccatcacctgccgtgccagctccagtgtgaagtta
catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagttcaa
atcgcttcagtggtcagtggttctgggacctcttattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
15 gtcagcagtggtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacggggtggcgggtggctcggg
cggtgggtgggtcgggtggcgccgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
ccctgtccctcacatgcactgtctctgggttctcattaaccagcagtggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagttttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtgccagaatccac
20 ttgattactggggccaaggagtcaggtcacagtctcctctgatcaggagcccaaattctgtgacaaaactcacacatgcc
accgtgccagcacctgaactcctggggggaccgtcagttctcttccccccaaaacccaaggacacctcatgatctc
ccggacccctgaggtcacatgcgtggtggtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggac
ggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctc
accgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcg
25 agaaaacaatctccaaagccaaaggcagccccgagaaccacaggtgtacacctgcccccatcccgggatgagctga
ccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgg
gcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgt
ggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcag
aagagcctctccctgtctccgggtaaatgatctaga

30

AA

1D8 scFv IgG WTH WTCH2CH3 (amino acid sequence) (SEQ ID NO:107)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPTTIAASPGEKVTITCRASSSVSYM
YWYQQKSGASPKLWIYDTSKLASGVPNRFSGSGSGTSYSLAINTMETEDAAT
5 YYCQQWSSTPLTFGSGTKLEIKRGGGGSGGGGSGGGGSQVQLKEAGPGLVQP
TQTLSLTCTVSGFSLTSDGVHWIRQPPGKGLEWMGIYYDGGTDYNSAIKSRL
SISRDTSKSQVFLKINSLQTDDTAMYYCARIHFDYWGQGVMVTVSSDQEPKS
CDKTHTCPPELPGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKV
10 SNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI
AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVM
HEALHNHYTQKSLSLSPGK

NT

1D8 scFv IgG MTH MTCH2CH3-CD80 (nucleotide sequence) (SEQ ID NO:108)

aagcttatggatttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctccagtgtgaagta
catgtactggtagcagaagtcaggcgccctccctaaactctggatttatgacacatccaagctggcttctggagttccaa
atcgcttcagtggcagtggtctgggacctcttattctctcgaatcaacaccatggagactgaagatgctgccacttattact
20 gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcgggtggctcggg
cggtgggtgggtcgggtggcgccgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
ccctgtccctcatatgcactgtctctgggttctcattaaccagcgtggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagtttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtccagaatccac
25 ttgattactggggccaaggagtcaggtcacagtctcctctgatctggagcccaaatctctgacaaaactcacacaagccc
accgagcccagcacctgaactcctggggggatcgtcagttctcttccccccaaaacccaaggacacctcatgatctc
ccggacccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggac
ggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctc
accgtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaacaaagccctcccagcccccatcg
30 agaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctga
ccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatgg
gcagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgt

ggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcag
aagagcctctccctgtctccgggtaaacggatccttcgaacctgctcccatcctgggccattacctaatactcagtaaattg
aatTTTTgtgatatgctgcctgacctactgctttgccccaaagatgcagagagagaaggaggaatgagagattgagaaggga
aagltacgccctgtataaatcgata

5

AA

1D8 scFv IgG MTH MTCH2CH3-CD80 (amino acid sequence) (SEQ ID NO:109)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPTTIAASPGEKVTITCRASSSVSYM
YWYQQKSGASPKLWIYDTSKLASGVPNRFSGSGSGTSYSLAINTMETEDAAT
10 YYCQQWSSTPLTFGSGTKLEIKRGGGSGGGGSGGGGSQVQLKEAGPGLVQP
TQTLSLTCTVSGFSLTSDGVHWIRQPPGKGLEWMGIYYDGGTDYNSAIKSRL
SISRDTSKSQVFLKINSLQTDDTAMYYCARIHFDYWGQGMVTVSSDLEPKSS
DKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVK
FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKV
15 NKALPAIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIA
VEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMH
EALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVNGIFVICCLTYCFAPRCR
RRRNERLRRESVRPV

20 NT

1D8 scFv IgG WTH WTCH2CH3-CD80 (nucleotide sequence) (SEQ ID NO: 110)

aagcttatggatttcaagtcagatttccagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaaccatagctgcatctccaggggagaagggtccatcacctgccgtgccagctccagtgaagtta
catgtactggtaccagcagaagtcaggcgccctcccctaaactctggatttatgacacatccaagctggcttctggagtccaa
25 atcgcttcagtggcagtggtctgggacctctattctctcgcaatcaacacatggagactgaagatgctgccacttattact
gtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
cggtgggtgggtcgggtggcgccgatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
ccctgtccctcacatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
30 acacctccaagagccaagtttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtgccagaatccac
tttgattactggggccaaggagtcaggtcacagctctcctctgatctggagcccaaatcttgacaaaactcacacatgcc

accgtgccagcacctgaactcctggggggaccgtcagtcctccttccccccaaaaccaaggacacccctcatgatctc
 ccggacccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggac
 ggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctc
 accgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcg
 5 agaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctga
 ccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgg
 gcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcacctg
 ggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcag
 aagagcctctccctgtctccgggtaaagcggatccttcgaacctgctcccatcctgggccattacctaatactcagtaaattg
 10 aattttgtgatatgctgcctgacactactgctttgccccaaagatgcagagagagaaggaggaatgagagattgagaaggga
 aagtgtacgccctgtataaatcgata

AA

1D8 scFv IgG WTH WTCH2CH3-CD80 (amino acid sequence) (SEQ ID NO:

15 111)
 MDFQVQIFSLLISASVIMSRGVDIVLTQSPTTIAASPGEKVTITCRASSSVSYM
 YWYQQKSGASPKLWIYDTSKLGSGVGNRFSGSGSGTSYSLAINTMETEDAAT
 YYCQQWSSTPLTFGSGTKLEIKRGGGSGGGGSGGGGSQVQLKEAGPGLVQP
 TQTLSTCTVSGFSLTSDGVHWIRQPPGKGLEWMGIYYDGGTDYNSAIKSRL
 20 SISRDTSKSQVFLKINSLQTDAMYYCARIHFDYWGQGMVTVSSDLEPKS
 CDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
 KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKV
 SNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI
 AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSFSVM
 25 HEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVNGIFVICCLTYCFAPRCR
 ERRRNERLRRESVRPV

NT

Anti human CD3 scFv WTH WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 112)

aagcttatggattttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacatccaga
5 tgacacagactacatcctccctgtctgcctctctgggagacagagtcacccatcagttgcagggcaagtcaggacattcgca
attatttaaactggatcagcagaaaccagatggaactgttaaactcctgatctactacacatcaagattacactcaggagtcc
catcaaggttcagtgccagtggtctggaacagattattctctcaccattgccaacctgcaaccagaagatattgccacttact
tttgccaacagggttaatacgttccgtggacgttcggtggaggcaccaaactggtaaccaaaggagctcggtggcggg
ggctcgggcgggtgggtgggtggcggcggtatcgcgatgaggtccagctgcaacagctctggacctgaactgggtga
10 agcctggagcttcaatgtcctgcaaggcctctggttactcattcactggctacatcgtgaactggctgaagcagagccatgg
aaagaaccttgagtggattggacttattaatccatacaaaggtcttactacctacaaccagaaattcaagggaaggccacat
taactgtagacaagtcacccagcacagcctacatggagctcctcagctctgacatctgaagactctgcagctcttactgtgca
agatctgggtactatggtgactcggactggacttcgatgtctggggcgagggaccacggtcaccgtctcctctgatcagg
agcccaaacttctgtacaaaactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttctcctctt
15 ccccccaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtggtgggtggacgtgagccacgaa
gacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcag
tacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtcaa
ggtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggt
gtacaccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatecca
20 gcgacatcgccgtggagtgggagagcaatgggcagccgggagaaactacaagaccacgcctcccgtgctggactccg
acggctccttctcctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaactcttctcatgctccgtga
tgcatagaggtctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaatgatctaga

AA

25 **Anti human CD3 scFv WTH WTCH2CH3 (amino acid sequence)** (SEQ ID NO: 113)

MDFQVQIFSLLISASVIMSRGVDIQMTQTTSSLSASLGDRVITISCRASQDIRNY
LNWYQQKPDGTVKLLIYYTSRLHSGVPSRFSGSGSGTDYSLTIANLQPEDIAT
YFCQQGNTLPWTFGGGKLVTKRELGGGGSGGGGSGGGGSIDEVQLQQSGP
30 ELVKPGASMSCKASGYSFTGYIVNWLKQSHGKNLEWIGLINPYKGLTTYNQK
FKGKATLTVDKSSSTAYMELLSLTSEDSAVYYCARSGYYGDSWDYFDVWGA
GTTVTVSSDQEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV

TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL
HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDK
SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

5

NT

2H7-antiCD40 scFv MTH (SSS) MTCH2WTCH3 (2h7-40.2.220Ig + restriction sites) (nucleotide sequence) (SEQ ID NO: 114)

aagcttgccgcatggattttcaagtcagattttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
10 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttcagggccagctcaagttaag
ttacatgactggtagcagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgcttcggtgctgggaccaagctggagctgaaaggtggcgggtggctcg
ggcgggtggtagctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
15 cctcagtgaaagatgtcctgcaaggcttctggctacacattaccaggttacaatatgactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtggtgtactatagtaacttacttggtacttcagtgctctggggcacagggaccacggtcaccgtctcttctgataatcca
actctgaagaagcaaaagaaagaggaggccaaaaaggaggaagccaagaaatctaacagcgtcgacattgttctgactca
20 gtctccagccaccctgtctgtgactccaggagatagagtctctcttctcctgcagggccagccagagtattagcgactacttac
actggtatcaacaaaaatcacatgagctctcaaggcttctcatcaaatatgcttccattccatctctgggatccccctcagggtt
cagtggcagtggtacagggtcagatttctcagtatcaacagtggtgaacctgaagatgttggaatttattactgtcaaca
tggtcacagctttccgtggacgttcggtggaggcaccaagctggaaatcaaacggggtggcgggtggctcgggcggaggt
gggtcgggtggcggcggtatctcagatccagttggtgcaatctggacctgagctgaagaagcctggagagacagtcagga
25 tctcctgcaaggcttctgggtatgccttcacaactactggaatgcagtggtgcaagagatgccaggaaagggttgaaagt
gattggctggataaacaccccactctggagtgccaaaatatgtagaagacttcaaggacggtttgccttctctttgaaacct
ctgccaaactgcatatttacagataagcaacctcaaagatgaggacacggctacgtatttctgtgtgagatccgggaatgg
taactatgacctggcctactttgcttactggggccaagggacactggtcactgtctctgatcaggagcccaaatcttctgaca
aaactcacacatccccaccgtccccagcacctgaactcctggggggatcgtcagttcttcttccccccaaaaccaagg
30 acaccctcatgatctcccgaccctgaggtcacatgcgtggtggtagctgagccacgaagaccctgaggtcaagttc
aactggtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgt
gtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccct

ccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatc
ccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtaaaggcttctatcccagcgacatcgccgtggagt
gggagagcaatgggcagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctac
agcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaa
5 ccactacacgcagaagagcctctccctgtctccgggtaaataatgatctaga

AA

2H7-antiCD40 scFv MTH (SSS) MTCH2WTCH3 (2H7-40.2.220Ig) (amino acid sequence) (SEQ ID NO: 115)

10 MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPFTFGAGTKLELKGSGSGSGSGSGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
15 VTVSSDQSNSEEAKKEEAKKEEAKKSNSVDIVLTQSPATLSVTPGDRVSLSCR
ASQSIDYLVHWYQQKSHESPRLLIKYASHSISGIPSRFSGSGSGSDFTLSINSVEP
EDVGIYYCQHGHSFPWTFGGGTKLEIKRGGGGSGGGGSGGGGSGQIQLVQSGP
ELKKPGETVRISCKASGYAFTTTGMQWVQEMPGKGLKWIGWINTPLWSAKI
CRRLQGRFAFSLETSANTAYLQISNLKDEDTATYFCVRSGNGNYDLAYFAYW
20 GQGTLVTVSDQEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPE
VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVD
KSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

25

NT

5B9 VH (includes the VH leader peptide) (nucleotide sequence) (SEQ ID NO: 116)

atggctgtcttggggctgctcttgcctggtagacattccaagctgtgcctatcccaggtgcagctgaagcagtcaggacc
30 tggcctagtgcagtcctcacagagcctgccatcacctgcacagtctctggtttctcattaactacctatgctgtacactgggtt
cgccagctccaggaaaggtctggagtggctgggagtgatatggagtggggaatcacagactataatgcagctttcatat

ccagactgagcatcaccaaggacgattccaagagccaagttttctttaaataaacagctgcaacctaataacacagccat
ttattactgtgccagaaatgggggtgataactacccttattactatgctatggactactgggggtcaaggaacctcagtcaccgt
ctcctca

5 NT

5B9 VH (missing the leader) (nucleotide sequence) (SEQ ID NO: 117):

caggtgcagctgaagcagtcaggacctggcctagtgcagtcctcacagagcctgtccatcacctgcacagtctctggttct
cattaactacctatgctgtacactgggttcgccagtctccaggaaagggctctggagtggctgggagtgatatggagtgggtg
aatcacagactataatgcagctttcatatccagactgagcatcaccaaggacgattccaagagccaagttttctttaaataa
10 cagctgcaacctaataacacagccatttattactgtgccagaaatgggggtgataactacccttattactatgctatggacta
ctgggggtcaaggaacctcagtcaccgtctcctca

AA

5B9 VH (includes leader peptide) (amino acid sequence) (SEQ ID NO: 118)

15 MAVLGLLFCLVTFPSCVLSQVQLKQSGPGLVQSSQSL SITCTVSGFSLTTYAV
HWVRQSPGKGLEWLGVWSSGITDYNAAFISRLSITKDDSKSQVFFKMNSLQ
PNDTAIYYCARNGGDNYPYYAMDYWGQGTSVTVSS

5B9 VH (no leader peptide) (amino acid sequence) (SEQ ID NO: 119)

20 QVQLKQSGPGLVQSSQSL SITCTVSGFSLTTYAVHWVRQSPGKGLEWLGVW
SSGITDYNAAFISRLSITKDDSKSQVFFKMNSLQ PNDTAIYYCARNGGDNYPY
YYAMDYWGQGTSVTVSS

NT

25 **5B9 VL (nucleotide sequence) (SEQ ID NO: 120)**

atgaggttctctgctcagcttctggggctgcttgctctggatccctggatccactgcagatattgtgatgacgcaggctgca
ttctccaatccagtcactcttggacatcagcttccatctcctgcaggcttagtaagagtctcctacatagtaatggcatcactta
tttgattggatctgcagaagccaggccagctcctcagctcctgatttatcagatgtccaaccttgccctcaggagtcaccaga

cagggtcagtagcagtggggcaggaactgattcacactgagaatcagcagagtgagggtgaggatgtgggtgtttattac
tgtgctcaaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacgg

AA

5 **5B9 VL (amino acid sequence)** (SEQ ID NO: 121)

MRFSAQLLGLLVLWIPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSHNG
ITYLYWYLQKPGQSPQLLIYQMSNLAGVVPDRFSSSGSGTDFTLRISRVEAED
VGVYYCAQNLELPLTFGAGTKLELKR

10 NT

5B9 scFv (nucleotide sequence) (SEQ ID NO: 122)

aagcttgccgccatgaggttctctgctcagcttctggggctgcttgtgctctggatccctggatccactgcagatattgtgatg
acgcaggtgcattctccaatccagtcactcttgaacatcagcttccatctctgcaggtctagtaagagtctcctacatagt
aatggcatcacttatttatttggtatctgcagaagccaggccagtcctcctcagctcctgatttatcagatgtccaacctgcctc
15 aggagtcacagacaggttcagtagcagtgggtcaggaactgattcacactgagaatcagcagagtgagggtgaggatg
tggtgtttattactgtgctcaaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacggggtgg
cgggtggctcgggcgggtgggtgggtcgggtggcggcgatcgtcacaggtgcagctgaagcagtcaggacctggcctagt
gcagtcctcacagacgtgtccatcacctgcacagtcctctggtttctcattaactacctatgtgtacactgggttcgccagtct
ccaggaaagggtctggagtgggtgggagtgatatggagtgggtggaatcacagactataatgcagctttcatatccagactg
20 agcatcaccaaggacgattccaagagccaagttttctttaaataaacagctctgcaacctaatacacagccatttattactgt
gccagaaatgggggtgataactacccttattactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctct

AA

5B9 scFv (amino acid sequence) (SEQ ID NO: 123)

25 MRFSAQLLGLLVLWIPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSHNG
ITYLYWYLQKPGQSPQLLIYQMSNLAGVVPDRFSSSGSGTDFTLRISRVEAED
VGVYYCAQNLELPLTFGAGTKLELKRGGGGSGGGGSGGGGSSQVQLKQSGP
GLVQSSQSLSITCTVSGFSLTTYAVHWVRQSPGKLEWLGVWISGGITDYNA
AFISRLSITKDDSKSQVFFKMNSLQPNDAIYYCARNGGDNYPYYYAMDYW
30 GQGTSVTVSS

NT

5B9 scFv-hmtIgG1-hCD80 (nucleotide sequence) (SEQ ID NO: 124)

aagcttgccgcatgaggttctctgctcagcttctggggctgcttgtgctctggatccctggatccactgcagatattgtgatg
acgcaggctgcatttccaatccagtcactcttggacatcagcttccatctcctgcaggtctagtaagagtctcctacatagt
5 aatggcatcacttatttgtattggtatctgcagaagccaggccagctcctcagctcctgatttatcagatgtccaacctgcctc
aggagtcccagacaggttcagtagcagtggtcaggaactgatttcacactgagaatcagcagagtggaggctgaggatg
tggtgttttattactgtgctcaaaatctagaacttccgctcacttccggtgctgggaccaagctggagctgaaacggggtgg
cgggtggctcgggcggtggtgggtcgggtggcgccggtatgacacaggtgcagctgaagcagtcaggacctggcctagt
gcagtcctcacagagcctgtccatcacctgcacagctcttggtttctattaactacctatgctgtacactgggttcgccagtct
10 ccaggaaagggctgagtggtgggtggtggtgatatggagtgggtggaatcacagactataatgcagctttcatatccagactg
agcatcaccaaggacgattccaagagccaagttttcttaaaatgaacagtctgcaacctaatacacagccatttattactgt
gccagaaatgggggtgataactacccttattactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctctga
tctggagcccaaacttctgacaaaactcacacaagcccaccgagcccagcacctgaactcctggggggatcgtcagctctt
cctcttccccccaaaacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgagcc
15 acgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggagg
agcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaa
gtgcaaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaacc
acaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttct
atcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctgg
20 actccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgct
ccgtgatgatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaacgggatccttcgaacc
tgctcccatcctgggccattaccttaatctcagtaaatggaattttgtgatatgctgcctgacctactgctttgccccaaagatgc
agagagagaaggaggaatgagagattgagaagggaaagtgtacgccctgtataaatcgatactcgag

25 AA

5B9 scFv-hmtIgG1-hCD80 (amino acid sequence) (SEQ ID NO: 125)

MRFSAQLLGLLVLWIPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSNG
ITYLYWYLQKPGQSPQLLIYQMSNLSGVLPDRFSSSGSGTDFTLRISRVEAED
VGVIYCAQNLELPLTFGAGTKLELKRGGGSGGGGSGGGGSSQVQLKQSGP
30 GLVQSSQSLSITCTVSGFSLTTYAVHWVRQSPGKGLEWLGVIWSGGITDYNA
AFISRLSITKDDSKSQVFFKMNSLQPNDAIYYCARNGGDNYPIYYAMDYW
GQGTSVTVSSDLEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPE

VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVD
KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISVN
5 GIFVICCLTYCFAPRCRERRRNERLRRESVRPV

NT

2e12 scFv WTH CH2 CH3 (2e12 scFv-WthIgG-CD80) (nucleotide sequence)
(SEQ ID NO: 126)

10 aagcttatggatttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttcttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaagtggtgaatatta
tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
tctgggggtccctgccaggttagtggcagtggtgtgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaagtaggaaggttccttgacgttcggtggaggcaccaagctggaaatcaaacggggtgg
15 cggtggctcggggcggaggtgggtggcgggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcct
ccaggaaagggtctggagtgggtgggaatgatatgggggtgatggaagcacagactataattcagctctcaatccagactg
agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgcaaactgatgacacagccagataactact
gtgccagagatggttatagtaactttcattactatgttatggactactgggggtcaaggaacctcagtcaccgtctcctcagatct
20 ggagcccaaactctgtgacaaaactcacacatgccaccgtgccagcacctgaactcctgggggggaccgtcagctttcct
cttccccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacg
aagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagc
agtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgc
aaggctctcaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacag
25 gtgtacaccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccc
agcgacatcgccgtggagtgggagagcaatgggcagccgggagaacaactacaagaccacgcctcccgtgctggactcc
gacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtg
atgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaagcggatccttgaacctgctc
ccatcctgggccattacctaatactcagtaaatggaattttgtgatatgctgcctgacctactgctttgccccaaagtgcagag
30 agagaaggagggaatgagagattgagaagggaaggtgtacgccctgtataaatcgat

AA

2e12 scFv WTH CH2 CH3 (2e12 scFv-WthIgG-CD80) (amino acid sequence)

(SEQ ID NO: 127)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPASLAVSLGQRATISCRASESVEYY
5 VTSLMQWYQQKPGQPPKLLISAASNVESGVPARFSGSGSGTDFSLNIHPVEED
DIAMYFCQQSRKVPWTFGGGTKLEIKRGGGSGGGGSGGGGSQVQLKESGP
GLVAPSQSLITCTVSGFSLTGYGVNWVRQPPGKLEWLGMWGDGSTDYNS
ALKSRLSITKDNSKSQVFLKMNSLQTDRTARYYCARGYSNFIHYVMDYW
GQGTSVTVSSDLEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMISRT
10 EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT
VLHQQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL
KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTV
DKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKADPSNLLPSWAITLISV
NGIFVICCLTYCFAPRCRERRRNERLRRESVRPV

15

NT

2H7-human IgE Fc (CH2-CH3-CH4) (nucleotide sequence) (SEQ ID NO: 128)

aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatctgtctgcattctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
20 ttacatgcactggtaccagcagaagccaggatcctcccccaccctggattatgccccatccaacctggcttctggagtc
cctgctcgcttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggagctgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagtgccggtggtcgc
ggcggtggtgagcttgaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
25 gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaacttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctgatcacgtctgct
ccagggacttcaccccgccaccgtgaagatcttacagtcgtctgacgacggcgggcacttcccccgaccatcca
gctcctgtgcctcgtctctgggtacacccaggactatcaacatcacctggctggaggacgggcaggtcatggacgtgg
30 actgtccaccgcctctaccacgcaggaggggtgagctggcctccacacaaagcgagctcacctcagccagaagcactg
gctgtcagaccgcacctacacctgccagggtcacctatcaagggtcacaccttgaggacagcaccaagaagtgtgcagattc
caaccgagaggggtgagcgcctacctaagccggcccgagccggttcgacctgttcacccgaagtcgcccacgatcacc

tgtctggtggtggacctggcaccagcaaggggaccgtgaacctgacctgggtcccgggccagtgggaagcctgtgaacc
actccaccagaaaggaggagaagcagcgcaatggcacgttaaccgtcacgtccaccctgccgggtgggcacccgagact
ggatcgagggggagacctaccagtgcagggtgacccacccccacctgccagggccctcatgcggtccacgaccaag
accagcggcccgcgltgctgccccggaagtctatgcgtttgcgacgccggagtggccggggagccgggacaagcgcac
5 cctcgctgcctgatccagaacttcacgtgcctgaggacatctcggtgcagtggctgcacaacgaggtgcagctcccgacg
cccggcacagcagcagcagccccgcaagaccaagggtccggcttctctcttcagccgcctggaggtgaccaggg
ccgaatgggagcagaaagatgagttcatctgccgtgcagtcctatagggcagcagccccctcacagaccgtccagcgagc
ggtgtctgtaaattcccggtaaatgataatctaga

10 AA

2H7 scFv IgE (CH2-CH3-CH4) (amino acid sequence) (SEQ ID NO: 129)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPFTFGAGTKLELKGGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
15 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSDHVCSDFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTINITWLE
DGQVMDVDLSTASTTQEGELASTQSELTLSQKHWLSDRTYTCQVTYQGHTF
EDSTKKCADSNPRGVSAÿLSRPSPFDLFIRKSPTITCLVVDLAPSKGTVNLTWS
20 RASGKPVNHSTRKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTHPHLPR
ALMRSTTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDISVQWL
HNEVQLPDARHSTTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAVHEAA
SPSQTVQRAVSVNPGK

25 NT

2H7 scFv MH (SSS) MCH2WTCH3 (nucleotide sequence) (SEQ ID NO: 130)

aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatacgtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaagggtcacaatgacttgaggggccagctcaagtgaag
ttacatgcactggtagcagagaagccaggatcctcccccacacctggatttatgccccatccaacctggcttctggagtc
30 cctgctcgcttcagtggcagtggtctgggacctcttactctctcacaatcagcagagtgaggagctgaagatgctgccactt
attactgccagcagtgaggatttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg

5 ggcggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
 cctcagtgaaagatgtcctgcaaggcttctggctacacattaccaggtacaatatgcactgggtaaagcagacacctagaca
 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
 actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 10 gagtgggtgactatagtaactcttactgggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcaggag
 cccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggatcgtcagttctcttctccc
 cccaaaaccaagacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaagac
 cctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgaggagagcagtag
 aacagcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactgggtgaatggcaaggagtacaagtgaaggt
 15 ctccaacaaagccctccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgccccatcccggtgatgagctgaccaagaaccaggtcagcctgacctgctgtgtaaaggcttctatccagcg
 acatcgccgtggagtgaggagagcaatgggcagccggagaacaactacaagaccacgcctcccggtgtggactccgacg
 gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

15

AA

2H7 scFv MH (SSS) MCH2WTCH3 (amino acid sequence) (SEQ ID NO: 131)

20 MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
 CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
 VTVSSDQEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVV
 VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
 25 LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLT
 CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
 QGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

5B9 scFv MTHWTCH2CH3 (nucleotide sequence) (SEQ ID NO: 132)

aagcttgccgccaatgaggttctctgctcagcttctggggctgcttgctctggatccctggatccactgcagatattgtgatg
acgcaggctgcattctccaatccagtcactcttggacatcagcttccatctcctgcaggctctagtaagagtctctacatagt
5 aatggcatcacttatttatttggtatctgcagaagccaggccagctctcctcagctcctgatttatcagatgtccaaccttgctc
aggagtcccagacaggttcagtagcagtggtcaggaactgatttcacactgagaatcagcagagtggaggctgaggatg
tgggtgtttattactgtgctcaaaatctagaacttccgctcacgttcggtgctgggaccaagctggagctgaaacggggtgg
cggtggtcgggcggtggtgggtcgggtggcggcggatcgtcacaggtgcagctgaagcagtcaggacctggcctagt
gcagtcctcacagagcctgtccatcacctgcacagctcttggtttctattaactacctatgctgtacactgggttcgccagtct
10 ccaggaaagggctcggagtgggtgggtgatatggagtgggtggaatcacagactataatgcagctttcatatccagactg
agcatcaccaaggacgattccaagagccaagttttcttaaaatgaacagctctgcaacctaatacagacagccatttactgt
gccagaaatgggggtgataactacccttattactatgctatggactactggggtcaaggaacctcagtcaccgtctcctctga
tcaggagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagctctt
cctctccccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagcc
15 acgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggagg
agcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaa
gtgcaaggtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaacc
acaggtgtacacctgccccatcccggtgatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttct
atcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctgg
20 actccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgct
ccgtgatgatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

5B9 scFv MTHWTCH2CH3 (amino acid sequence) (SEQ ID NO: 133)

25 MRFSAQLLGLLVLPWPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSNG
ITYLYWYLQKPGQSPQLLIYQMSNLAGVFPDRFSSSGSGTDFTLRISRVEAED
VGVYYCAQNLELPLTFGAGTKLELKRGGGSGGGGSGGGGSSQVQLKQSGP
GLVQSSQSLSITCTVSGFSLTTYAVHWVRQSPGKGLEWLGVIWSGGITDYNA
AFISRLSITKDDSKSQVFFKMNSLQPNDTAIYYCARNGGDNYPIYYAMDYW
30 GQGTSVTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPE
VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK

NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD
KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Human IgG1 hinge mutations

5 NT

2H7 scFv- MTH (CSS) WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 134)

Nucleotide:

aagcttgccgcatggattttcaagtgcagattttcagcttcctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcaaggccagctcaagtgaag
10 ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgcttcagtggcagtggtctgggacctcttactctctcacaatcagcagagtgaggagctgaagatgctgccactt
attactgccagcagtgaggttttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg
ggcgggtgggtgagctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
15 gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggctctattctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaattctgtgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagttctcttcc
ccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacgaaga
20 ccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaaggcttctatcccagc
gacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
25 ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

2H7 scFv- MTH (CSS) WTCH2CH3 (amino acid sequence) (SEQ ID NO: 135):

30 MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYM
H
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY

CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSSDQEPKSCDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV
5 VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
WLNQKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

10 NT

2H7 scFv- MTH (SCS) WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 136):

aagcttgccgccatggattttcaagtcagatgttcagcttctgctaatacagtcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctcccccaccctggattatgccccatccaacctggcttctggagtc
15 cctgtcgttcagtggcagtggtctgggaccttactcttcacaatcagcagagtgaggctgaagatgtgccactt
attactgccagcagtgagggttttaaccccccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctc
ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactg
20 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcgggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcaggag
cccaaattcttgacaaaactcacacatgccaccgtccccagcacctgaactcctggggggaccgtcagcttctcttcc
cccaaaaacccaaggacaccctcatgatctccggaccctgaggtcacatgcgtggtgggtggacgtgagccacgaaga
ccctgagggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
25 caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagcccccatcgaaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acaccctgcccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctatcccagc
gacatcgccgtggagtgggagagcaatgggcagccggagacaactacaagaccacgcctcccgtgctggactccgac
ggctccttctctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
30 catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

2H7 scFv- MTH (SCS) WTCH2CH3 (amino acid sequence) (SEQ ID NO: 137):

NT

15 **2H7 scFv- MTH (SSC) WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 138):**

348

ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

5 **2H7 scFv- MTH (SSC) WTCH2CH3 (amino acid sequence)** (SEQ ID NO: 139):

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
10 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSSDQEPKSSDKTHTSPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRW
15 QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

HIgGMHcys1 (nucleotide sequence) (SEQ ID NO: 140)

ggt ggt gat cag gag ccc aaa tct tct gac aaa act cac aca tg

20

NT

HIgGMHcys2 (nucleotide sequence) (SEQ ID NO: 141)

ggt ggt gat cag gag ccc aaa tct tgt gac aaa act cac aca tct cca ccg tgc

25 NT

HIgGMHcys3 (nucleotide sequence) (SEQ ID NO: 142)

ggt ggt gat cag gag ccc aaa tct tgt gac aaa act cac aca tgt cca ccg tcc cca gca cct

NT

HuIgG1 MTCH3Y405 (nucleotide sequence) (SEQ ID NO: 143)

gggcagccccgagaaccacaggtgtacaccctgccccatcccgggaggagatgaccaagaaccaggtcagcctgac
ctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaag
5 accacgcctcccgtgctggactccgacggctccttctacctctatagcaagctcaccgtggacaagagcaggtggcagca
ggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccggg
taaatga

AA

10 **HuIgG1 MTCH3Y405** (amino acid sequence) (SEQ ID NO: 144)

GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFYLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS
PGK

15 NT

HuIgG1 MTCH3A405 (nucleotide sequence) (SEQ ID NO: 145)

gggcagccccgagaaccacaggtgtacaccctgccccatcccgggaggagatgaccaagaaccaggtcagcctgac
ctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaag
accacgcctcccgtgctggactccgacggctccttcgccctctatagcaagctcaccgtggacaagagcaggtggcagca
20 ggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccggg
taaatga

AA

HuIgG1 MTCH3A405 (amino acid sequence) (SEQ ID NO: 146)

25 GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFALYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS
PGK

NT

HuIgG1 MTCH3A407 (nucleotide sequence) (SEQ ID NO: 147)

Gggcagccccgagaaccacaggtgtacaccctgccccatcccgggaggagatgaccaagaaccaggtcagcctgac
ctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagagcaatgggcagccggagaacaactacaag
5 accacgcctcccgtgctggactccgacggctccttctcctcgccagcaagctcaccgtggacaagagcaggtggcagca
ggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccggg
taaatga

AA

10 **HuIgG1 MTCH3A407 (amino acid sequence)** (SEQ ID NO: 148)

GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLASKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS
PGK

15 NT

HuIgG1 MTCH3Y405A407 (nucleotide sequence) (SEQ ID NO: 149)

gggcagccccgagaaccacaggtgtacaccctgccccatcccgggaggagatgaccaagaaccaggtcagcctgac
ctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagagcaatgggcagccggagaacaactacaag
accacgcctcccgtgctggactccgacggctccttctacctcgccagcaagctcaccgtggacaagagcaggtggcagc
20 aggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgg
gtaaatga

AA

HuIgG1 MTCH3Y405A407 (amino acid sequence) (SEQ ID NO: 150)

25 GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFYLASKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS
PGK

NT

HuIgG1 MTCH3A405A407 (nucleotide sequence) (SEQ ID NO: 151)

gggcagccccgagaaccacaggtgtacacctgccccatccccgggaggagatgaccaagaaccaggtcagcctgac
ctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaag
5 accacgcctcccgtgctggactccgacggctccttcgccctgccagcaagctcaccgtggacaagagcaggtggcagc
aggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgg
gtaaatga

AA

10 **HuIgG1 MTCH3A405A407 (amino acid sequence)** (SEQ ID NO: 152)

GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFALASKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS
PGK

15 NT

2H7 scFv MTH (SSS) WTCH2MTCH3Y405 (nucleotide sequence) (SEQ ID NO: 153)

aagcttgccgcatggattttcaagtgagattttcagcttctgtaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagtgaag
20 ttacatgcactggtaccagcagaagccaggatcctccccaaacctggatttatgccccatccaacctggcttctggagtc
cctgtctgcttcagtgggcagtggtctgggaccttactctctcacaatcagcagagtgaggagctgaagatgctgccactt
attactgccagcagtgaggtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
ggcggtgggtgatctggaggaggtgggagctctcaggcttctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacatttaccagttacaatatgcactgggtaaagcagacacctagaca
25 gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagttctctcttcc
cccaaaaaccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtgggtggacgtgagccacgaaga
30 ccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag

gltccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
 acacctgcccccatcccgaggagatgaccaagaaccaggtcagcctgacctgcctggtaaaaggcttctatcccagc
 gacatgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
 ggctccttctaccttatagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
 5 catgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgggtaaatgatctaga

AA

2H7 scFv MTH (SSS) WTCH2MTCH3Y405 (amino acid sequence) (SEQ ID NO: 154)

10 MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
 CQQWSFNPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
 15 VTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV
 VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
 LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT
 CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFYLYSKLTVDKSRWQ
 QGNVFSCSVMHEALHNHYTQKSLSLSPGK

20

NT

2H7 scFv MTH (SSS) WTCH2MTCH3A405 (nucleotide sequence) (SEQ ID NO: 155)

aagcttgccgcatggatttcaagtcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
 25 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagtgaag
 ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
 cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
 attactgccagcagtgaggtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg
 ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
 30 cctcagtgaaatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg

actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggctctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagttcttcttcc
cccaaaaaccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtggacgtgagccacgaaga
5 ccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acaccctgcccccatcccgggaggagatgaccaagaaccaggtcagcctgacctgcctggtaaaaggcttctatcccagc
gacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
10 ggctccttcgccctctatagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctctcatgctccgtgatg
catgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgggtaaatga

AA

2H7 scFv MTH (SSS) WTCH2MTCH3A405 (nucleotide sequence) (SEQ ID NO:
15 156)

mdfqvqifsfllisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiapsnlasg
vparfsgsgsgtsysltisrveadaatyycqwsfnpptfgatgkklkdggsgggsgggssqaylqqsgaelvr
pgasvkmsckasgytftsynmhvwkqtrqglewigaiypngdtsynqkfkkgkatltvdksstaymqlssltسد
savyfcarvvyysnsywyfdvwgtgtvtvssdqepkssdkthtsppspapellggpsvflfppkpkdtlmsrtpevt
20 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
skakgqprepqvtylppsreemtknqvslclvkgfypsdiavewesngqpennykttpvldsdgsfalyskltvdk
srwqqgnvfscsvmhealhnhytqkslsispkg

NT

2H7 scFv MTH (SSS) WTCH2MTCH3A407 (nucleotide sequence) (SEQ ID NO:
25 157)

aagcttccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctccccaaacctggatttatgccccatccaacctggcttctggagtc
30 cctgctcgttcagtggcagtggtctgggaccttactctctacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcgggtgctgggaccaagctggagctgaaagatggcgggtggctcg

ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
 cctcagtgaagatgtcttgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactg
 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 5 gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
 cccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagtcttctcttcc
 ccccaaaaccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtgggtgacgtgagccacgaaga
 ccctgaggtcaagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
 caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
 10 gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtg
 acaccctgcccccatcccgggaggagatgaccaagaaccaggtcagcctgacctgcctggtaaaaggcttctatcccagc
 gacatcgccgtggagtgggagagcaatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgac
 ggctccttctctcgcagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
 catgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgggtaaatga

15

AA

2H7 scFv MTH (SSS) WTCH2MTCH3A407 (amino acid sequence) (SEQ ID NO: 158)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
 20 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
 CQQWSFNPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
 VTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV
 25 VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
 LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT
 CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLASKLTVDKSRWQ
 QGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

2H7 scFv MTH (SSS) WTCH2MTCH3Y405A407 (nucleotide sequence) (SEQ ID NO: 159)

aagcttgccgccaatggattttcaagtcagattttcagcttcctgctaatacagtgcttcagtcataattgccagaggacaaattg
5 ttctctcccagctcagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgaggtttaaccccccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
ggcggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagctggtgagcgctgggg
10 cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagtccaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaattctctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagttctcttcc
15 ccccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga
ccctgaggtcaagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acaccctgccccatcccgggaggagatgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagc
20 gacatcgccgtggagtgggagagcaatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgac
ggctccttctacctgccagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgat
gcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgggtaaatga

AA

25 **2H7 scFv MTH (SSS) WTCH2MTCH3Y405A407 (amino acid sequence)** (SEQ ID NO: 160)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGSSQAYLQQSGAELVRPG
30 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
VTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV

VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFIYPSDIAVEWESNGQPENNYKTTTPVLDSGFSYFYLASKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK

5

NT

2H7 scFv MTH (SSS) WTCH2MTCH3A405A407 (nucleotide sequence) (SEQ ID NO: 161)

aagcttgccgcatggattttcaagtcagattttcagcttcctgctaatactgcttcagtcataattgccagaggacaaattg
10 ttctctcccagctccagcaatcctgtctgcattctccaggggagaaggtcacatgacttcagggccagctcaagtgaag
ttacatgcactggtagcagcagaagccaggatcctccccaaacctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggacctcttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgaggtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
ggcggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
15 cctcagtgaaagatgctcgaaggcttctggctacacatttaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagtcaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtggtgtactatagtaactttacttggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagttcttcttcc
20 ccccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga
ccctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acaccctgccccatcccgggaggagatgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagc
25 gacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
ggctccttcgccctcgccgaagctcaccgtggacaagagcagtggtgcagcaggggaacgtcttctcatgctccgtgat
gcatgaggctctgcacaaccactacacgcagaagagcctctccctgtccccgggtaaatga

AA

2H7 scFv MTH (SSS) WTCH2MTCH3A405A407 (amino acid sequence) (SEQ ID NO: 162)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
5 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV
10 VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFALASKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK

15 NT

2H7 scFv MTH (SCC) WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 163)

aagcttgccgcatggatttcaagtcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctccccaaacctggatttatgccccatccaacctggcttctggagtc
20 cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtgctc
ggcgggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacatttaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
25 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaacttacttggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcaggag
cccaaattcttgacaaaactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttctctctcc
ccccaaaacccaaggacacctcatgatctccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga
ccctgaggtcaagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
30 caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaaagccctcccagccccatcgaaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctatcccagc

gacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

5 AA

2H7 scFv MTH (SCC) WTCH2CH3 (amino acid sequence) (SEQ ID NO: 164)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYM
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGSSQAYLQQSGAELVRPG
10 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSSDQEPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
WLNQKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSL
15 TCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

2H7 scFv MTH (CSC) WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 165)

20 aagcttgccgccatggatttcaagtgcagattttagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattctccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctccccaaaccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactcttcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgaggattttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggaggctcg
25 ggcgggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
30 cccaaatcttgtgacaaaactcacacatctccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcc
ccccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga

ccctgagggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
 caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
 gtctccaacaaagccctcccagcccccatcgagaaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
 acacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaaggcttctatcccagc
 5 gacatcgccgtggagtgaggagagcaatgggcagccggagacaactacaagaccacgcctcccgtgctggactccgac
 ggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
 catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

10 **2H7 scFv MTH (CSC) WTCH2CH3 (amino acid sequence)** (SEQ ID NO: 166)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
 WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
 CQQWSFNPPTFGAGTKLELKDGGGSGGGGSSQAYLQQSGAELVRPG
 ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
 15 KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFYFDVWGTGTT
 VTVSSDQEPKSCDKTHTSPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV
 VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
 WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSL
 TCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRW
 20 QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

2H7 scFv MTH (CCS) WTCH2CH3 (nucleotide sequence) (SEQ ID NO: 167)

aagcttgccgccatggattttcaagtcagattttcagcttcctgctaatactgcttcagtcataattgccagaggacaaattg
 25 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
 ttacatgcactggtaccagcagaagccaggatcctccccaaacctggatttatgccccatccaacctggcttctggagtc
 cctgctcgttcagtggcagtggtctgggacctcttactctctcacaatcagcagagtggaggctgaagatgctgccactt
 attactgccagcagtgaggtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
 ggcggtggatctggaggaggtgggagctctcaggcttatctacagcagctctgggctgagctggtgaggcctgggg
 30 cctcagtgaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttaagggaaggccacactg

actgtagacaaatccctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcaggag
cccaaatcttgtgacaaaactcacacatgtccaccgtccccagcacctgaactcctggggggaccgtcagtcttctcttcc
cccaaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtgggtggacgtgagccacgaaga
5 ccctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagta
caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
acaccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggctcaaggcttctatcccagc
gacatcgccgtggagtgggagagcaatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgac
10 ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctctcatgctccgtgatg
catgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

AA

2H7 scFv MTH (CCS) WTCH2CH3 (amino acid sequence) (SEQ ID NO: 168)

15 MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
CQQWSFNPPTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
20 VTVSSDQEPKSCDKTHTCPPSPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD
WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

25

NT

HuIgAHlgA-T4-ORF (nucleotide sequence) (SEQ ID NO: 169)

tgatcagccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgac
tgtcactgcaccgaccggccctcgaggacgtgctcttaggttcagaagcagtcctcacgtgcacactgaccggcctgaga
30 gatgcctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgtgttcaaggaccacctgaccgtgacctctgt
ggctgctacagcgtgtccagtgtctgccgggctgtgccgagccatggaacctgggaagaccttcacttgactgctgcc

taccccgagtgccaagacccccgctaaccgccaccctctcaaaatccggaaacacattccggcccagggtccacctgctgcc
 gccgccgtcggaggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtg
 ctggttcgctggctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccgagcaggagcccagc
 cagggcaccaccaccttcgctgtgaccagcactgcgcgtggcagccgaggactggaagaagggggacaccttctct
 5 gcatggtgggccacgagggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgt
 caatgtgtctgtgtcatggcggaggtggacgcggatccttcgaac

AA

HuIgAHlgA-T4-ORF (amino acid sequence) (SEQ ID NO: 170)

10 dqpvpstptpsptpspspschprlslhrpaledlllgseailtctltglrdasgvftwtpssgksavqgppdrdlcgc
 ysvssvlpgcaepwnhgtftctaaypeskptlatlsksgntfrpevhllpppseelalnelvtltclargfspkdvlvrvl
 qgsqelprekyltwasrqepsqgtttfavtsilrvaaedwkkgdtfscmvghealplftqktidrlagkpthvnsvv
 maevdadpsn

15 NT

1D8-IgAH IgA-T4-CD80 (nucleotide sequence) (SEQ ID NO: 171)

aagcttatggattttcaagtgcagatttcagcttcctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgt
 cactcagctctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctccagtgtgaagtta
 catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
 20 atcgcttcagtggtcagtggttctgggacctcttattctctcgcaatcaacacatggagactgaagatgctgccacttattact
 gtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
 cgggtgggtgggtcgggtggcggggatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
 cccctgtccctcatatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt
 ctggaatggatgggaataatatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
 25 acacctccaagagccaagttttcttaaaaatcaacagtctgcaaactgatgacacagccatgtattactgtgccagaatccac
 ttgattactggggccaaggagtcaggtcagctcctctgatcagccagttccctcaactccacctaccccatctccctca
 actccacctaccccatctccctcatgtgccacccccgactgtcactgcaccgaccggccctcgaggacctgctcttaggtt
 cagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcaggtgtcaccttcacctggacgccctcaagtggg
 aagagcgctgttcaaggaccactgaccgtgacctctgtggctgtacagcgtgtccagtgtcctgccgggctgtgccgag
 30 ccatggaacatgggaagaccttcacttgcactgtgcctaccccgagtccaagacccccgctaaccgccaccctctcaaaa
 tccggaaacacattccggcccagggtccacctgtgccgccgccgtcggaggagctggccctgaacgagctggtgacg

ctgacgtgcctggcacgtggcttcagccccaaggatgtgctgggtcgtggctgcaggggtcacaggagctgccccgcg
 agaagtacctgacttgggcatcccggcaggagcccagccagggcaccaccaccttcgtgtgaccagcatactgcgcgt
 ggcagccgaggactggaagaagggggacaccttctcctgcatggtgggccacgagggcctgccgctggccttcacaca
 gaagaccatcgaccgcttggcgggtaaacccacccatgtcaatgtgtctgttgcattggcggaggtggacgcggatccttc
 5 gaacaacctgctcccatcctgggccattacctaatactcagtaaattggaattttgtgatatgctgcctgacactactgctttgcc
 caagatgcagagagagaaggaggaatgagagattgagaagggaagtgtacgccctgtataaatcgatac

AA

1D8 scFv IgAH IgA-T4-CD80 (amino acid sequence) (SEQ ID NO: 172)

10 MDFQVQIFSLLISASVIMSRGVDIVLTQSPTTIAASPGEKVTITCRASSSVSYM
 YWYQQKSGASPKLWIYDTSKLASGVPNRFSGSGSGTSYSLAINTMETEDAAT
 YYCQQWSSTPLTFGSGTKLEIKRGGGGSGGGGSGGGGSQVQLKEAGPGLVQP
 TQTLSLTCTVSGFSLTSDGVHWIRQPPGKGLEWMGIYYDGGTDYNSAIKSRL
 SISRDTSKSQVFLKINSLQTDDETAMYYCARIHFDYWGQGMVTVSSDQPVPS
 15 TPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAILTCTLTGLRDASGVTFT
 WTPSSGKSAVQGPDRDLGCGYSVSSVLPGCAEPWNHGKTFTCTAAYPESKT
 PLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFSPKDVLRWLQ
 GSQELPREKYLTWASRQEPSQGTTFFAVTSILRVAEDWKKGDTFSCMVGHE
 ALPLAFTQKTIDRLAGKPTHVNVSVVMAEVDADPSNNLLPSWAITLISVNGIF
 20 VICCLTYCFAPRCRERRRNERLRRESVRPV

NT

human IgE Fc (CH2-CH3-CH4) ORF (nucleotide sequence) (SEQ ID NO: 173):

tgatcacgtctgctccagggacttcaccccccccaccgtgaagatcttacagtcgtcctgcgacggcggcgggcacttccc
 25 cccgaccatccagctcctgtgcctcgtctctgggtacaccccagggactatcaacatcacctggctggaggacgggcagg
 tcatggacgtggacttgcaccgcctctaccacgcaggaggggtgagctggcctccacacaaagcgagctcacctcagc
 cagaagcactggctgtcagaccgcctacacctgccaggtcacctatcaaggtcacacctttgaggacagcaccaagaa
 gtgtgcagattccaacccgagaggggtgagcgcctacctaaagccggcccagcccgttcgacctgttcacgcaagtcgc
 ccacgatcacctgtctgggtggacctggcaccagcaaggggaccgtgaacctgacctggtcccgggccagtgggaa
 30 gcctgtgaaccactccaccagaaaggaggagaagcagcgcgaatggcacgttaaccgtcacgtccaccctgccgggtggg
 caccggagactggatcgagggggagacctaccagtgagggtgacccacccccacctgccaggggcctcatcgcggtc

cacgaccaagaccagcggcccgctgctgccccggaagtctatgcgtttgcgacgccggagtggccggggagccggg
acaagcgcacccctgcctgcctgatccagaacttcctgcctgaggacatctcggcgcagtggctgcacaacgaggtgcag
ctcccgacgccccggcacagcacgacgcagccccgcaagaccaagggtccggcttcttcgtcttcagccgcctggag
gtgaccagggccgaatgggagcagaaagatgagttcatctgccgtgcagtccatgaggcagcgagccccctcacagacc
5 gtccagcgagcgggtgctgtaaatcccggtaaagcggatcctcgaa

AA

human IgE Fc (CH2-CH3-CH4) ORF (amino acid sequence) (SEQ ID NO: 174):

DHVCSRDFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTINITWLEDGQV
10 MDVDLSTASTTQEGELASTQSELTLSQKHWLSDRITYTCQVTYQGHTFEDSTK
KCADSNPRGVSAYLSRPSFDLFIRKSPTITCLVVDLAPSKGTVNLTWSRASGK
PVNHSTRKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTHPHLPRALMRS
TTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDISVQWLHNEVQ
LPDARHSTTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAVHEAASPSQTV
15 QRAVSVNPGKADPS

NT

1D8 scFv-human IgE Fc (CH2-CH3-CH4)-CD80 (nucleotide sequence) (SEQ ID NO: 175)

aagcttatggattttcaagtcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctcctcaacaaccatagctgcattctccaggggagaaggtcaccatcacctgccgtgccagctccagtgtgaagtta
catgtactggtaccagcagaagtcaggcgcctccctaaactctggatttatgacacatccaagctggcttctggagttccaa
atcgcttcagtggcagtggtctgggacctcttattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
gtcagcagtgaggtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcgggtggctcggg
25 cgggtgggtgggtcgggtggcggcgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
ccctgtccctcatatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctcaagagccaagtttcttaaaaatcaacagctctgaaactgatgacacagccatgtattactgtccagaatccac
tttgattactggggccaaggagtcaggtcacagctcctctgatcacgtctgctccagggacttcacccgccaccgtga
30 agatcttacagtcgtcctgcgacggcggcgggcacttcccccgacctccagctcctgtgcctcgtctctgggtacaccc
cagggactatcaacatcacctggctggaggacgggcaggtcatggacgtggactgtccaccgcctctaccacgcagga

ggggtgagctggcctccacacaaagcgagctcaccctcagccagaagcactggctgtcagaccgcacctacacctgccag
 gtcacctatcaaggtcacacctttgaggacagaccaagaagtgtgcagattccaacccgagaggggtgagcgcctacct
 aagccggcccagcccgttcgacctgttcacccgaagtcgcccacgatcacctgtctgggtggacctggcaccagca
 aggggaccgtgaacctgacctgggtcccgggccagtgggaagcctgtgaaccactccaccagaaaggaggagaagcag
 5 cgcaatggcacgttaaccgtcacgtccaccctgccgggtgggcacccgagactggatcgagggggagacctaccagtgc
 aggggtgacccacccccacctgccagggccctcatcggtccacgaccaagaccagcggcccgctgctgccccgga
 agtctatgcgtttgcgacgccggagtggccggggagccgggacaagcgcaccctcgcctgcctgatccagaactcatg
 cctgaggacatctcgggtgcagtggctgcacaacgaggtgcagctcccgacgccgggcacagcacgacgcagccccg
 caagaccaagggtccggcttctcgtcttcagccgcctggaggtgaccagggccgaatgggagcagaaagatgagttc
 10 atctgccgtgcagtccatgaggcagcgagccctcacagaccgtccagcgagcgggtgtctgtaaattcccggtaaagcg
 atccttcgaagctcccatcctgggccattaccttaatctcagtaaatggaattttgtgatatgctgcctgacctactgctttgcc
 ccaagatgcagagagagaaggaggaatgagagattgagaagggaagtgtacgccctgtataaatcgata

AA

15 **1D8-scFv-human IgE Fc (CH2-CH3-CH4)-CD80 (amino acid sequence) (SEQ ID NO: 176)**

MDFQVQIFSLLISASVIMSRGVDIVLTQSPTTIAASPGEKVTITCRASSSVSYM
 YWYQQKSGASPKLWIYDTSKLASGVPNRFSGSGSGTSYSLAINTMETEDAAT
 YYCQQWSS'TPLTFGSGTKLEIKRGGGGSGGGGSGGGGSQVQLKEAGPGLVQP
 20 TQTLSLTCTVSGFSLTSDGVHWIRQPPGKGLEWMGIYYDGGTDYNSAIKSRL
 SISRDTSKSQVFLKINSLQTDDTAMYYCARIHFDYWGQGMVTVSSDHVCSR
 DFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTINITWLEDGQVMDVDLS
 TASTTQEGELASTQSELTLSQKHWLSDRITYTCQVTYQGHTFEDSTKKCADSN
 PRGVSAYLRSRSPFDLFIRKSPTITCLVVDLAPSKGTVNLTWSRASGKPVNHST
 25 RKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTHPHLPRALMRSTTKTSG
 PRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDISVQWLHNEVQLPDARH
 STTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAVHEAASPSQTVQRAVSV
 NPGKADPSKLPWAITLISVNGIFVICCLTYCFAPRCRERRRNERLRRESVRPV

30 NT

5B9-IgAH IgA-T4-CD80 (nucleotide sequence) (SEQ ID NO: 177)

aagcttgccgccatgaggttctctgctcagcttctggggctgcttgtgctctggatccctggatccactgcagatattgtgatg
 acgcaggctgcatttccaatccagtcactcttgaacatcagcttccatctcctgcaggtctagtaagagtctcctacatagt
 aatggcatcacttatttatttggtatctgcagaagccaggccagctcctcagctcctgatttatcagatgtccaaccttgccctc
 aggagtcacagacaggttcagtagcagtggttcaggaactgatttcacactgagaatcagcagagtggaggtgaggatg
 5 tgggtgtttattactgtgctcaaaatctagaacttccgctcacgttcggtgctgggaccaagctggagctgaaacgggggtg
 cgggtggctcgggcggtggtgggtcgggtggcggcggtatgctcacagggtgcagctgaagcagtcaggacctggcctagt
 gcagtcctcacagagcctgtccatcacctgcacagtctctggtttctcattaactacatgtgtacactgggttcgccagtct
 ccaggaaagggtctggagtggtggtggtgatatggagtgggtggaatcacagactataatgcagctttcatatccagactg
 agcatcaccaaggacgattccaagagccaagttttctttaaataaacagtctgcaacctaataacacagccatttattactgt
 10 gccagaaatgggggtgataactacccttattactatgctatggactactgggtcaaggaacctcagtcaccgtctcctctga
 tcagccagttccctcaactccactaccccatctccctcaactccactaccccatctccctcatgtgccacccccgactgt
 cactgcaccgaccggccctcaggagctgctcttaggttcagaagcgtcctcacgtgcacactgaccggcctgagagat
 gcctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgctgtcaaggaccacctgacctgacctctgtggc
 tgtacagcgtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttgcactgtgcctac
 15 cccgagtcacaagacccccgtaaccgccacctctcaaaatccggaacacattccggccccgaggtccacctgctgccgc
 cggcgtcggaggagctggccctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagcccaaggatgtgct
 ggttcgtggtgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccgccaggagcccagcca
 gggcaccaccaccttcgtgtgaccagcactgctgctggcagccgaggactggaagaagggggacaccttctcctgc
 atggtgggccacgaggccctgccgctggccttcacagaagaccatcgaccgcttggcgggtaaacccacctatgca
 20 atgtgtctgtgtcatggcggaggtggacgcggatccttgaacaacctgctcccatcctgggccattaccttaatctcagta
 aatggaattttgtgatgtgctgacctactgcttggcccaagatgcagagagagaaggaggaatgagagattgagaa
 gggaaagtgtacgccctgtataaatcgatac

AA

25 **5B9-IgAH IgA-T4-CD80 (amino acid sequence)** (SEQ ID NO: 178)

MRFSAQLLGLLVLWIPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSNG
 ITYLYWYLQKPGQSPQLLIYQMSNLAGVPPDRFSSSGSGTDFTLRISRVEAD
 VGVYYCAQNLELPLTFGAGTKLELKRGGGSGGGGSGGGGSSQVQLKQSGP
 GLVQSSQSLSITCTVSGFSLTTYAVHWVRQSPGKGLEWLGVWSGGITDYNA
 30 AFISRLSITKDDSKSQVFFKMNSLQPNDTAIYYCARNGGDNYPIYYAMDIW
 GQGTSVTVSSDQVPSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAIL
 TCTLTGLRDASGVFTFTWTPSSGKSAVQGPPDRDLGCYSVSSVLPGCAEPWN

gaatthttgtgatatgctgcctgacctactgctttgcccccaagatgcagagagagaaggaggaatgagagattgagaaggg
aaagtgtacgccctgtataaatcgata

AA

5 **5B9-scFv-human IgE Fc (CH2-CH3-CH4)-CD80 (amino acid sequence) (SEQ ID NO: 180)**

MRFSAQLLGLLVLWIPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSNG
ITYLYWYLQKPGQSPQLLIYQMSNLAGVDPDRFSSSGSGTDFTLRISRVEAED
VGVYYCAQNLELPLTFGAGTKLELKRGGGGSGGGGSGGGGSSQVQLKQSGP
10 GLVQSSQSLSITCTVSGFSLTTYAVHWVRQSPGKGLEWLGVIWSGGITDYNA
AFISRLSITKDDSKSQVFFKMNSLQPNDAIYYCARNGGDNYPYYAMDYW
GQGTSVTVSSDHVCSRDFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTI
NITWLEDGQVMDVDLSTASTTQEGELASTQSELTLSQKHWLSDRTYTCQVTY
QGHTFEDSTKKCADSNPRGVSAAYLSRPSFDLFIRKSPTITCLVVDLAPSKGTV
15 NLTWSRASGKPVNHSTRKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTH
PHLPRALMRSTTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDIS
VQWLHNEVQLPDARHSTTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAV
HEAASPSQTVQRAVSVNPGKADPSKLPWAITLISVNGIFVICCLTYCFAPRCR
ERRRNERLRRESVRPV

20

NT

2e12-scFv-IgAH IgA-T4-CD80 (nucleotide sequence) (SEQ ID NO: 181)

aagcttatggattttcaagtcagattttcagcttctgctaatacagtcgctcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttctttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgtgaatatta
25 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
tctgggggtccctgccaggttttagtggcagtggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaagtaggaaggttccttgacgttcggtggaggcaccaagctggaaatcaaacgggggtgg
cgggtggctcgggcggaggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcct
30 ccaggaaagggtctggagtggctgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactg
agcatcaccaaggacaactccaagagccaagttttcttaaaaatgaacagctctgcaaactgatgacacagccagataactact

gtgccagagatggttatagtaactttcattactatgttatggactactgggggtcaaggaacctcagtcaccgtctcctcagatc
agccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtca
ctgcaccgaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgc
ctcaggtgtcaccttcacctggacgcccctcaagtgggaagagcgctgtcaaggaccacctgacctgacctctgtggctg
5 ctacagcgtgtccagtgctctgccgggctgtgccgagccatggaacctgggaagaccttcacttgactgctgcctacc
cgagtccaagacccccgtaaccgccaccctctcaaatccggaacacattccggcccagggtccacctgctgccgccg
ccgtcggaggagctggccctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagcccaaggatgtgctgg
ttcgtggctgcaggggtcacaggagctgccccgcgagaagtacgtgacttgggcatccggcgaggagcccagccagg
gcaccaccaccttcgtgtgaccagcatactgcgcgtggcagccgaggactggaagaagggggacaccttctcctgcat
10 ggtgggccacgaggccctgccgtggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgtcaat
gtgtctgttgcattggcggaggtggacgcggatccttcgaacaacctgctcccatcctgggccattaccttaatctagtaaa
tggaattttgtgatatgctgcctgacctactgctttccccaagatgcagagagagaaggaggaatgagagattgagaagg
gaaagtgtacgccctgtataaatcgatac

15 AA

2e12-scFv-IgAH IgA-T4-CD80 (amino acid sequence) (SEQ ID NO: 182)

MDFQVQIFSLLISASVIMSRGVDIVLTQSPASLAVSLGQRATISCRASESVEYY
VTSLMQWYQQKPGQPPKLLISAASNVESGVPARFSGSGSGTDFSLNIHPVEED
DIAMYFCQQSRKVPWTFGGGKLEIKRGGGGSGGGGSGGGGSQVQLKESGP
20 GLVAPSQSLITCTVSGFSLTGYGVNWVRQPPGKLEWLGMWGDGSTDYNS
ALKSRLSITKDNSKSQVFLKMNSLQTDDTARYYCARDGYSNFHYVMDYW
GQGTSVTVSSDQVPSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEAIL
TCTLTGLRDASGVTFWTPSSGKSAVQGPPDRDLCGCYSVSSVLPGCAEPWN
HGKTFTCTAAYPESKTPLTATLSKSGNTFRPEVHLLPPPSEELALNELVTLTCL
25 ARGFSPKDVLVRWLQGSQELPREKYLTWASRQEPSQGTTTTFAVTSILRVAAE
DWKKGDTFSCMVGHEALPLAFTQKTIDRLAGKPTHVNVSVVMAEVDADPSN
NLLPSWAITLISVNGIFVICCLTYCFAPRCRERRRNERLRRESVRPV

NT

30 **2e12-scFv-human IgE Fc (CH2-CH3-CH4)-CD80 (nucleotide sequence) (SEQ ID NO: 183)**

aagcttatggattttcaagtcagattttcagcttcctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
 caccaaatctccagcttctttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaaagtgttgatatta
 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
 tctgggggtccctgccaggtttagtggcagtggggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
 5 ttgcaatgtatttctgtcagcaaagtaggaaggttccttggacgttcgggtggaggcaccaagctggaaatcaaacggggtgg
 cgggtggctcgggagggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
 cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcct
 ccaggaaagggtctggagtggtgggaatgatatggggtgatggaagcacagactataatcagctctcaaatccagactg
 agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgcaaactgatgacacagccagataactact
 10 gtccagagatggttatagtaactttcattactatgttatggactactggggtcaaggaacctcagtcaccgtctcctcagatc
 acgtctgctccaggagcttcaccccgcccaccgtgaagatcttacagtcgtcctgcgacggcggcgggcacttcccccg
 accatccagctcctgtgcctcgtctctgggtacacccagggactatcaacatcacctggctggaggacgggcaggtcatg
 gacgtggacttgtccaccgcctctaccacgcaggagggtgagctggcctccacaaaagcgagctcacctcagccaga
 agcactggctgtcagaccgcacctacacctgccaggtcacctatcaaggtcacacctttgaggacagcaccaagaagtg
 15 gcagattccaacccgagaggggtgagcgcctacctaagccggcccagcccgttcgacctgttcacccgaagtcgccc
 cgtcacctgtctggtgggtggacctggcaccagcaaggggaccgtgaacctgacctgggtcccgccagtggaagc
 ctgtgaaccactccaccagaaaggaggagaagcagcgcaatggcacgttaaccgtcacgtccaccctgccgggtgggca
 cccgagactggatcgagggggagacctaccagtgcagggtgacccaccccccacctgccaggggccctcatcggtcca
 cgaccaagaccagcggcccgctgctgccccggaagtctatgcgtttgcgacgccggagtggccggggagccgggac
 20 aagcgcaccctgcctgcctgatccagaacttcagcctgaggacatctcgggtgcagtggtgcacaacgaggtgcagct
 cccggagccccggcacagcacgacgcagccccgcaagaccaagggtccggcttcttcgtcttcagccgcctggaggt
 gaccagggccgaatgggagcagaaagatgagttcatctgccgtgcagtcctatgaggcagcgagccccctcacagaccgt
 ccagcgagcgggtgtctgtaaatcccggtaaagcggatccttcgaagctccatcctgggccattacctaatactcagtaaatg
 gaattttgtgatatgctgcctgacctactgctttgccccagatgcagagagagaaggagggaatgagagattgagaagg
 25 aaagtgtacgccctgtataaatcgata

AA

2e12-scFv-human IgE Fc (CH2-CH3-CH4)-CD80 (amino acid sequence) (SEQ ID
NO: 184)

30 MDFQVQIFSLLISASVIMSRGVDIVLTQSPASLAVSLGQRATISCRASESVEYY
 VTSLMQWYQQKPGQPPKLLISAASNVESGVPARFSGSGSGTDFSLNIHPVEED
 DIAMYFCQQSRKVPWTFGGGTLKLEIKRGGGSGGGGSGGGGSGVQLKESGP

GLVAPSQSL SITCTVSGFSLTGYGVNWVRQPPGKGLEWLGMWGDGSTDYNS
 ALKSRLSITKDNSKSQVFLKMNSLQTDDTARYYCARGYSNFHYVMDYW
 GQGTSVTVSSDHVCSRDFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTI
 NITWLEDGQVMDVDLSTASTTQEGELASTQSELTLSQKHWLSDRTYTCQVTY
 5 QGHTFEDSTKKCADSNPRGVSA YLSRSPFDLFIRKSPTITCLVVDLAPSKGTV
 NLTWSRASGKPVNHSTRKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTH
 PHLPRALMRSTTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFM PEDIS
 VQWLHNEVQLPDARHSTTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAV
 HEAASPSQTVQRAVSVNPGKADPSKLPSWAITLISVNGIFVICCLTYCFAPRCR
 10 ERRRNERLRRESVRPV

NT

500A2 scFv (nucleotide sequence) (SEQ ID NO: 185)

atgttgatacatctcagctccttgggctttactcttctggattcagcctccagaagtacatagtctgactcagactccagc
 15 cactctgtctctaattctggagaaagagtcacaatgacctgaagaccagtcagaatattggcacaatcttacactggtatca
 ccaaaaaccaaaggaggctccaagggtctcatcaagtatgcttcgcagtcattcctgggatccccctccagattcagtggc
 agtgggttcggaacagatttactctcagcatcaataacctggagcctgatgatcggaaattattactgtcaacaaagtaga
 agctggcctgtcacgttcggctcctggcaccaagctggagataaaacggggtggcggtggctcgggcggaggtgggtcg
 ggtggcggcggatctcaggtcaagctgcagcagtcgggttctgaactagggaacctggggcctcagtgaaactgtcctg
 20 caagacttcaggctacatattcacagatcactatatttctgggtgaaacagaagcctggagaaagcctgcagtgatagga
 aatgtttatggtggaaatggtgtacaagctacaatcaaaaattccaggggcaaggccacactgactgtagataaaatctctag
 cacagcctacatggaactcagcagcctgacatctgaggattctgccatctattactgtgaagaaggccggtagcgacggg
 ccatgctatggactactggggtcaggggatccaagttaccgtctcctctgatc

25 AA

500A2 scFv (amino acid sequence) (SEQ ID NO: 186)

MLYTSQLLGLLLFWISASRSDIVLTQTPATLSLIPGERVTMTCKTSQNIGTILH
 WYHQKPKEAPRALIKYASQSIPGIPSRFSGSGSETDFTLSINNLEPDDIGIYYCQ
 QSRSWPVTFPGTKLEIKRGGGGSGGGGSGGGGSQVKLQQSGSELGKPGASV
 30 KLSCKTSGYIFTDHYISWVKQKPGESLQWIGNVYGGNGGTSYNQKFQ GKATL
 TVDKISSTAYMELSSLTSEDSAIYYCARRPVATGHAMDYWGQGIQVTVSSD

NT

5' oligo:

Name : hIgAbcl5 (SEQ ID NO: 662)

5 Sequence : GTTGTTGATCAGCCAGTTCCTCAACTCCACCTACC

3' oligo:

Name : IgA3BB (SEQ ID NO: 663)

GTTGTTTTTCGAAGGATCCGCGTCCACCTCCGCCATGACAACAGA

10

5' oligo:

Name : IgGWT3 (SEQ ID NO: 187)

GTTGTTTTTCGAAGGATCCGCTTTACCCGGAGACAGGGAGAGGCTCTT

15 3' oligo:

Name : hIgGWT5 (SEQ ID NO: 188)

GTTGTTAGATCTGGAGCCCAAATCTTGTGACAAAACCTCACACATG

5' oligo:

20 Name : FADD5 (SEQ ID NO: 189)

Sequence :

GTTGTGGATCCTTCGAACCCGTTCTTGGTGCTGCTGCACTCGGTGTCG

3' oligo:

25 Name : FADD3 (SEQ ID NO: 190)

Sequence :

GTTGTTATCGATCTCGAGTTATCAGGACGCTTCGGAGGTTAGATGCGTC

NT

FADD-CSSCFV (nucleotide sequence) (SEQ ID NO: 191):

gtggatccttcgaacccgttcctggtgctgctgcactcgggtgctccagcctgtcgagcagcgagctgaccgagctcaag
5 ttcctatgcctcgggcgcgtgggcaagcgcaagctggagcgcgtgcagagcggcctagacctcttctccatgctgctgga
gcagaacgacctggagccccgggcacaccgagctcctgcgcgagctgctgcctccctgcggcgccacgacctgctgcg
gcgcgtcgacgacttcgagggcgggggcggcggccggggccgcgcctggggaagaagacctgtgtgcagcatttaacg
tcatatgtgataatgtggggaaagattggagaaggctggctcgtcagctcaaagtctcagacaccaagatcgacagcatcg
aggacagatacccccgcaacctgacagagcgtgtgcgggagtcactgagaatctggaagaacacagagaaggagaaac
10 gcaacagtggcccacctggtgggggctctcaggtcctgccagatgaacctggtggctgacctggtacaagaggttcagca
ggccccgtgacctccagaacaggagtgggggccatgtccccgatgtcatggaactcagacgcattctacctccgaagcgctct
gataactcgagatcgataacaac

AA

15 **FADD-CSSCFV (amino acid sequence)** (SEQ ID NO: 192):

VDPSNPFLVLLHSVSSSLSSSELTFLKFLCLGRVVGKRLERVQSGLDLFSMLLE
QNDLEPGHTELLRELLASLRRHDLLRRVDDFEAGAAAGAAPGEEDLCAAFNV
ICDNVGKDWRRRLARQLKVSDTKIDSIEDRYPRNLTERVRESLRIWKNTKEN
ATVAHLVGALRSCQMNLVADLVQEVQQARDLQNRSGAMSPMSWNSDASTS
20 EAS

NT

HCD28tm5B (nucleotide sequence) (SEQ ID NO: 193)

GTTGTGGATCCTCCCTTTTGGGTGCTGGTGGTGGTTGGTGTCTCCTGGCTTGC
25 TATAGCTTG

NT

HCD28tm3S (nucleotide sequence) (SEQ ID NO: 194)

GTTGTTTCGAACCCAGAAAATAATAAAGGCCACTGTTACTAGCAAGCTAT
AGCAAGCCAG

NT

5 **HCD28tm5'** (nucleotide sequence) (SEQ ID NO: 195)

GTTGTGGATCCTCCCTTTTGGGTGCTGGTGGT

NT

HCD28tm3' (nucleotide sequence) (SEQ ID NO: 196)

10 GTTGTTTCGAACCCAGAAAATAATAAAGGCCAC

NT

HCD80tm5' (nucleotide sequence) (SEQ ID NO: 197)

GTTGTGGATCCTCCTGCTCCCATCCTGG

15

NT

HCD80tm3' (nucleotide sequence) (SEQ ID NO: 198)

GTTGTTTCGAACGGCAAAGCAGTAGGTCAGGC

20 NT

MFADD5BB (nucleotide sequence) (SEQ ID NO: 199)

GTTGTGGATCCTTCGAACCCATTCCTGGTGCTGCTGCACTCGCTG

NT

25 **MFADD3XC** (nucleotide sequence) (SEQ ID NO: 200)

GTTGTTATCGATCTCGAGTCAGGGTGTTTCTGAGGAAGACAC

NT

Murine FADD Nucleotide sequence (full length, but without flanking –Ig or transmembrane sequences) (**nucleotide sequence**) (SEQ ID NO: 201):

5 gtggatccttgaacatggacccattcctgggtgctgctgcactcgtgtccggcagcctgtcgggcaacgatctgatggag
ctcaagttcttgtgccgcgagcgcgtgagcaaacgaaagctggagcgcgtgcagagtggcctggacctgttcacggtgct
gctggagcagaacgacctggagcgcgggcacaccgggctgctgcgcgagttgctggcctcgtgcgccgacacgatct
actgcagcgcctggacgacttcgaggcggggacggcgaccgctgcgccccgggggaggcagatctgcaggtggcat
ttgacattgtgtgtgacaatgtggggagagactggaaaagactggcccgcgagctgaagggtgtctgaggccaagatggat
10 gggattgaggagaagtacccccgaagtctgagtgagcgggtaaggagagctctgaaagtctggaagaatgctgagaag
aagaacgcctcgggtggccgactggtcaaggcgtgcggacctgcaggctgaatctggtggctgacctggtggaagaa
gcccaggaatctgtgagcaagagtgagaatatgtccccagtactaagggttcaactgtgtcttctcagaaacaccctgac
tcgagatcgat

15 AA

Murine FADD (amino acid sequence) (SEQ ID NO: 202)

VDPSNMDPFLVLLHSLSGSLSGNDLMELKFLCRERVSKRKLERVQSGLDLFT
VLLEQNDLERGHTGLLRELLASLRRHDLLQRLDDFEAGTATAAPPGEADLQV
AFDIVCDNVGRDWKRLARELKVSEAKMDGIEEKYPRSLSERVRESLKVWKN
20 AEKKNASVAGLVKALRTCRLNLVADLVEEAQESVSKSENMSPVLRDSTVSSS
ETP

NT

MCASP3-5 (nucleotide sequence) (SEQ ID NO: 203)

25 GTTGTGGATCCTTCGAACATGGAGAACAACAAAACCTCAGTGGATTCA

NT

MCASP3-3 (nucleotide sequence) (SEQ ID NO: 204)

GTTGTTATCGATCTCGAGCTAGTGATAAAAGTACAGTTCTTTCGT

NT

MCASP8-5 (nucleotide sequence) (SEQ ID NO: 205)

GTTGTTTCGAACATGGATTTCCAGAGTTGTCTTTATGCTATTGCTG

5 NT

MCASP8-3 (nucleotide sequence) (SEQ ID NO: 206)

GTTGTTATCGATCTCGAGTCATTAGGGAGGGAAGAAGAGCTTCTTCCG

NT

10 **hcasp3-5**(nucleotide sequence) (SEQ ID NO: 207)

GTTGTGGATCCTTCGAACATGGAGAACACTGAAAACCTCAGTGGAT

NT

hcasp3-3 (nucleotide sequence) (SEQ ID NO: 208)

15 GTTGTTATCGATCTCGAGTTAGTGATAAAAATAGAGTTCTTTTGTGAG

NT

hcasp8-5 (nucleotide sequence) (SEQ ID NO: 209)

GTTGTGGATCCTTCGAACATGGACTTCAGCAGAAATCTTTATGAT

20

NT

hcasp8-3 (nucleotide sequence) (SEQ ID NO: 210)

GTTGTTATCGATGCATGCTCAATCAGAAGGGAAGACAAGTTTTTTTCT

NT

HuIgGMHWC (nucleotide sequence) (SEQ ID NO: 648)

gtt gtt gat cag gag ccc aaa tct tct gac aaa act cac aca tct cca ccg tcc cca gca cct gaa etc
ctg ggt gga ccg tca gtc ttc c

5

NT

2H7-human IgE (CH2-CH3-CH4) (nucleotide sequence) (SEQ ID NO: 128)

aagcttgccgcatggattttcaagtgcagattttcagcttcctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaagggtcacaatgacttgcagggccagctcaagtgaag
10 ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggacctcttactctctcacaatcagcagagtggaggctgaagatgctgccactt
attactgccagcagtgaggttttaacccaccacggtcgggtgctgggaccaagctggagctgaaaggtggcgggtggctcg
ggcgggtgggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
15 gggcctggaatggattggagctatattatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggctctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctgatcacgtctgct
ccagggacttcaccccgccaccgtgaagatcttacagtcgtctcgacggcgggcggcacttcccccgaccatcca
gctcctgtgcctcgtctctgggtacaccccagggactatcaacatcacctggctggaggacgggcaggtcatggacgtgg
20 acttgtccaccgcctctaccacgcaggaggggtgagctggcctccacaaaagcgagctcacctcagccagaagcactg
gctgtcagaccgcacctacacctgccaggtcacctatcaagggtcacacctttgaggacagcaccaagaagtgtgcagattc
caacccgagaggggtgagcgcctacctaagccggcccagccggttcgacctgttcacgcaagtcgccacgatcacc
tgtctgggtgggtggacctggcaccagcaaggggaccgtgaacctgacctgggtcccgggccagtggggaagcctgtgaacc
actccaccagaaaggaggagaagcagcgcgaatggcacgttaaccgtcacgtccaccctgccgggtgggcacccgagact
25 ggatcgaggggggagacctaccagtgcagggtgacccacccccacctgccaggggcctcatgcggtccacgaccaag
accagcggcccgctgctgccccggaagtctatgcgtttcgacgccggagtggccggggagccgggacaagcgcac
cctcgcctgcctgatccagaacttcacgtgaggacatctcgggtgcagtgggtgcacaacgaggtgcagctcccgagc
ccgggcacagcacgacgcagccccgcaagaccaagggtccggcttcttctcagccgcctggaggtgaccaggg
ccgaatgggagcagaaagatgagttcatctgccgtgcagtcctatgaggcagcagccccctcacagacctccagcgagc
30 ggtgtctgtaaatcccggtaaataataatctaga

AA

2H7 scFv IgE (CH2-CH3-CH4) (amino acid sequence) (SEQ ID NO:129)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
5 CQQWSFNPPTFGAGTKLELKGGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSDHVCSDFTPPTVKILQSSCDGGGHFPPTIQLLCLVSGYTPGTINITWLE
DGQVMDVDLSTASTTQEGELASTQSELTLSQKHWLSDRTYTCQVTYQGHTF
10 EDSTKKCADSNPRGVSAYLSRPSFDLFIKSPSTITCLVVDLAPSKGTVNLTWS
RASGKPVNHSTRKEEKQRNGTLTVTSTLPVGTRDWIEGETYQCRVTHPHLPR
ALMRSTTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDISVQWL
HNEVQLPDARHSTTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAVHEAA
SPSQTVQRAVSVNPGK

15

NT

2H7 scFv MH (SSS) MCH2WTCH3 (nucleotide sequence) (SEQ ID NO:130)

aagcttgcgcccatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagtgaag
20 ttacatgcactggtaccagcagaagccaggatcctcccccaccctggattatgccccatccaacctggcttctggagtc
cctgctcgcttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgaggagctgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg
ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtctgcaaggcttctggctacacatttaccagttacaatatgcactgggtaaagcagacacctagaca
25 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaatttctgacaaaactcacacatccccaccgtcccagcacctgaactcctggggggatcgtcagttctcttctccc
cccaaaaccaagacacccctcatgatctccggaccctgaggtcacatgcgtggtggtgagctgagccacgaagac
30 cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
aacagcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
ctccaacaaagccctccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta

cacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatcccagcg
acatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctcatgctccgtgatgc
atgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaataatgatctaga

5

AA

2H7 scFv MH (SSS) MCH2WTCH3 (amino acid sequence) (SEQ ID NO:131)

MDFQVQIFSLLISASVIIARGQIVLSQSPAILSASPGEKVTMTCRASSSVSYMH
WYQQKPGSSPKPWIYAPSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYY
10 CQQWSFNPPFTFGAGTKLELKDGGGSGGGGSGGGGSSQAYLQQSGAELVRPG
ASVKMSCKASGYTFTSYNMHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKG
KATLTVDKSSSTAYMQLSSLTSEDSAVYFCARVVYYSNSYWFVDVWGTGTT
VTVSSDQEPKSSDKTHTSPSPAPELLGGSSVFLFPPKPKDTLMISRTPEVTCVV
VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
15 LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK

NT

20 **5B9 scFv MTHWTCH2CH3 (nucleotide sequence)** (SEQ ID NO: 132)

aagcttgccgcatgaggttctctgctcagcttctggggctgctgtgctctggatccctggatccactgcagatattgtgatg
acgcaggctgcattctccaatccagtcactcttgaacatcagcttccatctcctgcaggcttagtaagagctcctacatagt
aatggcatcacttatttgatttggtatctgcagaagccaggccagtcctcagctcctgatttatcagatgtccaacctgcctc
aggagtcccagacaggttcagtagcagtggtgcaggactgatttcacactgagaatcagcagagtgagggtgaggatg
25 tgggtgtttactgtgctcaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacgggggtgg
cggtggctcgggcgggtgggtgggtcgggtggcggcgatcgctcacaggtgcagctgaagcagtcaggacctggcctagt
gcagtcctcacagagcctgtccatcacctgcacagctctctggtttctattaactacatgctgtacactgggttcgccagtct
ccaggaaagggcttgagtggtgggagtgatatggagtgggtggaatcacagactataatgcagcttcatatccagactg
agcatcaccaaggacgattccaagagccaagtttctttaaaatgaacagcttgcaacctaatacacagccatttactgt
30 gccagaaatgggggtgataactacccttattactatgctatggactactgggtcaaggaacctcagtcaccgtctcctctga
tcaggagcccaaatcttctgacaaaactcacatccccaccgtccccagcacctgaactcctggggggaccgtcagctctt

cctctccccccaaaaccaagacacccctcatgatctcccgaccctgaggtcacatgctgggtggacgtgagcc
 acgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaagccgcgggagg
 agcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaa
 gtgcaaggctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaacc
 5 acaggtgtacacccctgccccatcccggtgatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttct
 atcccagcgacatcgccgtggagtgggagagcaatgggcagccggagacaactacaagaccacgcctcccgtgctgg
 actccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctcatgct
 ccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

10 AA

5B9 scFv MTHW/CH2CH3 (amino acid sequence) (SEQ ID NO: 133)

MRFSAQLLGLLVLWIPGSTADIVMTQAAFSNPVTLGTSASISCRSSKSLLSHNSG
 ITYLYWYLQKPGQSPQLLIYQMSNLASGVPDRFSSSGSGTDFTLRISRVEAED
 VGVYYCAQNLELPLTFGAGTKLELKRGGGGSGGGGSGGGGSSQVQLKQSGP
 15 GLVQSSQSLSITCTVSGFSLTTYAVHWVRQSPGKGLEWLGVWSGGITDYNA
 AFISRLSITKDDSKSQVFFKMNSLQPNDTAIYYCARNGGDNYPIYYYAMDYW
 GQGTSVTVSSDQEPKSSDKTHTSPSPAPELLGGPSVFLFPPKPKDTLMISRTPE
 VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
 LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK
 20 NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD
 KSRWQQGNVFSQSVMHENHNTQKSLSLSPGK

1. 2H7 scFv with alternative VHL11 mutations:

Nucleotide sequence (SEQ ID NO: 211)

25 Aagcttgccgcatggattttcaagtcagattttcagcttctgctaatacagtccttcagtcataattgccagaggacaaatt
 gttctctccagctctccagcaatcctgtctgcatctccaggggagaaggtcacatgacttgaggccagctcaagtgtaa
 gttacatgactggtaccagcagaagccaggatcctccccaaaccctggatttatgccccatccaacctggcttctggagt
 cctgtctgcttcagtggcagtggtctgggaccttactctctcaaatcagcagagtgaggctgaagatgctgccact
 tattactgccagcagtgagtttaacccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
 30 ggcggtggtgatctggaggaggtgggagc tctcaggcttatctacagcagctctggggctgag (**one of the
 following: tcn, acn, gan, can, aan, cgn, agn**)
 gtgaggcctggggcctcagtgaaatgtcctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagc
 agacacctagacagggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagg
 caaggccacactgactgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctcggt
 35 ctatttctgtgcaagagtgggtgtactatagtaactcttactgggtacttcgatgtctggggcacagggaccacggtcaccgtctc
 ttctgatcag

Amino acid sequence (SEQ ID NO: 212)

mdfqvqifsflisaviiargqivlsqspailsaspgekvtmtrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyycqqwsfnpptfgagtklelkdggsgggsgggssqaylqqsgae
(one of the following: s, t, d, e, q, n, r, k, h)

5 vrpgasvkmsckasgytftsymbhwwkqtpqrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlsslts
edsavy fcarvvyysnsywyfdvwtgtttvssdq

2. VHL11 deletion

Nucleotide sequence: (SEQ ID NO: 213)

10 Aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaatt
gttctctcccagctctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttgaggccagctcaagtgtaa
gttacatgcactggtagcagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagt
ccctgctcgttcagtggcagtggtctgggacctctactctctcacaatcagcagagtgagggtgaagatgctgccact
tattactgccagcagtgaggtttaacccacccacgttcgggtgctgggaccaagctggagctgaaagatggcggtggctcg
15 ggcggtgggtgatctggag- gaggtgggagc tctcaggcttatctacagcagctctggggctgaggtgaggcctgggg-
cctcagtgaaagatgctctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctattatccaggaaatggtgatacttctcacaatcagaagttaagggaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctattctgtgcaa
gaggggtgactatagtaactcttactggtagctcgtctggggcacaggaccacggtcaccgtctcttctgatcag

20 Amino acid sequence: (SEQ ID NO: 214)

mdfqvqifsflisaviiargqivlsqspailsaspgekvtmtrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyycqqwsfnpptfgagtklelkdggsgggsgggssqaylqqsgaevrp
gasvkmsckasgytftsymbhwwkqtpqrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltseds
avyfcarvvyysnsywyfdvwtgtttvssdq

25

3. 2H7 VL L106 with alternative mutations

Nucleotide sequence: (SEQ ID NO: 215)

aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttgaggccagctcaagtgtaa
30 ttacatgcactggtagcagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagt
cctgctcgttcagtggcagtggtctgggacctctactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgcca- gcagtgaggtttaac ccacccacgttcgggtgctgggaccaagctggag (tcn, agn, aan, cgn,
can, gan, and non-natural derivatives of these codons)
aaagatggcggtggctcggcggtggtggatctggaggaggtgggagctc

35 Amino acid sequence: (SEQ ID NO: 216)

mdfqvqifsflisaviiargqivlsqspailsaspgekvtmtrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfs gsgsgtsysltisrveadaatyycqqwsfnpptfgagtkle (s, t, r, k, h, q, n, d, e, and non-
natural derivatives of these amino acids at position 106) kdggsgggsgggsgggss

40 4. YL L106 deletion

Nucleotide sequence: (SEQ ID NO: 217)

aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttgaggccagctcaagtgtaa

ttacatgcactgggtaccagcagaagccaggatcctcccccaccctggatttatgccccatccaacctggcttctggagtc
cctgtcgccttcagtgagcagtggtctgggacctcttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgcttcggtgctgggaccaagctggagaaaagatggcgggtggctcgggc
ggtggtggatctggaggaggtgggagctc

5 Amino acid sequence: (SEQ ID NO: 218)

mdfqvqifsflisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwi yapsnlasg
vparfsg sgsgtsysltisrveadaatyccqqsfnptfgagtklekdggsgggsgggsgss

5. IgE CH3 CH4

10 Nucleotide sequence: (SEQ ID NO: 219)

tccaacccgagaggggtgagcgcctacctaagccggcccagcccgttcgacctgttcacccgaagtcgcccacgatca
cctgtcgggtgggtggacctggcaccagcaaggggaccgtgaacctgacctgggtccgggagtgaggagcctgtgaa
ccactccaccagaaaggaggagaagcagcgcaatggcacgttaaccgtcacgtccaccctgccgggtgggcacccgaga
ctggatcgagggggagacctaccagtgcagggtgacccacccccacctgccaggggccctcatcggtccacgaccaa
15 gaccagcggcccgcgtgctgccccggaagtctatgcgttgcgacgccggagtgccggggagccgggacaagcgca
ccctcgctgctgatccagaacttcacgtgaggacatctcggtgcagtggtgcacaacgaggtgcagctcccgac
gccccgcacagcacgacgcagccccgcaagaccaagggctccggcttctctcttcagccgcctggaggtgaccagg
gccgaatgggagcagaaagatgagttcatctgccgtgcagtcacatgaggcagcgagcccctcacagaccgtccagcga
gcggtgtctgtaaatcccgtaaatgataatctagaa

20 Amino acid sequence: (SEQ ID NO: 220)

snprgvsaylsrpspfdlfrksptitclvdlapskgtvnltwsrasgkpvnhstrkeekqrmgtltvtstlpvgrdwieg
etyqcrvthphlpralmrsttktspraapevyafatpewpgsrdrklaciqnfmpe disvqwlhnevqlpdarhstt
qprktkgsgffv fsrlevtraeweqkdeficravheaaspsqtvqravsvnpgk

25 6. hIgG1H/IgE WCH3 WCH4

Nucleotide sequence: (SEQ ID NO: 221)

tgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtccccagcatccaacccgagaggggtgagcgcct
acctaagccggcccagcccgttcgacctgttcacccgaagtcgcccacgatcacctgtctgggtggacctggcaccca
30 gcaaggggaccgtgaacctgacctgggtccgggagtgaggagcctgtgaacctccaccagaaaggaggagaag
cagcgcaatggcacgttaaccgtcacgtccaccctgccgggtgggcacccgagactggatcgagggggagacctaccag
tgcagggtgacccacccccacctgccaggggccctcatcggtccacgaccaagaccagcggcccgctgctgccccg
gaagtctatgcgttgcgacgccggagtg gccggggagccgggacaagcgacccctgcctgcctgatccagaac-
ttcatgcctgaggacatctcggtgcagtggtgcacaacgaggtgcagctcccgacccccggcacagcacgacgcag
ccccgcaagaccaagggctccggcttctctgcttccagccgcctggaggtgaccagggccgaatgggagcagaaagatg
35 agttcatctgccgtgcagtcacatgaggcagcgagcccctcacagaccgtccagcgagcgggtgtctgtaaatcccgtaaat
gataatctagaa

Amino acid sequence: (SEQ ID NO: 222)

dqepkssdkthtspspasnpgrgvsaylsrpspfdlfrksptitclvdlapskgtvnltwsrasgkpvnhstrkeekqr
ngltltvtstlpvgrdwiegetyqcrvthphlpralmrsttktspraapevyafatpewpgsrdrklaciqnfmpe dis
40 vqwlhnevqlp darhsttqprktkgsgffv fsrlevtraeweqkdeficravheaaspsqtvqravsvnpgk

7. IgE WCH2 WCH3 WCH4

Nucleotide sequence: (SEQ ID NO: 223)

Tgatcacgtctgtccagggaattcacccccgccaccgtgaagatcttacagtcgtctgcgacggcgggcggaacttcc
ccccgaccatccagctcctgtgccctgtctctgggtacacccagggaactatcaacatcacctggctggaggacgggcag
gtcatggacgtggacttgtccaccgcctctaccacgcaggaggggtgagctggcctccacacaaagcgagctcacctcag
ccagaagcactggctgtcagaccgcacctacacctgccagggtcacctatcaaggtcacaccttgaggacagccaaga
agtgtgcagattccaacccgagaggggtgagcgcctacctaaagccggccagcccgttcgacctgttcacccgaagtcg
cccacgatcacctgtctggtgggtggacctggcaccagcaaggggaccgtgaacctgacctgggtcccgggccaagtggga
agcctgtgaaccactccaccagaaaggaggagaagcagcgcaatggcacgttaaccgtcacgtccaccctgccggtgg
gcacccgagactggatcgagggggagacctaccagtgcagggtgacccacccccacctgcccagggccctcatgcggt
ccacgaccaagaccagcggcccgcgtgctgccccggaagtctatgcgtttgcgacgccggagtggccggggagccgg
gacaagcgcaccctgcctgcctgatccagaacttcatgcttgaggacatctcgggtgcagtgggtgcacaacgaggtgca
gtccccggacgccgggcacagcacgacgcagccccgcaagaccaaggggtccggcttcttctgtcttcagccgcctgga
ggtgaccagggccgaatgggagcagaaagatgagttcatctgccgtgcagtccatgaggcagcgagcccctcacagac
cgtccagcgagcgggtgtctgtaaatccccggtaaatgataatctaga

Amino acid sequence: (SEQ ID NO: 224)

dhvcsrdftpptvkilqsscdggghfpptiqlclvsgytpgtinitwledgqvmdivlstaattqegelastqseltlsqk
hwlstdrtytcqvtyqghtfedstkkcadsnprgvsaylsrpsfdlfrksptitclvdlapskgtnltsrasgkpv
hstrkeekqrngltvtstlpvgtrdwiegetyqcrvthphlpralmrsttktsgpaaapevyafatpewpgsrdkrtlacli
qnfmpedisvqwlhnevqlpdarhsttqprkktgsgffvfrlevtraeweqkdeficravheaaspsqtvqravsvnp
gk

8. hIgG1H/IgE CH3 CH4 (ORF)

Nucleotide sequence: (SEQ ID NO: 225)

tgatcaggagcccaatctctgacaaaactcacacatccccaccgtccccagcatccaacccgagaggggtgagcgctt
acctaagccggcccagcccggtcgacctgttcatccgcaagtcgccacgatcacctgtctggtggtggacctggcaccga
gcaaggggaccgtgaacctgacctggtcccgggccagtggggaagcctgtgaaccactccaccagaaaggaggagaag
cagcgcaatggcacgttaaccgtcacgtccaccctgccggtgggcacccgagactggatcgaggggggagacctaccag
tgcagggtgaccacccccacctgccagggccctcatgcggtccacgaccaagaccagcggcccgctgctgccccg
gaagtctatgcgtttgcg- acgccggagtg gccggggagccgggacaagcgcaccctcgctgctgatccagaac-
ttcatgcctgaggacatctcggtgcagtggtgcacaacgaggtgcagctcccgacgccgggcacagcacgacgcag
ccccgcaagaccaagggtccggcttcttctcttcagccgcctggaggtgaccagggccgaatgggagcagaaagatg
agttcatctgccgtgcagtcctgaggcagcgagcccctcacagaccgtccagcgagcgggtgtctgtaaatcccggtaaa
cgggatcccttcgaa

Amino acid sequence: (SEQ ID NO: 226)

dqepkssdkhtspasprgvsaylsrspfdlfrksptitclvdlapskgtnlwsrasgkpvnhstrkeekqr
ngltvtstlpvgtrdwiegetyqcrvthphlpralmrsttktsgraaapevyafatpewpgsrkrtlacliqnfmpe
vqwlhnevqlpdarhsttqprkktkgsgffvfrlevtraeweqkdeficravheaaspsqtvravsvnpgksgsfe

9. 2H7 VHL11S scFv hIgG1(SSS-S)H hIgE WCH3 WCH4

Nucleotide sequence: (SEQ ID NO: 227)

aagcttgccgccaatggattttcaagtcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctccagctctccagcaatcctgtctgcattctccaggggagaaggtcacaatgacttgcagggccagctcaagtgtaa
ttacatgcactgggtaccagcagaagccaggatctcccccaaacctggatttatgccccatccaacctggcttctggagtc
cctgtctgcttcagtggcagtggtgtctgggacctcttactctctcacaatcagcagagtggaggctgaagatgctgccactt
attactgccagcagtggaagttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggctcg
ggcggtgggtggaatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggccctgggg

cctcagtgaagatgtcctgcaaggcttctggctacacatttaccagttacaatatgactgggtaaagcagacacctagac-
 agggcctggaatggattggagctatttaccaggaaatggtgatacttctacaatcagaagttcaagggcaagggcacact
 gactgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgca
 agagtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcagga
 5 gccccaaatcttctgacaaaactcacacatccccaccgtcctcagcatccaacccgagagggggtgagcgcctacctaagcc
 ggcccagcccgttcgacctgttcacccgaagtcgcccacgatcacctgtctggtggtggacctggcaccagcaagggg
 accgtgaacctgacctgggtcccgggcccagtggaagcctgtgaaccactccaccagaaaggaggagaagcag
 cgcaatggcacgttaaccgtcacgtccaccctgccggtgggacccgagactggatcgagggggagacctaccagtgc
 aggggtgaccacccccacctgccagggccctcatgcggtccacgaccaagaccagcggccgcgtgctgccccgga
 10 agtctatcggttgcgacgccggagtggccggggagccgggacaagcgcaccctcgcctgcctgatccagaacttcatg
 cctgaggacatctcgggtcagtggtgcacaacgaggtgcagctcccgacgcccggcacagcacgacgcagccccg
 caagaccaagggtccggcttctcgtcttcagccgcctggaggtgaccagggccgaatgggagcagaagatgagttc
 atctgccgtgcagtcctatgaggcagcgagcccctcacagaccgtccagcgagcgggtgtctgtaaatcccggtaaatgata
 atctaga

15 Amino acid sequence: (SEQ ID NO: 228)

Mdfqvqifsfllisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyycqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
 pgasvkmsckasgytfsynmhvwkqtrpqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltسد
 savyfcarvvyysnsywyfdvwgtgtvtvssdqepkssdkthtspssasnprgvsaylsrpsfdlfrksptitclv
 20 vdlapskgtnltwsrasgkpvnhstrkeekqrmgtltvtstlpvgtrdwieetyqcervthphlpralmrsttktsgrpra
 pevyafatpewpgrdkrtlacliqnfmpeidvqwlhnevqlpdarhsttqprktksgffvfrlevtraeweqkde
 ficravheaaspsqtvqravsvnpgk

10. 2H7 VHL11S scFv hIgG1(SSS-P)H hIgE WCH3 WCH4

25 Nucleotide sequence: (SEQ ID NO: 229)

aagcttgccgcatggatttcaagtgcagattttagcttctgctaatactgcttcagtcataattgccagaggacaaattg
 ttctctcccagctctccagcaatctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagtgtaa
 ttacatgactggtaccagcagaagccaggatcctccccaaacccctggattatgccccatccaacctggcttctggagtc
 cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
 30 attactgccagcagtgaggatttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcggtggtctg
 ggcgggtggtgatctggaggagtgaggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaatgtcctgcaaggcttctggctacacatttaccagttacaatatgactgggtaaagcagacacctagaca
 gggcctggaatggattggagctatttaccaggaaatggtgatacttctacaatcagaagttcaagggcaagggcacactg
 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 35 gagtgggtgactatagtaacttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
 cccaaatcttctgacaaaactcacacatccccaccgtcccagcatccaacccgagaggggtgagcgcctacctaagccg
 gccagcccgttcgacctgttcacccgaagtcgcccacgatcacctgtctggtggtggacctggcaccagcaagggga
 ccgtgaacctgacctggtcccgggcccagtggaagcctgtgaaccactccaccagaaaggaggagaagcagcgcaatg
 gcacgttaaccgtcacgtccaccctgccggtgggacccgagactggatcgagggggagacctaccagtgcaggggtg
 40 acccaccacccacctgccagggccctcatgcggtccacgaccaagaccagcggcccgcgtgctgccccggaagtctat
 gcgtttgcgacgccggagtggccggggagccgggacaagcgcaccctcgcctgcctgatccagaacttcatgcctgag
 gacatctcgggtcagtggtgcacaacgaggtgcagctcccgacgcccggcacagcacgacgcagccccgcaagac
 caagggctccggcttctcgtcttcagccgcctggaggtgaccagggccgaatgggagcagaagatgagttcatctgcc
 gtgcagtcctatgaggcagcgagcccctcacagaccgtccagcgagcgggtgtctgtaaatcccggtaaatgataatctaga

45 Amino acid sequence: (SEQ ID NO: 230)

mdfvqvifsfllisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyycqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvrp

gasvkmsckasgytftsynmhwwkqtrpqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltse
avyfcarvvyysnsywyfdvwtgtttvssdqepkssdkhtsppspasprgvsaylsrpsfdlfrksptitclvv
dlapskgtvnltsrasgkpvnhstrkeekqrmgtltvtstlpvgtrdwiegetyqcrvthphlpralmrsttktsgraap
evyafatpewpgsrdrkrlacliqnfmpeidsvqwlhnevqlpdarhsttqprktksgffvfrlevtraeweqlkdefi
5 cravheaaspsqtvravsvnpgk

10. 2H7 VL L106S

Nucleotide sequence: (SEQ ID NO: 231)

10 aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaagggtcacaatgactgcagggccagctcaagtgttaag
ttacatgcactggtaccagcagaagccaggatcctccccaaaccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtcgc
ggcgggtggtgagctggaggaggtgggagctc

15 Amino acid sequence: (SEQ ID NO: 232)

mdfqvqifsfllisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiaypsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgatgkleskdggsgggsgggsgggss

11. 2H7 VL L106S scFv

20 Nucleotide sequence: (SEQ ID NO: 233)

aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaagggtcacaatgactgcagggccagctcaagtgttaag
ttacatgcactggtaccagcagaagccaggatcctccccaaaccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
25 attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtcgc
gagggtggtgagctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagctggtgaggcctggggc
ctcagtgaaagatgtcctgcaaggcttctggtacacattaccagttacaatatgactgggttaaagcagacacctagacag
ggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactga
ctgtagacaaatcctccagcagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaag
30 agtggtgtactatagtaacttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 234)

mdfqvqifsfllisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiaypsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgatgkleskdggsgggsgggsgggssqaylqqsgaelvr
pgasvkmsckasgytftsynmhwwkqtrpqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltse
35 savyfcarvvyysnsywyfdvwtgtttvssdq

12. 2H7 scFv VL L106S VHL11S scFv

Nucleotide sequence: (SEQ ID NO: 235)

40 Aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaatt
gttctctcccagctccagcaatcctgtctgcattccaggggagaagggtcacaatgactgcagggccagctcaagtgttaag
gttacatgcactggtaccagcagaagccaggatcctccccaaaccctggatttatgccccatccaacctggcttctggagtc
ccctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccact
tattactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtcgc

ggcgggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctg-
gggcctcagtgaaagtgctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctag
acagggcctggaatggattggagctatttatccaggaaatgggtgatacttctacaatcagaagttcaagggcaaggccac
actgactgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtg
5 caagagtgggtgactatagtaactcttactggcttctgatgtctggggcacagggaccacggtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 236)

mdfqvqifsflisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyccqwsfnpptfgagtkleskdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftslnmhvkwqtpqrglewigaiypngdtsynqkfkkgkatltvdksstaymqlsslteds
10 savyfcarvvyysnsyw yfdvwgtgtvtvssdq

10. Human IgD hinge linker with attached restriction sites

Nucleotide: (SEQ ID NO: 237)

gtggatccagggtcgaagtctccaaaggcacaggcctcctccgtgccactgcacaaccccaagcagaggggcagcctcg
15 ccaaggcaaccacagccccagccaccaccgtaacacaggaagaggaggagaagagaagaagaaggagaaggaga
aagaggaacaag aagagagagagacaaagaccgggtgcagtcgacg

Amino acid: (SEQ ID NO: 238)

vdpgskspkaqassvptaqpqaegslakattapattntgrggeekkkkekeeeqeeretktgavd

20 Sequence of Native IgD hinge domain:

(includes a cysteine residue-we truncated the hinge prior to that residue for these constructs:)

Nucleotide: (SEQ ID NO: 239)

gagtctccaaaggcacaggcctcctccgtgccactgcacaaccccaagcagaggggcagcctcgccaaggcaaccaca
25 gccccagccaccaccgtaacacaggaagaggaggagaagagaagaagaaggagaaggagaaagaggaacaagaa
gagagagagac aaagacaccagagtgtccgagcc- acaccagcctcttggcgtctacctgtaacccct

Amino acid sequence: (SEQ ID NO: 240)

Espkaqassvptaqpqaegslakattapattntgrggeekkkkekeeeqeeretktpecpshtqplgvyltp

30 12. 2H7 VHL11S

Nucleotide sequence: (SEQ ID NO: 241)

caggcttatctacagcagctctggggctgagtcggtgaggcctggggcctcagtgaaatgtcctgcaaggcttctggctac
acatttaccagttacaatatgactgggtaaagcagacacctagacaggcctggaatggattggagctatttatccaggaa
atgggtgatacttctacaatcagaagttcaagggcaaggccacactgactgtagacaaatcctccagcacagcctacatgc
35 agctcagcagcctgacatctgaagactctgcggtctatttctgtgcaagagtgggtgactatagtaactcttactggctactcga
tgtctggggcacagggaccacgg tcacctgtcttct

Amino acid sequence: (SEQ ID NO: 242)

qaylqqsgaesvrpgasvkmsckasgytftslnmhvkwqtpqrglewigaiypngdtsynqkfkkgkatltvdks
staym qlssltedsavyfcarvvyysnsyw yfdvwgtgtvtvss

40

13. 2H7 VH L11S scFv

Nucleotide sequence: (SEQ ID NO: 243)

aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
5 ttctctcccagctctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgttaag
ttacatgcaactggtagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggacctcttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgagtttaacccacccacgttcggtgctgggaccaagctggagctgaaagatggcggaggctcg
ggcgggtggtagcttgaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
cctcagtgaaatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
10 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtggtgtactatagtaactcttactggtacttcgatgtctggggcacaggggaccac ggtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 244)

mdfqvqifsfllisaviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
15 vparfsgsgsgtsysltisrveadaatyccqqsfnpptfagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftsynmhvwkqtrqglewigaiypngdtsynqkfkakatltvdkssstaymqlssltse
savycarvvyysnsywyfdvwgtgtvtvssdq

14. 2H7 scFv VH L11S hIgG1 (CSC-S)H WCH2 WCH3

20 Nucleotide sequence: (SEQ ID NO: 245)

aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgttaag
ttacatgcaactggtagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggacctcttactctctcacaatcagcagagtgagggtgaagatgctgccactt
25 attactgccagcagtgagtttaacccacccacgttcggtgctgggaccaagctggagctgaaagatggcggaggctcg
ggcgggtggtagcttgaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcgtgaggcctggggc
ctcagtgaaatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagacag
ggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactga
ctgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaag
30 agtggtgtactatagtaactcttactggtacttcgatgtctggggcacaggggaccacggtcaccgtctcttctgatcaggagc
ccaaatcttgtgacaaaactcacacatctccaccgtgctcagcacctgaactcctgggtggaccgtcagttctcttcccc
ccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaagacc
ctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtaca
acagcacgtaccgtgtggtcagcgtctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggtc
35 tccaacaaagccctccagcccccacgagaaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtac
acctgcccccatccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccaagcga
catcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacgg
ctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcat
gaggctctgcacaaccactacacgc agaagagcctctccctgtctccgggtaaatgatctaga

40 Amino acid sequence: (SEQ ID NO: 246)

mdfqvqifsfllisaviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftsynmhvwkqtrqglewigaiypngdtsynqkfkakatltvdkssstaymqlssltse
savycarvvyysnsywyfdvwgtgtvtvssdqepkscdkthtspcsapellggpsvflfppkpkdtlmsrtpevt
45 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiekti

skakgqprepqvylppsrdeitknqvslclvkgfypsdiavewesngqpennyktpvpvldsdgsfflyskltvdk
rwqqgnvfscsvmhhealnhytqkslspsgk

15. 2H7 scFv VH L11S IgE WCH2 WCH3 WCH4

5 Nucleotide sequence: (SEQ ID NO: 247)

aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgactggtaccagcagaagccaggtacctccccaaacctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctcacaatcagcagagtgagggtgaagatgctgccactt
10 attactgccagcagtgaggattttaacccaccacgttcgggtgctgggaccaagctggagctgaaagatggcgggtgctc
ggcgggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctgtgaggcctggggc
ctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagacag
ggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggaaggccacactga
ctgtagacaaatctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaag
15 agtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcacgtctg
ctccagggaacttaccgccaccgtgaagatcttacagtcgtcctgcgacggcggcgggcaacttcccccgaccatcc
agctcctgtgcctcgtctctgggtacacccagggaactatcaacatcacctgggtggaggacgggcaggtcatggacgtg
gacttgtccaccgcctctaccacgcaggaggggtgagctggcctccacaaaagcgagctcacctcagccagaagcact
ggctgtcagaccgcactacacctgccaggtcacctatcaagggtcacacctttgaggacagcaccaagaagtgtgcagatt
20 ccaacccgagaggggtgagcgcctacctaagccggccagcccgttcgacctgttcacccgaagtcgccacgatcac
ctgtctggtggtggacctggcaccagcaaggggaccgtgaacctgacctggctccggggcagtggaagcctgtgaac
cactccaccagaaaggaggagaagcagcgcaatggcacgttaaccgtcacgtccaccctgccggtgggcacccgagac
tggatcgagggggagacctaccagtcaggggtgacccaccccccacctgccagggccctcatgcggtccacgaccaag
accagcggcccgctgctgccccggaagtctatgcgtttgcgacgccggagtgccggggagccgggacaagcgcac
25 cctcgcctgcctgatccagaacttcacgtcctgaggacatctcgggtcagtggtgcacaacgaggtgcagctcccggacg
ccgggcacagcagcagcagccccgaagaccaagggctccggcttctctcagccgcctggaggtgaccaggg
ccgaatgggagcagaaagatgagttcatctgccgtgcagtcctatgaggcagcagccccctcacagacegtccagcgagc
ggtgtctgtaaatcccggttaagtataatctaga

Amino acid sequence: (SEQ ID NO: 248)

mdfvqvifsfllisaviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiaypsnlasg
vparfsgsgsgtsysltisrveadaatyycqqsfnpptfgagtkleldggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftsnnmhvkwqtrpglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltse
savyfcarvvyysnsywyfdvwtgtttvtvssdhvcsrdftptvkilqsscdggghfpptiqlclvsgytpgtinitwl
edgqvmvdvldstattqegelastqseltlsqkhwlsdrtycqvytqgthfedstkkcadsnprgvsaylsrspfdlfr
35 ksptitclvdlapskgtvnltsrasgkpvnhstrikeekqrmgtltvtstlpvgtrdwiegetyqcrvthphlpralmrst
tktsgpaaapevyafatpewpgsrdkrtlacliqnfmpeidsvqwlhnevqlpdarhsttqprtkgsgffvfrlevtra
eweekdeficravheaaspsqtvravsvnpgk

16. 2H7 scFv VH L11S mIgE WCH2 WCH3 WCH4 (SEQ ID NO: 249)

40 Nucleotide sequence:

aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgactggtaccagcagaagccaggtacctccccaaacctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctcacaatcagcagagtgagggtgaagatgctgccactt
45 attactgccagcagtgaggattttaacccaccacgttcgggtgctgggaccaagctggagctgaaagatggcgggtgctc
ggcgggtggtgatctggag- gaggtgggagc tctcaggcttatctacagcagctctggggctgagctgtgaggcctg-

gggcctcagtgaaatgtcctgcaaggcttctggctacacatttaccagttacaatatgcactgggtaaaagcagacacctag
acagggcctggaatggattggagctatttatccaggaaatgggatacttctacaatcagaagttcaagggcaaggccac
actgactgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggctatttctgtg
caagagtgggtgactatagtaactcttactggctacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcac
5 gttcgacctgtcaacatcactgagcccaccttgagctactccattcatcctgcgaccccaatgcattccactccaccatcca
gctgtactgcttcatttatggccacatcctaaatgatgtctctgtcagctgggtaaatggacgatcgggagataactgatacactt
gcacaaactgttctaatacaggaggaaggcaactagcctctacctgcagtaactcaacatcactgagcagcaatggatg
tctgaaagcaccttcacctgcaaggtcacctcccaaggcgtagactatttggcccacactcggagatgccagatcatgag
ccacgggggtgtgattacctacctgatccaccagccccctggacctgtatcaaaacgggtgtcccaagcttacctgtctgg
10 tgggtggacctggaaagcgagaagaatgtcaatgtgacgtggaaccaagagaagaagacttcagctcagcatccagtggtg
tacactaagcaccacaataacgccacaactagtatcacctccatcctgcctgtagttgccaaggactggattgaaggctacg
gctatcagtgcatagtggaccaccctgattttccaagcccattgtgcgttccatccaagaccccaggccagcgctcagc
ccccgaggtatatgtgtccaccaccagaggaggagagcggaggacaaacgcacactcacctgtttgatccagaacttctt
ccctgaggatattctgtgcagtggtgggggatggcaactgatctcaaacagccagcacagtaccacaacaccctga
15 aatccaatggctccaatcaaggcttcttcatcttcagtcgcctagaggtcgccaagacactctggacacagagaaaacagtt
cacctgccaagtgatccatgaggcacttcagaaaccaggaactggagaaaacaatatccacaagccttggtaacctt
ccctccgtccatcctagtaacttagag

Amino acid sequence (SEQ ID NO: 250)

mdfqvqifsfllisaviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
20 vparfsgsgsytisrveadaatyycqqsfnpptfgagtkleldggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytfsynmhvkwqtrpglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltsgd
savycarvvyysnsywyfdvwtgtttvssdhvrpniteptlellhsscdpnafhstiqlycfyghilndvsvswl
mddreitdtlaqtvlikeegklastcsklniteqqwmsestfickvtsqgvdylahtrcpdheprgvitylippspldlyq
ngapklclvdlseknvntwnqekktvsasqwytkhhnnattsitsilpvvakdwiegygyqivdhpdpkp
25 ivrsitktpgqrsapevyvfppeeesedkrtltclqnffpedisvqwlgdgklisnshsttplksngsnqgffifsrle
vaktlwtqrkqftcqihealqkprklektistslgntslrps

17. 2H7 scFv VH L11S hIgA WH WCH2 T4CH3 (SEQ ID NO: 251)

Nucleotide sequence:

30 aagcttggccgcatggatttcaagtgacagatttcagcttctgtaatacagtgcttcagtcataattgccagaggacaaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaagggtcacaatgacttcagggccagctcaagtgtaa
ttacatgcactggtagcagagaagccaggatcctcccccacccctggatttatcccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgaggatttaacccaccacgcttcggtgctgggaccaagctggagctgaaagatggcgggtggtcgc
35 ggcggtgggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctctgtgaggcctggggc
ctcagtgaaatgtcctgcaaggcttctggctacacatttaccagttacaatatgcactgggtaaaagcagacacctagacag
ggcctggaatggattggagctatttatccaggaaatgggtgatacttctacaatcagaagttcaagggcaaggccacactga
ctgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggctatttctgtgcaag
agtgggtgactatagtaactcttactggctacttcgatgtctggggcacaggaccacggtcaccgtctcttctgatcagccagt
40 tcctcaactccactacccatctccctcaactccactacccatctccctcatgctgccacccccgactgtcactgcacc
gaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcaggt
gtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccacctgacctgtggtgctacagc
gtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttacttcagctgctgctaccccgagtc
aagaccccgctaaccgccacctctcaaaatccggaaacacattccggcccagggtccacctgctgccgccgctcgg
45 aggagctggccctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagccccaaggatgtgctggttcgctgg
ctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccggcaggagcccagccagggcaccac
caccttcgctgtgaccagcactatgcgcgtggcagccaggactggaagaagggggacaccttctcctgcatggtgggc

cacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaaccacccatgtcaatgtgtctgtt
gtcatggcggaggtggactgataatctaga

Amino acid sequence: (SEQ ID NO: 252)

mdfqvqifsflisasviiargqivlsqpailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
5 vparfsgsgsgtsysltisrveadaatyycqqsfnppftgagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftsynmhvwkqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltsgd
savyfcarvvyysnsywyfdvwgtgtvtvssdqpvpstptpsptpspscchprlsrlhrpaledlllgseailtcl
tglrdasgvtftwtpssgksavqgppdrdlcgcysvssvlpgcaepwnhgktfctaaypesktpatlslksngntfrpev
hllpppseelalnelvtlclargfspkdvlrwlqgsqelprekyltwasrqepsqgtttfavtsilrvaaedwkkgdtfsc
10 mvghalplaftqktidrlagkpthvnsvvmaevd

18. 2H7 scFv VH L11S mIgA WH WCH2 T4 CH3

Nucleotide sequence: (SEQ ID NO: 253)

aagcttgccgccatggatttcaagtgcagatttcagcttctgtaatcagtgcttcagtcataattgccagaggacaaattg
15 ttctctcccagctccagcaatcctgtctgcattccaggggagaagggtcacaatgacttcagggccagctcaagtgtaa
ttacatgcactggtagcagcagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtgctcg
ggcgggtggtgatctggaggagtgaggagctctcaggcttatctacagcagctctggggctgagctgtgaggcctggggc
20 ctacgtgaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagacag
ggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggcaagccacactga
ctgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaag
agtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcacatctgt
tctctctactactcctcctccacctcctgcccagccagcctgtcactgcagcggccagctcttgaggacctgctcctggg
25 ttcatagccagcatcacatgtactctgaatggcctgagagatctgagggagctgtcttcacctgggagccctccactggg
aaggatgcagtgacagaagaaagctgtgcagaattcctgcggctgtacagtgtgtccagcgtctgcctggctgtgctgag
cgctggaacagtgccgcacattcaagtgcacagttacceatcctgagctctgacaccttaactggcacaattgccaaagta
cagtgaacaccttcccacccaggtccacctgctaccgccgccgctcgaggagctggccctgaatgagctcgtgtccctg
acatgcctggtgcgagcttcaaccctaaagaagtgtggtgcgatggctgcattggaaatgaggagctgtccccagaaag
30 ctacctagtgtttgagcccctaaaggagccaggcgaggagccaccctacctggtgacaagcgtgttgctgtatcag
ctgaaatctgaaacagggtgaccagtactcctgcattggtgggccacgaggccttgccatgaacttcaccagaagacc
atcgaccgtctgtcgggtaaacccaccaatgtcagcgtgtctgtgatcatgtcagaggagattgataatctagat

Amino acid sequence: (SEQ ID NO: 254)

mdfqvqifsflisasviiargqivlsqpailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
35 vparfsgsgsgtsysltisrveadaatyycqqsfnppftgagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftsynmhvwkqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltsgd
savyfcarvvyysnsywyfdvwgtgtvtvssdhicspptppppscqpslsrlhrpaledlllgdsaitctlnlrdpeg
avftwepstgkdavqkkavqnsqgcysvssvlpgcaerwnsgasfkctvthpesdltgtiakvtvntfppqvhlppp
seelalnelvslclvrafnpkvllrvlhgneelspesylvfeplkepgegattylvtslrvsaeiwwkqgdqyscmvg
40 healpmnftqktidrlsgkptnsvsvimsegd

A. mIgA WCH2 T4CH3

Nucleotide sequence: (SEQ ID NO: 255)

Gttgttgatcacatctgttctcctactactcctcctccacctcctgccagccagcctgtcactgcagcggccagctcttg
aggacctgctcctgggttcagatgccagcatcacatgtactctgaatggcctgagagatcctgaggagctgtcttcacctg
45 ggagccctccactgggaaggatgcagtgacagaagaaagctgtgcagaattcctgcggctgctacagtgtgtccagcgtcc

5 tgcctggctgtgctgagcgtggaacagtggcgcatcattcaagtgcacagttacccatcctgagtctgacaccttaactgg
 cacaattgccaaagtcacagtgaacaccttcccaccccagggtccacctgctaccgccgccgtcggaggagctggccctga
 atgagctcgtgtccctgacatgctggtgcgagctttcaaccctaaagaagtgtggtgcgatggctgcatggaaatgagg
 agctgtccccagaaagctacctaagtgtttgagcccctaaaggagccaggcgagggagccaccacctacctggtgacaag
 cgtgttgcgtgtatcagctgaaatctggaacagggtgaccagtactcctgcatggtggggccacgagggccttgcccatgaa
 cttcaccagaagaccatcgaccgtctgtcgggtaaacccaccaatgtcagcgtgtctgtgatcatgtcagaggagattga
 taatctagat

Amino acid sequence: (SEQ ID NO: 256)

10 dhicspptppppscqpslslrpaledlllgsdasitctlnlrdpegavftwepstgkdavqkkavqnsccysvssvl
 pgcaerwnsgasfkctvthpesdtltgtiaktvntfppqvhllpppseelalnelvsltlvrafnpkvllvrwlhgneel
 spesylvfeplkepeggattylvtsvlrsvaeiwwkqgdqyscmvghealpmnftqktidrlsgkptnsvsvimsegd

20. K322S CH2 region

Nucleotide sequence: (SEQ ID NO: 257)

15 cctgaactcctggggggaccgtcagtcttctcttcccccaaaaaccaaggacaccctcatgatctcccgaccctgag
 gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
 ataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
 ggactggctgaatggcaaggagtacaagtgtcgtctccaacaaagccctcccagccccatcgagaaaacaatctcca
 aagccaaa

20 Amino acid sequence: (SEQ ID NO: 258)

pellggpsvflfppkpkdtlmsirtpvctvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvl
 hqdwlnkeykcsvsnkalpapietiskak

21. K322S CH2 WCH3

25 Nucleotide sequence: (SEQ ID NO: 259)

cctgaactcctggggggaccgtcagtcttctcttcccccaaaaaccaaggacaccctcatgatctcccgaccctgag
 gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
 ataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
 ggactggctgaatggcaaggagtacaagtgtcgtctccaacaaagccctcccagccccatcgagaaaacaatctcca
 30 aagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgaccaagaaccaggtca
 gcctgacctgcctggtcaaaggcttctatccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaa
 ctacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtg
 gcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctcctgtc
 tccgggtaaatgatctaga

35 Amino acid sequence: (SEQ ID NO: 260)

Pellggpsvflfppkpkdtlmsirtpvctvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvl
 hqdwlnkeykcsvsnkalpapietiskakgqprepqvylppsdeltknqvsltlvkgyfypsdiavewesngqp
 ennyktppvl dsdgsfflyskltvdkrsrwqqgnvfscsvmhealnhytqkslsispk

40 1. K322L CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 261)

tgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagt
cttctcttccccccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgctggtggtggacgtgag
ccacgaagaccctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcggga
ggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtac
5 aagtgcctggtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaac
cacaggtgtacaccctgccccatccccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttc
tatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctg
gactccgacggctccttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgc
tccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaatgatctaga

10 Amino acid sequence: (SEQ ID NO: 262)

Dqepkssdkthtspssapellggpsvflfppkpkdtlmsirtpevtcvvvdvshedpevkfnwyvdgvevhnakt
kpreeqynstyrvsvltvlhqdwlngkeykclvsnkalspapietkiskakgpprepqvytlppsrdeitknqvsitclv
kgfypsdiavewesngqpennyktpvldsdgsfflyskltvdksrwqqgnvfscsvmhhealhnhytqkslsisp
k

15

22. 2H7 scFv VHL11S hIgG1 (SSS-S)H K322SCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 263)

aagcttgccgccatggatttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacatgacttcagggccagctcaagtgaag
20 ttacatgactggtaccagcagaagccaggtacctccccaaaccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtggaggtgaagatgctgccactt
attactgccagcagtggttttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtcgc
ggcgggtggtggtatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagaca
25 gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactg
actgtagacaaatctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagttcttcttccc
cccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgctggtggtggacgtgagccacgaagac
30 cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagaca
acagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgtcggtc
tccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtac
acctgccccatccccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcga
catcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacgg
35 ctctcttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcat
gaggctctgcacaaccactacacg agaagagcctctcctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 264)

mdfqvqifsflisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiyapsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfagtklelkdggsgggsgggssqaylqqsgaesvr
40 pgasvkmsckasgytftsynmhvkwqtrqglewigaiypngdtsynqkfkgkatltvdksstaymqslstsed
savyfcarvvyysnsywyfdvwtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmsirtpevt
cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykcsvsnkalspapietki
skakgpprepqvytlppsrdeitknqvsitclvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdks
rwqqgnvfscsvmhhealhnhytqkslsispk

45

23. 2H7 scFv VHL11S hIgG1 (SSS-S)H K322LCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 265)

aagcttgccgcatggatttcaagtgcagatttccagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtagcagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
5 cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgcttcggtgctgggaccaagctggagctgaaagatggcgggtgctcg
ggcgggtggtggtatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
10 actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtctcagcacctgaactcctggggggaccgtcagcttctcttccc
cccaaaacccaaggacacccctcatgatctcccgaccctgaggtcacatgcgtggtggtgagctgagccacgaagac
cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgaggagagcagtagc
15 aacagcacgtaccgtgtggtcagcgtctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgcctggt
ctccaacaaagccctccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
caccctgccccatcccggtgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccacgcg
acatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcecggtgctggactccgacg
gtcctcttctctacagcaagctaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctcgtgatgc
20 atgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 266)

mdfqvqifslhisasviiargqivlsqspailsaspgkvmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnppfagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytfsynmhwwkqtrpglewigaiypngdtsynqkfkgkatltvdkssstaymqlsslteds
25 savyfcarvvyysnsywyfdvwgtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmisrtpet
cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkēykcclvsnkalpapiēkti
skakgqprepvytlppsrdeltnqvsltlvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdk
rwqqgnvfscsvmhēalhnhytqkslsispk

30 24. 2H7 scFv VHL11S hIgG1 (CSS-S)H K322SCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 267)

aagcttgccgcatggatttcaagtgcagatttccagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttaeatgcactggtagcagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
35 cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgcttcggtgctgggaccaagctggagctgaaagatggcgggtgctcg
ggcgggtggtggtatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
40 actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaatcttgtgacaaaactcacacatccccaccgtctcagcacctgaactcctggggggaccgtcagcttctcttccc
cccaaaacccaaggacacccctcatgatctcccgaccctgaggtcacatgcgtggtggtgagctgagccacgaagac
cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgaggagagcagtagc
45 aacagcacgtaccgtgtggtcagcgtctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgcctcgt
ctccaacaaagccctccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
caccctgccccatcccggtgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccacgcg

acatcgccgtggagtgaggagcaatgggcagccggagacaactacaagaccacgcctcccgtgctggactccgacg
gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
atgaggctctgcacaaccactacacgc agaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 268)

- 5 mdfqvqifsfhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftslnmhvkwqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltse
savyfcarvvyysnsywyfdvwtgtvtvssdqepkscdkthtspssapellggpsvflfppkpkdtlmsrtpevt
cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykcsvsnkalpapieti
10 skakgqprepvytlppsrdeitknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdks
rwqqgnvfscsvmhealhnhytqkslslspgk

25. P331S CH2

Nucleotide sequence: (SEQ ID NO: 269)

- 15 cctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacaccctcatgatctcccgaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagtcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcctccatcgagaaaacaatctcca
aagccaaa

- 20 Amino acid sequence (SEQ ID NO: 270)

pellggpsvflfppkpkdtlmsrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvl
hqdwlngkeykckvsnkalpasiektiskak

26. P331S CH2 WCH3

- 25 Nucleotide sequence: (SEQ ID NO: 271)

- cctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacaccctcatgatctcccgaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagtcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcctccatcgagaaaacaatctcca
30 aagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccggtatgagctgaccaagaaccaggtca
gcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaa
ctacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtg
gcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgc
tccgggtaaatgatctaga

- 35 Amino acid sequence (SEQ ID NO: 272)

pellggpsvflfppkpkdtlmsrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvl
hqdwlngkeykckvsnkalpasiektiskakgqprepvytlppsrdeitknqvsltlvkgfypsdiavewesngqp
ennyktpvldsdgsfflyskltvdksrwqqgnvfscsvmhealhnhytqkslslspgk

- 40 27. 2H7 scFv VH L11S (SSS-S)H P331S CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 273)

aagcttgccgcatggattttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
 ttacatgactggtagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
 cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
 5 attactgccagcagtgaggatttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtc
 ggcgggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagaca
 gggcctggaatggattggagctattatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
 actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 10 gactgactatagtaactcttactggtacttcgatgtctggggcacagggaaccacggtcaccgtctcttctgatcaggagccca
 aatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcttccccca
 aaaccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtgacgtgagccacgaagaccctg
 aggtcaagttcaactgtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtagaca
 gcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtcaaggtctcc
 15 aacaaagccctcccagcctccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacc
 ctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacat
 cgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctc
 ctctctctctacagcaagctcaccgtggacaagagcaggtggcagcagggaacgtcttctcatgctccgtgatcatga
 ggctctgcacaaccactacacgc agaagagcctctccctgtctccgggtaaatgatctaga

20 Amino acid sequence (SEQ ID NO: 274)

mdffqvqifslhisasviiargqivlsqspailsaspgkvmtcrassvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsytstisrveadaatyccqwsfnppftgagtkleldggsgggsgggssqaylqqgsaevr
 pgasvkmsckasgytfsynmhvkwqtrpglewigaiypngdtsynqkfkgtatlvdkssstaymqslstsd
 savyfcarvvyysnywyfdvwgtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmisrtpevt
 25 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeyckkvsnkpalpasiekti
 skakgqprepvytlppsrdeltnqvslclvkgyfypsdiavewesngqpennkytppvldsdgsfflyskltvdk
 rwqqgnvfscsymhealhnhytqkslslspgk

28. 2H7 scFv VH L11S (CSS-S)H P331S CH2 WCH3

30 Nucleotide sequence: (SEQ ID NO: 275)

aagcttgccgcatggattttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
 ttacatgactggtagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
 cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
 35 attactgccagcagtgaggatttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggtc
 ggcgggtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagaca
 gggcctggaatggattggagctattatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
 actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 40 gactggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaaccacggtcaccgtctcttctgatcaggag
 cccaaatcttgtgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcttccc
 cccaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtgacgtgagccacgaagac
 cctgaggtcaagttcaactgtgtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtag
 aacagcagctaccgtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtcaaggt
 45 ctccaacaaagccctcccagcctccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtac
 acctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcga
 catcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacgg

ctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcat
gaggctctgcacaaccactacacgc agaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence (SEQ ID NO: 276)

5 mdfqvqifslhisasviiargqivlsqspailsaspgkvtmtcrasssvsymhwyqqkpgsspkpwiapsnlasg
vparfsgsgtsysltisrveadaatyccqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftslnmhvkwqtrqglewigaiypngdtsynqtkgkatltvdksstaymqlssltse
savycarvvysnsywyfdvwtgtvtvssdqepkscdkthtspssapellggpsvflfppkpkdtlmisrtpevt
cuvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpasiecti
10 skakgqprepvytlppsrdeletknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdks
rwqqgnvfscsvmhlehnhytqkslslspgk

29. T256N CH2 region

Nucleotide sequence: (SEQ ID NO: 277)

15 Cctgaactcctggggggaccgtcagttcttcttccccccaaaacccaaggacaccctcatgatctcccgaaccctgag
gtcacatgcgtgggtggacgtgagccacgaagaccctgaggtcaagtcaactggtagcggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtgtgcagcgtcctcaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctcc
aaagccaaa

Amino acid sequence (SEQ ID NO: 278)

20 pellggpsvflfppkpkdtlmisrnpvctuvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltv
lhqdw lngkeykckvsnkalpapietiskak

30. T256N CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 279)

25 cctgaactcctggggggaccgtcagttcttcttccccccaaaacccaaggacaccctcatgatctcccgaaccctgag
gtcacatgcgtgggtggacgtgagccacgaagaccctgaggtcaagtcaactggtagcggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtgtgcagcgtcctcaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctcc
aaagccaaaggcagccccgagaaccacaggtgtacacctgcccccatccgggatgagctgaccaagaaccagggtc
30 agcctgacctgcctggtcaaaggcttctatcccagcgacatgccgtggagtgaggagcaatgggcagccggagaaca
actacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggt
ggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgt
ctccgggtaaatgatctaga

Amino acid sequence (SEQ ID NO: 280)

35 pellggpsvflfppkpkdtlmisrnpvctuvvdvshedpevkfhwyvdgvevhnaktkpreeqynstyrvsvltv
lhqdwlngkeykckvsnkalpapietiskakgqprepvytlppsrdeletknqvsltlvkgfypsdiavewesngq
pennyktpv ldsdgsfflyskltvdksrwqqgnvfscsvmhlehnhytqkslslspgk

31. 2H7 scFv VH L11S (SSS-S)H T256N CH2 WCH3

40 Nucleotide sequence: (SEQ ID NO: 281)

aagcttgcggccatggatttcaagtgcagattttagcttctgtaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaagggtcacaatgacttcagggccagctcaagtgtaa

ttacatgactggtaccagcagaagccaggatcctcccccaccctggatttatgccccatccaacctggcttctggagtc
 cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtggaggctgaagatgctgccactt
 attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggtggtcgc
 5 ggcgggtggtggtatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagtcaagggaaggccacactg
 actgtagacaaaatctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgtatcaggag
 cccaaatcttctgacaaaactcacacatccccaccgtctcagcacctgaactcctggggggaccgtcagcttctcttccc
 10 cccaaaacccaaggacacccctcatgatctcccgaaccctgaggtcacatgcgtgggtgggacgtgagccacgaagac
 cctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
 aacagcacgtaccgtgtggtcagcgtctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaggttctatcccagcg
 15 acatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
 gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacagcagaagagcctctcctgtctccgggtaaatgatctaga

Amino acid sequence (SEQ ID NO: 282)

mdfqvqifslhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 20 vparfsgsgsytisrveadaatyccqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
 pgasvkmsckasgytfsynmhvkwqtrpqrlewigaiypngdtsynqkfkghkathvdkssstaymqlssltse
 savyfcarvvyysnywyfdvwtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmismpevt
 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
 25 skakgqprepqvylppsrdeltknqvsltclvkgyfypsdiavewesngqpennyktpvpldsdgsfflyskltvdk
 rwqqgnvfscsvmhcalhnhytqkslslspgk

32. 2H7 scFv VH L11S (CSS-S)H T256N CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 283)

aagcttgccgccatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
 30 ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
 ttacatgactggtaccagcagaagccaggatcctcccccaccctggatttatgccccatccaacctggcttctggagtc
 cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtggaggctgaagatgctgccactt
 attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggtggtcgc
 35 ggcgggtggtggtatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagtcaagggaaggccacactg
 actgtagacaaaatctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
 gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgtatcaggag
 40 cccaaatcttctgacaaaactcacacatccccaccgtctcagcacctgaactcctggggggaccgtcagcttctcttccc
 cccaaaacccaaggacacccctcatgatctcccgaaccctgaggtcacatgcgtgggtgggacgtgagccacgaagac
 cctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
 aacagcacgtaccgtgtggtcagcgtctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaggttctatcccagcg
 45 acatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
 gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacagcagaagagcctctcctgtctccgggtaaatgatctaga

Amino acid sequence (SEQ ID NO: 284)

mdfqvqifslhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiyapsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftsynmhvkwqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltsed
5 savyfcarvvyysnsywyfdvwgtgtvtvssdqepkscdkthtsppssapellggpsvflfppkpkdtlmismpev
tcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpapiekt
iskakgqprepvytlppsrdeitknqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdk
rwqqgnvfscsvmhleahnhytqkslslspgk

10 33. RTPE/QNAK (255-258) CH2

Nucleotide sequence: (SEQ ID NO: 285)

cctgaactcctggggggaccgtcagtccttcttcccccaaaacccaaggacaccctcatgatctcccagaacgctaag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagtcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
15 ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctcc
aaagccaaa

Amino acid sequence (SEQ ID NO: 286)

pellggpsvflfppkpkdtlmisqnakvtcvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvlt
vlhqdwlngkeykckvsnkalpapiektiskak

20

34. RTPE/QNAK (255-258)CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 287)

cctgaactcctggggggaccgtcagtccttcttcccccaaaacccaaggacaccctcatgatctcccagaacgctaag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagtcaactggtacgtggacggcgtggaggtgc
25 ataatgccaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctcc
aaagccaaaaggcagccccgagaaccacaggtgtacaccctgcccccatccgggatgagctgaccaagaaccagggtc
agcctgacctgctgtcaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatgggcagccggagaaca
actacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggt
30 ggagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgt
ctccgggtaaatgatctaga

Amino acid sequence (SEQ ID NO: 288)

pellggpsvflfppkpkdtlmisqnakvtcvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvlt
vlhqdwlngkeykckvsnkalpapiektiskakgqprepvytlppsrdeitknqvsltlvkgfypsdiavewesng
35 qpennykttppvldsdgsfflysklvdkswqqgnvfscsvmhleahnhytqkslslspgk

35. 2H7 scFv VH L11S (SSS-S)H RTPE/QNAK (255-258)CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 289)

aagcttgccgcatggattttcaagtcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
40 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
ttacatgactggtaccagcagaagccaggatcctccccaaacccctggattatgccccatccaacctggcttctggagtc
cctgctgcttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggctgaagatgctgccactt
attactgccagcagtgaggtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg

ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtagggcctgggg
 cctcagtgaagatgtcctgcaaggctcttgctacacattaccagttacaatatgactgggtaaagcagacacctagaca
 gggcctgggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactg
 actgtagacaaaatctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggcttatttctgtgcaa
 5 gagggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacgggtcaccgtctcttctgatcaggag
 cccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagctcttcttccc
 cccaaaacccaaggacacccctcatgatctcccagaacgctaaggtcacatgcgtgggtgggacgtgagccacgaagac
 cctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
 aacagcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaagggt
 10 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctggtcaaaggcttctatcccagcg
 acatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgtgactccgacg
 gtccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacacg cagaagagcctctccctgtctccgggtaaatgatctaga

15 Amino acid sequence (SEQ ID NO: 290)

mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrassvsymhwyqqkpgsspkpwiaypsnlasg
 vparfsgsgsgtsysltisrveaadaatyycqqsfnpptfagtkleldggsgsgsgsgsgssqaylqqsgaesvr
 pgasvkmsckasgytfsynmhvwkqtrpqglewigaiypngdtsynqtkgkatltvdkssstaymqlssltsed
 20 savyfcarvvyysnsywyfdvwgtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmlsqnakv
 tcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpapiekt
 iskakgpprepqvytlppsrdeitknqvsitclvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdk
 rwqqgnvfscsvmhealhnhytqkslsispk

36. 2H7 scFv VH L11S (CSS-S)H RTPE/QNAK (255-258)CH2 WCH3

25 Nucleotide sequence: (SEQ ID NO: 291)

aagcttgcggccatggacaagtgcagatttccagcttctgtaatacagtgcttcagtcataattgccagaggacaaattgttct
 ctcccagctctccagcaatcctgtctgcattctccaggggagaaggtcacatgacttgcagggccagctcaagtgttaattac
 atgcactggtagcagcagaagccaggatcctccccaaaccctggatttatgccccatccaaacctggcttctggagtcctg
 ctgccttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccacttattac
 30 tgccagcagtgaggatttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcgggcg
 gtggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctggggcctca
 gtgaagatgtctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagacagggcc
 tggatggattggagctatttatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactgactgta
 gacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaagagtgg
 35 tgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacgggtcaccgtctcttctgatcaggagccaaat
 ctltgtgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagcttcttcttccccccaaa
 acccaaggacacccctcatgatctcccagaacgctaaggtcacatgcgtgggtgggacgtgagccacgaagaccctgag
 gtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtagaacagc
 acgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaa
 40 caaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacct
 gccccatcccgggatgagctgaccaagaaccaggtcagcctgacctggtcaaaggcttctatcccagcgacatcg
 ccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgtgactccgacggctcct
 tcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcagagg
 ctctgcacaaccactacacg cagaagagcctctccctgtctccgggtaaatgatctaga

45 Amino acid sequence (SEQ ID NO: 292)

mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrassvsymhwyqqkpgsspkpwiaypsnlasg
 vparfsgsgsgtsysltisrveaadaatyycqqsfnpptfagtkleldggsgsgsgsgsgssqaylqqsgaesvr

pgasvkmsckasgytftsynmhvwkqtrpqglewigaiypngdtsynqkfkgtatltvdkssstaymqlsslt
 savyfcarvvyysnywyfdvwtgtvtvssdqepkscdkthtspssapellggpsvflfppkpkdtl
 misqnakv
 tcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalp
 iekt
 iskakgqpprepqvytlppsrdeltnqvslclvkgyfypsdiavewesngqpennykttpvldsdgsfflyskltvdks
 5 rwqqgnvfscsvmhealhnhytqkslsispk

36. K290Q CH2 region

Nucleotide sequence: (SEQ ID NO: 293)

10 cctgaactcctggggggaccgtcagtccttcttcccccaaaacccaaggacaccctcatgatctcccggaccctgag
 gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaactggtacgtggacggcgtggaggtgc
 ataatccaagacacagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
 ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacaatctcc
 aaagccaaa

Amino acid sequence: (SEQ ID NO: 294)

15 pellggpsvflfppkpkdtl misrtpevtcvvvvdvshedpevkfnwyvdgvevhnaktqpreeqynstyrvvsvltvl
 hqdwlngkeykckvsnkalp iektiskak

37. K290Q CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 295)

20 cctgaactcctggggggaccgtcagtccttcttcccccaaaacccaaggacaccctcatgatctcccggaccctgag
 gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaactggtacgtggacggcgtggaggtgc
 ataatccaagacacagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
 ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacaatctcc
 aaagccaaaaggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgaccaagaaccaggtc
 25 agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagaaca
 actacaagaccagcctcccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggt
 ggcagcaggggaactcttctcatgctcctgatgcatgaggtctgcacaaccactacacgcagaagagcctctccctgt
 ctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 296)

30 pellggpsvflfppkpkdtl misrtpevtcvvvvdvshedpevkffwyvdgvevhnaktqpreeqynstyrvvsvltvl
 lhqdwlngkeykckvsnkalp iektiskakgqpprepqvytlppsrdeltnqvslclvkgyfypsdiavewesngq
 pennykttpvldsdgsfflyskltvdksrwqqgnvfscsvmhealhnhytqkslsispk

38. 2H7 scFv VH L11S (SSS-S)H K290Q CH2 WCH3

35 Nucleotide sequence: (SEQ ID NO: 297)

aagcttgccgccatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggggcagctcaagtgtaa
 ttacatgcactggtaccagcagaagccaggatcctccccaaaccctggattatgccccatccaacctggcttctggagtc
 cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
 40 attactgccagcagtgaggtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg
 ggcggtggtgagctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagtcggtgaggcctgggg
 cctcagtgaaatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagac
 agggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaaggtcaagggaaggccacact

gactgtagacaaatctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcgggtctatttctgtgca
 agagtgggtactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctctctgatcagga
 gccccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagttctcttcc
 ccccaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaaga
 5 ccttgagggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacacagccgcgggaggagcagta
 caacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaag
 gtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgt
 acaccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagc
 gacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgac
 10 ggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatg
 catgaggctctgcacaaccactacag cagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 298)

mdfqvqifsfhlhisasviiargqivlsqpailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgatkleldggsgggsgggssqaylqqsgaesvr
 15 pgasvkmsckasgytftsymbhwwkqtrqglewigaiypngdtsynqkflgkatltvdkssstaymqlsslted
 savyfcarvvyysnsywyfdvwtgtvtvssdqepkssdkthtspssapellggspsvflfppkpkdtlmisrtpevt
 cvvvdvshedpevkfnwyvdgvevhnaktqpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiiekti
 skakgqprepqvtylppsrdeitknqvslctlvkgfypsdiavewesngqpennkyktpvldsdgsfflyskltvdk
 rwqqgnvfscsvmhhealhnhytqkslslspgk

20

39. 2H7 scfv VH L11S (CSS-S)H K290Q CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 299)

aagcttgccgccatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttcgagggccagctcaagtgaag
 25 ttacatgcactggtaccagcagaagccaggatcctcccccaccctggattatgccccatccaacctggcttctggagtc
 cctgctcgttcagtgagcagtggtctgggacctcttactctcacaatcagcagagtggaggctgaagatgctgccactt
 attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtgctcg
 ggcggtggtgagctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaagatgctctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 30 gggcctggaatggattggagctattatccaggaaatggtgatacttctcacaatcagaagttcaagggcaaggccacactg
 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcgggtctatttctgtgcaa
 gagtgggtgactatagtaactcttactggtacttcgatgtctggggcacaggaccacggtcaccgtctctctgatcaggag
 cccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagttctcttctcc
 cccaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagac
 35 cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacacagccgcgggaggagcagtac
 aacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggt
 ctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcg
 acatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
 40 gctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgeacaaccactacag cagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 300)

mdfqvqifsfhlhisasviiargqivlsqpailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgatkleldggsgggsgggssqaylqqsgaesvr
 45 pgasvkmsckasgytftsymbhwwkqtrqglewigaiypngdtsynqkfkkgkathvdkssstaymqlsslted
 savyfcarvvyysnsywyfdvwtgtvtvssdqepkscdkthtspssapellggspsvflfppkpkdtlmisrtpevt
 cvvvdvshedpevkfnwyvdgvevhnaktqpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiiekti

skakgpprepqvylppsrdeltnqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdks
rwqqgnvfscsvmhcalhnhytqkslsispk

40. A339PCH2

5 Nucleotide sequence: (SEQ ID NO: 301)

cctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacaccctcatgatctccggaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacaatctcc
10 aaacccaaa

Amino acid sequence: (SEQ ID NO: 302)

pellggpsvflfppkpkdtlmsrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvl
hqdwlnkeykckvsnkalpapiektiskpk

15 41. A339P CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 303)

cctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacaccctcatgatctccggaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctaccgtcctgcacca
20 ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacaatctcc
aaacccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgaccaagaaccaggtc
agcctgacctgctgtgtaaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatgggcagccggagaaca
actacaagaccacgctccctgctggtgactccgacggctcttcttctctacagcaagctcaccgtggacaagagcaggt
ggcagcaggggaactgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgt
25 ctccgggtaatgatctaga

Amino acid sequence: (SEQ ID NO: 304)

pellggpsvflfppkpkdtlmsrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvl
hqdwlnkeykckvsnkalpapiektiskpkgpprepqvylppsrdeltnqvsltlvkgfypsdiavewesngci
pennykttppvldsdgsfflyskltvdksrwqqgnvfscsvmhcalhnhytqkslsispk

30

42. 2H7 scFv VHL11S (SSS-S)H A339P CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 305)

aagcttgccgcatggatttcaagtgcagatttcagttcctgctaactcagtgcttcagtcataattgccagaggacaaattg
ttctctcccagttccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgaggccagctcaagtgaag
35 ttacatgcactggtaccagcagaagccaggtacctccccaaacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgaggttttaacccaccacgttcgggtgctgggaccaagctggagctgaaagatggcgggtggctcg
ggcgggtggtggtatctggaggaggtgggagctctcagggttatctacagcagctctggggctgagtcggtgaggcctgggg
cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
40 gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
actgtagacaaaactccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagttcttcttctcc

cccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtgacgtgagccacgaagac
 cctgaggtcaagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
 aacagcacgtaccgtgtggtcagcgtcctaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaacccaaagggcagccccgagaaccacaggtgta
 5 caccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaggcttctatcccagcg
 acatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
 gtccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacacgc agaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 306)

10 mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyccqqsfnpptfagtklelkdgggsgggsgggssqaylqqsgaesvr
 pgasvkmsckasgytftslnmhvkwqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltسد
 savyfcarvvyysnsywyfdvwgtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmsrtpevt
 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
 15 skpkgqpprepqvylppsrdeitknqvslclvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdk
 rwqqgnvfscsvmhhealhnhytqkslslspgk

43. 2H7 scFv VHL11S (CSS-S)H A339P CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 307)

20 aagcttgccgcatggtatttcaagtgcagatttccagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatcctgtctgcattcctcaggggagaaggtcacaatgacttcagggccagctcaagtgaag
 ttacatgcactggtaccagcagaagccaggatcctcccccacccctggatttatccccatccaacctggcttctggagtc
 cctgctcgttcagtggtcagtggtctgggaccttactctctcacaatcagcagagtggtgaggtgaagatgctgccactt
 attactgccagcagtggtgatttaccacccacgctcgggtgctgggaccaagctggagctgaaagatggcgggtggtcgt
 25 ggcgggtggtgagctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagtcggtgaggcctgggg
 cctcagtgaaatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
 gggcctgggaatggattggagctattatccaggaaatggtgatacttctcacaatcagaagttcaagggcaaggccacactg
 actgtagacaaaactcctcagcagcctacatgcagctcagcagcctgacatctgaagactctgcgggtctatttctgtgcaa
 gagtgggtgtactatagtaacttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcaggag
 30 cccaaatcttgtacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcttccc
 cccaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtgacgtgagccacgaagac
 cctgaggtcaagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
 aacagcacgtaccgtgtggtcagcgtcctaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaacccaaagggcagccccgagaaccacaggtgta
 35 caccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaggcttctatcccagcg
 acatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
 gtccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacacgc agaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 308)

40 mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyccqqsfnpptfagtklelkdgggsgggsgggssqaylqqsgaesvr
 pgasvkmsckasgytftslnmhvkwqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltسد
 savyfcarvvyysnsywyfdvwgtgtvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmsrtpevt
 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
 45 skpkgqpprepqvylppsrdeitknqvslclvkgfypsdiavewesngcipennyktpvldsdgsfflyskltvdk
 rwqqgnvfscsvmhhealhnhytqkslslspgk

44. G28-1VH

Nucleotide sequence: (SEQ ID NO: 309)

5 gcggtccagctgcagcagctctggacctgagctggaaaagcctggcgcttcagtgaaagatttcctgcaaggcttctggttact
cattcactggctacaatatgaactgggtgaagcagaataatggaaagagccttgagtggattggaaatattgaccttattatg
gtggtactacctacaaccggaagttcaagggaaggccacattgactgtagacaaatcctccagcacagcctacatgcag
ctcaagagctgacatctgaggactctgcagctattactgtgcaagatcggtcggccctatggactactgggggtcaaggaa
cctcagtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 310)

10 avqlqqsgpelekp gasv kisckasgysftgynmnwvkqnnkslewnidpyyggtynrkfk gkatltvdksst
taymq lksltsedsavyy carsvgpmdywggtsvtvssdq

45. G28-1VL

Nucleotide sequence: (SEQ ID NO: 311)

15 aagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagctcagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaaatgtttacagtta
tttggcttggtatcagcagaaacagggaatctcctcagctcctggtctctttgcaaaaaccttagcagaaggtgtgccatc
aaggttcagtggcagtggtatcaggcacacagttttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
20 ggtgggtcgggtggcggcggtatcgta

Amino acid sequence: (SEQ ID NO: 312)

mvstaqlglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgs gtqfslkisslqpedsgsyfcqhhsdnptwfgggteleikgggsgggsgggsgggss

25 46. G28-1 scFv

Nucleotide sequence: (SEQ ID NO: 313)

aagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagctcagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaaatgtttacagtta
30 tttggcttggtatcagcagaaacagggaatctcctcagctcctggtctctttgcaaaaaccttagcagaaggtgtgccatc
aaggttcagtggcagtggtatcaggcacacagttttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
ggtgggtcgggtggcggcggtatcgtaagggtccagctgcagcagcttgacctgagctggaagcctggcgcttcag
tgaagatttcctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaaagagcctt
gagtggattggaatatgaccttattatggtggtactacctacaaccggaagttcaagggaaggccacattgactgtaga
35 caaatcctccagcacagcctacatgcagctcaagagctgacatctgaggactctgcagcttauactgtgcaagatcggtc
ggccctatggactactgggggtcaaggaacctcagtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 314)

mvstaqlglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwfgggteleikgggsgggsgggsgggssavqlqqsgpelekp gas
40 vkisckasgysftgynmnwvkqnnkslewnidpyyggtynrkfk gkatltvdkssttaymq lksltsedsav
ycarsvgpmdywgg tsvtvssdq

47. G28-1 VHL11S

Nucleotide sequence: (SEQ ID NO: 315)

gcgggtccagctgcagcagctctggacctgagtcggaaaagcctggcgcttcagtgagatttcctgcaaggcttctggttact
cattcactggctacaatatgaactgggtgaagcagaataatggaaagagccttgagtggattggaaatattgatccttattatg
5 gtggtactacctacaaccggaagtcaagggcaaggccacattgactgtagacaaatcctccagcacagcctacatgcag
ctcaagagcttgacatctgaggactctgcagctcttactgtgcaagatcggtcggccctatggactactggggtcaaggaa
cctcagtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 316)

avqlqqsgpesekpgasvkiskasgysftgynmnwvkqnnkslewnidpyyggtynrkfkkgkatltvdks
10 staym qlksltsedsavyycarsvgpmdywgqtsvtvssdq

48. G28-1 VHL11S scFv

Nucleotide sequence: (SEQ ID NO: 317)

aagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
15 gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgtttacagtta
tttggcttggtatcagcagaaacagggaaaatctcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtgccatc
aaggttcagtggtcagtggtatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcgggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
20 ggtgggtcgggtggcggcggtatcgtcagcgggtccagctgcagcagcttgacctgagtcggaaaagcctggcgcttcag
tgaagatttcctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaaagagcctt
gagtggattggaatatgtatccttattatggtggtactacctacaaccggaagtcaagggcaaggccacattgactgtaga
caaatcctccagcacagcctacatgcagctcaagagcttgacatctgaggactctgcagctcttactgtgcaagatcggtc
ggccctatggactactggggtcaaggaacctcagtcaccgtctcttctgatcag

Amino acid sequence: (SEQ ID NO: 318)

mvstaqlflglllwlgtgrcdiqmtqspaslsasvgetvtitrtsenvysylawyqqkqkspqllvsfaktlaegvpsrf
25 sgsrgsgtqfslkisslpedsqsyfcqhhsdnptwfgggteleikgggsgggsgggsgggssavqlqqsgpesekpgas
vkiskasgysftgynmnwvkqnnkslewnidpyyggtynrktkgkatltvdksstaymqlksltsedsav
ycarsvg- pmdywgqg tsvtvssdq

30 49. G28-1 scFv (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 319)

aagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgtttacagtta
35 tttggcttggtatcagcagaaacagggaaaatctcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtgccatc
aaggttcagtggtcagtggtatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcgggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
ggtgggtcgggtggcggcggtatcgtcagcgggtccagctgcagcagcttgacctgagtcggaaaagcctggcgcttcag
tgaagatttcctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaaagagcctt
gagtggattggaatatgtatccttattatggtggtactacctacaaccggaagtcaagggcaaggccacattgactgtaga
40 caaatcctccagcacagcctacatgcagctcaagagcttgacatctgaggactctgcagctcttactgtgcaagatcggtc
ggccctatggactactggggtcaaggaacctcagtcaccgtctcttctgatcatgatcaggagcccaaatcttctgacaaaa
ctcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagttctcttctccccccaaaacccaaggaca
ccctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacgaagaccctgaggtcaagttaact
ggtacgtggacggcgtggaggtgcataatgccaagacaagccgcgggaggagcagtagacaacagcacgtaccgtgtg

gtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgcagggtctccaacaaagccctccc
 agcccccacgcagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccg
 ggatgagctgaccaagaaccaggtcagcctgacctgctgggtcaaaggcttctatcccagcgacatcgccgtggagtggg
 agagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttctctctacagc
 5 aagctcacctgggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaacca
 ctacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 320)

mvstaqlglillwltggrcdiqmtqspaslsasvgevvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsr
 fsgsgsgtqfslkisslqpedsgsyfcqhhsdnptwtfgggteleikggggsgggsgggssavqlqqsgpelekpga
 10 svkiskcasgysftgynmnwvkqnnngkslewignidpyyggtynrkfkkgkatltvdkssstaymqlksltsedsav
 yycarsvgpmdywgqgtsvtvssdhqepkssdkthtspssapellggpsvflfppkpkdtlmisrtpvctcvvd
 vshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapietiskakg
 qprepvytlppsrdeltknqvsltclvkgyfypsdiavewesngqpennyktpvldsdgsfflyskhvdksrwqqg
 nvfscsvmhealnhytqkslslspgk

15

50. G28-1 scFv IgAW H IgG1WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 321)

aagcttcccgcctatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
 gactcagttctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaataatgtttacagtta
 20 ttggcttggtatcagcagaaacagggaataatctcctcagctcctggtctctttgcaaaaaccttagcagaaggtgtccatc
 aaggttcagtgagcagtgatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
 gtcaacatcattccgataatccgtggacgttcgggtggaggcaccgaactggagatcaaagggtggcgggtggctcgggagggt
 ggtgggtcgggtggcggcgatcgtcagcgggtccagctgcagcagcttgacctgagctggaaaagcctggcgcttcag
 tgaagatttctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaaagagcctt
 25 gagggttgattggaatattgatccttattatggtgtgactacctaaccggaagtcaagggaaggccacattgactgtaga
 caaatcctccagcacagcctacatgcagctcaagagcttgacatctgaggactctgcagctctattactgtgaagatcggtc
 ggccctatggactactggggtcaaggaaacctcagtcaccgctcttctgatcagccagttccctcaactccacctaccccatc
 tccctcaactccacctaccccatctccctcatgcgcacctgaactcctggggggaccgtcagttctcttccccccaaaac
 ccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtgggtggacgtgagccacgaagaccctgaggt
 30 caagttaactggtacgtggacggcgtggaggtgcataatgccaaagcaaaagccgcgggaggagcagtacaacagcac
 gtaccgtgtgtgcagcgtcctaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaaca
 aagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgc
 cccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctgggtcaaaggcttctatcccagcgacatcgcc
 gtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttct
 35 tctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggct
 ctgcacaaccactacacgcaga agagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 322)

mvstaqlglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
 sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwtfgggteleikggggsgggsgggssavqlqqsgpelekpgas
 40 vkiskcasgysftgynmnwvkqnnngkslewignidpyyggtynrkfkkgkatltvdkssstaymqlksltsedsav
 ycarsvgpmdywgqgtsvtvssdqvpstptpsptpspascapellggpsvflfppkpkdtlmisrtpvctcvv
 vdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapietiskak
 kgciprepvytlppsrdeltknqvsltclvkgyfypsdiavewesngcipennyktpvldsdgsfflyskltvdksrw
 qqgnvfscsvmhealnhytqkslslspgk

45

51. G28-1 scFv VHL11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 323)

aagcttggcccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgtttacagtta
tttggcttggatcagcagaacagggaatatctcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtgccatc
5 aaggttcagtggcagtggtatcaggcacacagttttcttgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
gggtgggtcgggtggcggcggtatcgtcagcgggtccagctgcagcagcttgacctgagtcggaaaagcctggcgcttcag
tgaagatttctgcaaggcttctggttactcattcactgggtacaatatgaactgggtgaagcagaataatgaaagagcctt
gagtggattggaatatgtatccttattatggtggtactacctacaaccggaagtcaagggcaaggccacattgactgtaga
10 caaatctccagcacagcctacatgcagctcaagagctgtgacatctgaggactctgcagctctattactgtgcaagatcgggtc
ggccctatggactactgggtcaaggaacctcagtcaccgtctcttctgatcaggagcccaaatcttctgacaaaactcaca
catccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcctccccccaaaaccaaggacacctca
tgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtac
gtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggcagc
15 gtctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtcaaggtctccaacaaagccctccagcccc
catcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccggtatga
gctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccagcgacatcgccgtggagtgggagagc
aatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctcctacagcaagct
caccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctgcacaaccactaca
20 cgcagaagagcctctccctgtc tccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 324)

mvstaqlfllllwtlgrcdiqmtqspaslsasvgetvtitrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslpedsqsyfcqhhsdnptwfgggteleikgggsgggsgggssavqlqqsgpesekpgas
vkiskcasgysftgynmnwvkqnnkslewignidpyyggtynrkfkakatltvdkssstaymqlksltsedsavy
25 ycarsvgpmdywgqgtsvtvssdhdpkssdkthtspssapellggpsvflfppkpkdtlmsrtpevtcvvvdv
shedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeyckvsnkalpapiektiskakgq
prepqvylpprsdeltnqvsltlvkgfypsdiavewesngqpennyktpvpldsdsfflyskltvdksrwqqgn
vfscsvmhealhnhytqkslslspgk

30 52. G28-1 scFv VHL11S (CSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 325)

aagcttggcccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgtttacagtta
tttggcttggatcagcagaacagggaatatctcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtgccatc
35 aaggttcagtggcagtggtatcaggcacacagttttcttgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
gggtgggtcgggtggcggcggtatcgtcagcgggtccagctgcagcagcttgacctgagtcggaaaagcctggcgcttcag
tgaagatttctgcaaggcttctggttactcattcactgggtacaatatgaactgggtgaagcagaataatgaaagagcctt
gagtggattggaatatgtatccttattatggtggtactacctacaaccggaagtcaagggcaaggccacattgactgtaga
40 caaatctccagcacagcctacatgcagctcaagagctgtgacatctgaggactctgcagctctattactgtgcaagatcgggtc
ggccctatggactactgggtcaaggaacctcagtcaccgtctcttctgatcaggagcccaaatcttgtgacaaaactcaca
catccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcctccccccaaaaccaaggacacctca
tgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagcaactggtacgt
ggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggcagcgt
45 cctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtcaaggtctccaacaaagccctccagcccc
atcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccggtatgag
ctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccagcgacatcgccgtggagtgggagagca

atgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctc
accgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacac
gcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 326)

5 mvstaqflgllllwhggrcdiqmtqspaslsasvgetvtitrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwfgggteleikgggsgggsgggsgssavqlqqsgpesekpgas
vkisckasgysftgynmnwvkqnnkslewnidpyyggynnrklikgkatltvdkssstaymqlksltsedsav
ycarsvgpmdywgqgtsvtvssdqepkscdkthtspssapellggpsvflfppkpkdtlmsrtpevtcvvvdvsh
edpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqpr
10 epqvytlppsrdeitknqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvf
scsvmhealhnhytqkslslspgk

53. G28-1 scFv VH L11S (CSC-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 327)

15 aagcttgccgcatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgtttacagtta
tttggcttggtatcagcagaacagggaatatcctcagctcctggtctcttttgcataaaccttagcagaaggtgtgccatc
aaggttcagtggcagtgatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgUcgggtggaggcaccgaactggagatcaaaggtggcggtggctcgggcgg
20 tgggtgggtcgggtggcgcgatcgtcagcggtccagctgcagcagcttggtgacctgagtcggaaaagcctggcgcttca
gtgaagatttctgcaaggttctggttactcattcactggctacaatatgaactgggtgaagcagaataatgaaagagcct
tgagtggattggaatatgtatccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgactgtag
acaaatcctccagcacagcctacatgcagctcaagagctgacatctgaggactctgcagctattactgtgcaagatcggt
cggccctatggactactgggtgcaaggaacctcagtcaccgtctcttctgatcaggagcccaaatcttgacaaaactcac
25 acatctccaccgtgtcagcacctgaactcctgggtggaccgtcagctcttcttccccccaaaacccaaggacacctca
tgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagtcaactggtac
gtggacggcgtggaggtgcataatccaagacaaagccgcgggaggagcagtacaacagcagtgaccgtgtggtcagc
gtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccc
catcgagaaaacaatctccaaagccaaaggcgagccccgagaaccacaggtgtacaccctgccccatccgggatga
30 gctgaccaagaaccaggtcagcctgacctgcctggtcaaaaggcttctatccaagcgacatcgccgtggagtgggagagc
aatgggcagccggagaacaactacaagaccacgctcctcctgctggactccgacggctccttctctctacagcaagct
caccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactaca
cgcagaagagcctctccctgtct ccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 328)

35 rnvstaqflgllllwltggrcdiqmtqspaslsasvgetvtitrtsenvysylawyqqkqgkspqllvsfaktlaegvpsr
fsgsgsgtqfslkisslqpedsgsyfcqhhsdnptwfgggteleikgggsgggsgggsgssavqlqqsgpesekpga
svkisckasgysftgynmnwvkqnnkslewnidpyyggtynrkfkkgkatltvdkssstaymqlksltsedsav
yycarsvgpmdywgqgtsvtvssdqepkscdkthtspcsapellggpsvflfjpkpkdtlmsrtpevtcvvvdvsh
edpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqpr
40 epqvytlppsrdeitknqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvf
scsvmhealhnhytqkslslspgk

54. G28-1 scFv VH L11S (SSC-P)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 329)

aagcttggccgcatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
 gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatgtttacagtta
 tttggcttggtatcagcagaaacagggaaaatctcctcagctcctgtctcttttgcaaaaaccttagcagaaggttgccatc
 aaggttcagtggtcagtggtatcaggcacacagttttctgaagatcagcagcctgcagcctgaagattctggaagtatttct
 5 gtaacacatccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcggt
 ggtgggtcgggtggcggcggtatcgtcagcgggtccagctgcagcagcttgacctgagtcggaaaagcctggcgcttcag
 tgaagatttctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaagagcctt
 gagtggattggaatatgaccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgactgtaga
 caaatctccagcacagcctacatgcagctcaagagtctgacatctgaggactctgcagcttattactgtgaagatcggtc
 10 ggccctatggactactgggtcaaggaacctcagtcaccgtctcttctgatcaggagcccaaatcttctgacaaaactcaca
 catccccaccgtgccagcacctgaactctggggggaccgtcagcttctcttccccccaaaacccaaggacacctca
 tgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacgaagacctgaggtcaagttcaactggtagt
 ggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgt
 cctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccagccccc
 15 atcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatccgggatgag
 ctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagagca
 atgggcagccggagaacaactacaagaccacgcctcccgctggtgactccgacggctccttctctctacagcaagctc
 accgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacac
 gcagaagagcctctccctgtc tccgggtaaatgatctaga

20 Amino acid sequence: (SEQ ID NO: 330)

mvstaqlfllllwtlgrcdiqmtqspaslsasvgetvtitrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
 sgsgsgtqfslkisslpedsqsyfcqhhsdnptwfgggteleikgggsgggsgggsgggssavqlqqsgpeseckpgas
 vkiskcasgysftgynmnwvkqnnkslewnidpyyggtytnrkfkakatltvdksstaymqlksltsedsavy
 ycarsvgpmdywgqgtsvtvssdqepkssdkhtsppcpapellggpsvflfppkpkdtlmisrtpetcvvvdvsh
 25 edpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqpr
 epqvytlppsrdeltknqvsltlvkgfypsdiavewesngqpennnykttppvldsdgsfflyskltvdksrwwqgnvf
 scsvmhealhnhytqkslslspgk

II. 54. HCTLA4 HIGG1 (SSS-S)H P238SCH2 WCH3

30 Nucleotide sequence: (SEQ ID NO: 331)

atggcttgcccttgatttcagcggcacaaggctcagctgaacctggctgccaggacctggccctgcactctcctgtttttctt
 ctcttcacccctgtcttctgcaaagcaatgcacgtggcccagcctgctgtggtactggccagcagccgaggtatgccagct
 ttgtgtgtgagtatgcatctccaggcaaagccactgaggtccgggtgacagtgttcggcaggtgacagccaggtgactg
 aagtctgtcggcaacctacatgacggggaatgagttgaccttcttagatgattccatctgcacgggcacctccagtggaa
 35 atcaagtgaacctactatccaaggactgagggccatggacacgggactctacatctgcaaggtggagctcatgtaccac
 cgccatactacctgggcataggaacggaaccagatttatgtaattgatccagaacctgcccagattctgatcaacccaa
 atcttctgacaaaactcacacatccccaccgtctcagcacctgaactcctggggggatcgtcagcttctcttccccccaa
 aacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtgggtggacgtgagccacgaagacctga
 ggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtagaacag
 40 cacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggttcca
 acaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacc
 tcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatc
 gccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctggtgactccgacggctcc
 ttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgag
 45 gctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatga

Amino acid sequence: (SEQ ID NO: 332)

maclgfqrhkaqlnlaartwpctllfflfpvfkamhvaqpavvlassrgiasfvceyaspgkatevrvvtlrlqadsqv
tevcaatymtgneltfddsiectgtssgnqvnltiqglramdtglyickvelmypppyylgigngtqiyvidpepcpdsd
qpkssdkthtspssapellggssvflfppkpkdtlmisrtpevtcvvvdvshedpevkfhwyvdgvevhnaktkpr
eeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgciprepqvytlppsrdeltknqvsltclvkg
5 fypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvfscsvmhealhnhytqkslslspgk

55. Fc2-2 VL

Nucleotide sequence: (SEQ ID NO: 333)

gttgtaagcttgccgccatggattcacaggcccaggttcttatgttactgctgctatgggtatctggtacctgtggggacattg
10 tgatgtcacagtctccatcctcctagctgtgtcagttggagagaaggtttctatgagctgcaagtccagtcagagcctttata
taatcacaatcaaaagaactacttggcctggtaccagcagataaccaggggcagtcctcctaaactgctgatttactgggcatcca
ctagggaatctggggctcctgatcgcttcacaggcagtggtatctgggacagatttactctcaccatcagcagagtgaagg
ctgaagacctggcagtttattactgtcagcaatattatacctatcctccacgctcggaggtggcaccaagctggaaataaaa
gggtggcgggtggctcgggcgggtgggtgggtcgggtggcggcgggagctcg

15 Amino acid sequence: (SEQ ID NO: 334)

mdsqaqvlmllllwvsgtcgdivmsqspsslavsvgekvsmckssqsllynhnqnknylawyqqipgqspklli
wastres gvpdrftgsgsgtdftltisrvkaedlavyyccqyytypptfgggtkleikggggsgggsgggss

56. FC2-2VH

20 Nucleotide sequence: (SEQ ID NO: 335)

Gggagctcgcaggtgcagttgaaggagtcaggacctggcctggtggcgccctcacagagcctgtccatcacatgcacc
gtctcagggttctcattaaccgtctatggtgttaactgggttcgacagcctccaggaaagggtctggactggctgggaatgat
atgggggtgatggaagcacagactataattcagctctcaaatccagactgagcatcagtaaggacaactccaagagccaagt
tttcttaaaaatggacagctctacaaactgatgacacagccaggtactactgtgccagagatcactatggtaccactatgctat
25 ggactactggggtaaggaacctcagtcaccgtctcctctgatcag

Amino acid sequence: (SEQ ID NO: 336)

gssqvqlkesgpplvapsqslsitctvsgfsltygvnwvrppgkglwdlwmgiwgdgstdynsalksrslskdnsk
sqvflkmdslqtdtdaryycardhygthyamdywqgtsvtvssdq

30 **57. FC2-2scFv**

Nucleotide sequence: (SEQ ID NO: 337)

gttgtaagcttgccgccatggattcacaggcccaggttcttatgttactgctgctatgggtatctggtacctgtggggacattg
tgatgtcacagtctccatcctcctagctgtgtcagttggagagaaggtttctatgagctgcaagtccagtcagagcctttata
taatcacaatcaaaagaactacttggcctggtaccagcagataaccaggggcagtcctcctaaactgctgatttactgggcatcca
35 ctagggaatctggggctcctgatcgcttcacaggcagtggtatctgggacagatttactctcaccatcagcagagtgaagg
ctgaagacctggcagtttattactgtcagcaatattatacctatcctccacgctcggaggtggcaccaagctggaaataaaa
gggtggcgggtggctcgggcgggtgggtgggtcgggtggcggcgggagctctcaggtgcagttgaaggagtcaggacctggc
ctgggtggcgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccgtctatggtgttaactgggttcg
ccagcctccaggaaagggtctggactggctgggaatgatatggggtgatggaagcacagactataattcagctctcaaatc
40 cagactgagcatcagtaaggacaactccaagagccaagtttcttaaaaatggacagctctacaaactgatgacacagccag
gtactactgtgccagagatcactatggtaccactatgctatggactactggggtcaaggaacctcagtcaccgtctcctctg
atcag

Amino acid sequence: (SEQ ID NO: 338)

mdsqaqvlmllllwvsgtcgdivmsqspsslavsvgekvsmckssqsllynhnqknylawyqqipgqspklliy
wastresvpdrftgsgsgtdftltsrvkaedlavyyccqyytypptfgggtkleikgggsgggsgggssqvqlkes
gpglvapsqslsitctvsgfsltygvnwvrppgkgldwlgmiwgdgstdynsalksrlsiskdnskqvflkmdsl
qtdtdaryycardhygt hyamdywgqgtsvtvssdq

5

58. FC2-2 VHL11S

Nucleotide sequence: (SEQ ID NO: 339)

gggagctctcaggtgcagttgaaggagtcaggacctggctcgggtggcgccctcacagagcctgtccatcacatgcaccgt
ctcagggttctcattaaccgtctatggtgttaactgggttcgccagcctccaggaaagggtctggactggctgggaatgatat
10 ggggtgatggaagcacagactataattcagctctcaaaccagactgagcatcagtaaggacaactccaagagccaagttt
tcttaaaaatggacagtctacaaactgatgacacagccaggtactactgtgccagagatcactatggtaccactatgctatg
gactactggggtcaaggaacctca gtcaccgtctcctctgatcag

Amino acid sequence: (SEQ ID NO: 340)

(gss)qvqlkesgpgsvapsqslsitctvsgfsltygvnwvrppgkgldwlgmiwgdgstdynsalksrlsiskdnsk
15 qsvfl kmdslqtdtdaryycardhygthyamdywgqgtsvtvssdq

59. FC2-2 VH L11S scFv

Nucleotide sequence: (SEQ ID NO: 341)

gttgtaagcttggcccatggattcacaggcccaggttcttatgttactgctgctatgggtatctggtacctgtggggacattg
20 tgatgtcacagcttccatcctccctagctgtgtcagttggagagaagggttctatgagctgcaagtccagtcagagcctttata
taatcacaaatcaaaagaactacttggcctggtagcagataccagggcagtcctctaaactgctgatttactgggcatcca
ctaggggaatctgggggtccctgatcgcttcacaggcagtgatctgggacagatttcactctcaccatcagcagagtgaagg
ctgaagacctggcagtttattactgtcagcaatattatacctatcctccacgttcggaggtggcaccgaagctggaaataaaa
gggtggcggtggctcgggcgggtggtgggtcgggtggcggggagctctcaggtgcagttgaaggagtcaggacctggc
25 tcggtggcgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccgtctatggtgttaactgggttcg
ccagcctccaggaaagggtctggactggctgggaatgatatgggtgatggaagcacagactataattcagctctcaaatc
cagactgagcatcagtaaggacaactccaagagccaagtttcttaaaaatggacagtctacaaactgatgacacagccag
gtactactgtgccagagatcactatggtaccactatgctatggactactgggtcaaggaacctcagtcaccgtctcctctg
atcag

30 Amino acid sequence: (SEQ ID NO: 342)

Mdsqaqvlmllllwvsgtcgdivmsqspsslavsvgekvsmckssqsllynhnqknylawyqqipgqspklliy
wastresvpdrftgsgsgtdftltsrvkaedlavyyccqyytypptfgggtkleikgggsgggsgggssqvqlk
esgpgsvapsqslsitctvsgfsltygvnwvrppgkgldwlgmiwgdgstdynsalksrlsiskdnskqvflkm
dslqtdtdaryycardhygt hyamdywgqgtsvtvssdq

35

60. FC2-2 (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 343)

gttgtaagcttggcccatggattcacaggcccaggttcttatgttactgctgctatgggtatctggtacctgtggggacattg
40 tgatgtcacagcttccatcctccctagctgtgtcagttggagagaagggttctatgagctgcaagtccagtcagagcctttata
taatcacaaatcaaaagaactacttggcctggtagcagataccagggcagtcctctaaactgctgatttactgggcatcca
ctaggggaatctgggggtccctgatcgcttcacaggcagtgatctgggacagatttcactctcaccatcagcagagtgaagg
ctgaagacctggcagtttattactgtcagcaatattatacctatcctccacgttcggaggtggcaccgaagctggaaataaaa
gggtggcggtggctcgggcgggtggtgggtcgggtggcggggagctctcaggtgcagttgaaggagtcaggacctggc
ctggtggcgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccgtctatggtgttaactgggttcg

ccagcctccaggaaagggtctggactggctgggaatgatatgggggtgatggaagcacagactataattcagctctcaaate
 cagactgagcatcagtaaggacaactccaagagccaagttttcttaaaaatggacagtctacaaactgatgacacagccag
 gtactactgtgccagagatcactatggtaccactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctctg
 atcaggagcccaaatctcttgacaaaactcacacatccccaccgtcctcagcacctgaactcctgggtggaccgtcagctctt
 5 cctcttccccccaaaaccaaggacacctcatgatctcccggaccctgaggtcacatgcgtgggtgggtggacgtgagcc
 acgaagacctgaggtcaagtcaactggtagctggacggcgtggaggtgcataatgccaagacaaagccgcgggagg
 agcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaa
 gtgcaaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaacc
 acaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttct
 10 atccaagcgacatcgccgtggagtgggagagcaatgggcagccgggagaacaactacaagaccacgcctcccgtgctgg
 actccgacggctccttctcctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgct
 ccgtgatgcatgaggctctgcaca accactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 344)

mdsqaqvlmlllwvsgtcgdivmsqspsslavsvgekvsmckssqsllynhnqnknylawyqqipgqspklliy
 15 wastresgvprftgsgsgtdfthisrvkaedlavyyccqyytyppftggggtkleikgggsgggsgggssqvqlke
 sgpglvapsqslsitctvsgfslivygvnwvrppgkglwdlwmigdgstdynsalksriskdnksqvflkmds
 lqtdtdaryycardhygthyamdywgqgtsvtvssdqepkssdkhtsppssapellggpsvflfppkpkdtlmsrpt
 evtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapi
 20 ektiskakgqprepvytlppsrdeltknqvsitclvkgfypsdiavewesngpennyktpvldsdgsttlyskltvd
 ksrwqqgnvfscsvmhealhnhytqkslspsgk

61. FC2-2 VHL11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 345)

gttgtaagcttgccgcatggattcacaggcccaggttctatgttactgctgctatgggtatctggtacctgtggggacattg
 25 tgatgtcacagtctccatcctccctagctgtgtcagttggagagaaggtttctatgagctgcaagtccagtcagagccctttata
 taatcacaatcaaaagaactacttgacctggtagcagataaccagggcagctcctaaactgctgatttactgggcatcca
 ctagggaatctggggtcctgatcgcttcacaggcagtgatctgggacagattcactctcaccatcagcagagtgaagg
 ctgaagacctggcagtttattactgtcgaatattatacctatcctccacgttcggaggtggcaccagctggaaataaaa
 ggtggcgggtggctcgggcggtggtgggtcgggtggcggggagctctcaggtgcagttgaaggagtcaggacctggc
 30 tcgggtggcgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccgtctatggtgtaactgggttcg
 ccagcctccaggaaagggtctggactggctgggaatgatatgggggtgatggaagcacagactataattcagctctcaaate
 cagactgagcatcagtaaggacaactccaagagccaagttttcttaaaaatggacagtctacaaactgatgacacagccag
 gtactactgtgccagagatcactatggtaccactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctctg
 atcaggagcccaaatctcttgacaaaactcacacatccccaccgtcctcagcacctgaactcctgggtggaccgtcagctctt
 35 cctcttccccccaaaaccaaggacacctcatgatctcccggaccctgaggtcacatgcgtgggtgggtgacgtgagcc
 acgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggagg
 agcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaa
 gtgcaaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaacc
 acaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttct
 40 atccaagcgacatcgccgtggagtgggagagcaatgggcagccgggagaacaactacaagaccacgcctcccgtgctgg
 actccgacggctccttctcctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgct
 ccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 346)

mdsqaqvlmlllwvsgtcgdivmsqspsslavsvgekvsmckssqsllynhnqnknylawyqqipgqspklliy
 45 wastresgvprftgsgsgtdfthisrvkaedlavyyccqyytyppftggggtkleikgggsgggsgggssqvqlk
 espgsvapsqslsitctvsgfslivygvnwvrppgkglwdlwmigdgstdynsalksriskdnksqvflkmds
 dslqtdtdaryycardhygthyamdywgqgtsvtvssdqepkssdkhtsppssapellggpsvflfppkpkdtlmsrpt

rtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalp
apiektiskakgqprepqvylppsrdeitknqvslclvkgfypsdiavewesngqpennykttppvldsdgsttlysk
ltvdksrwqqgnvfscsvmhleahnhytqkslsispk

5 **62. UCHL-1 VH**

Nucleotide sequence: (SEQ ID NO: 347)

atgggcaggcttacttctcattcctgctactgattgtcctgcatatgctctctccagattactctgaaagagtctggccctgg
gatcttgcagccctcccagaccctcagctctgacttgttcttctctgggtttcactgaccacttatggtataggagtaggttga
ttcgtcagcctccagggaagggtctggagtggctgacacacatttgggtgaatgataataagtactataacacagccctgag
10 gagccggctcacaatctccaaggattctccaacaaccaagtactcctcaagatcgccaatgtggacactgcagataccgc
cacatactactgtctctacggctac acttactggggccaagggactctggctactgtctctgca

Amino acid sequence: (SEQ ID NO: 348)

mgrltssflhlivpayvlsqitlkesgpgilqpsqtlsitcsfsgflttygigvgwirqppgkglewlthiwwndnkyynt
alrslrtiskdssnnqvllkianvtdadtatytyclygytywgqgtlvtvsa

15

63. UCHL-1 VL

Nucleotide sequence: (SEQ ID NO: 349)

atgaagttgcctgttaggctgttggtgctgatgttctggattcctgcttccatcagtgatgttgatgacccaaactccactctc
cctgcctgtcagctcttgagatcaggcctccatctcttgcagatctagtcagagccttctttacagtaatggaaacacctatta
20 catttggtacctgcagaagccaggccagctctccaaaactcctgatctacaaacttccaaccgattttctgggtcccagacag
gttcagtggtcagtgatcaggacagatttcacactcaagatcagcagagtggaggctgaggatctgggagtttatttctgc
tctcaaagtacacatgttccgtgga cgttcggtggaggcaccaagctggaaatcaaa

Amino acid sequence: (SEQ ID NO: 350)

mklpvrlvlmfwipasisdvmtqtplslpvsldqasiscrssqsllysngntylhwylqkpgqspklliyklsnrfs
25 gvpdrfsgsgsgtdftlkisrveaedlgvyfcsqsthvpwtfgggtkleik

64. UCHL-1 scFv

Nucleotide sequence: (SEQ ID NO: 351)

gttgtaagcttgccgcatgaagttgcctgttaggctgttggtgctgatgttctggattcctgcttccatcagtgatgttgatg
30 acccaaactccactctccctgcctgtcagctcttgagatcaggcctccatctcttgcagatctagtcagagccttctttacagta
atggaaacacctatttacattggtacctgcagaagccaggccagctctccaaaactcctgatctacaaacttccaaccgatttt
ctgggggtcccagacaggttcagtggtcagtgatcaggacagatttcacactcaagatcagcagagtggaggctgaggat
ctgggagtttatttctgctctcaaagtacacatguccgtggacgttcggtggaggcaccaagctggaaatcaaagatggcg
gtggctcgggcggtgttgatctggaggaggtgggagctctcagattactctgaaagagtctggccctgggatcttcag
35 cctcccagaccctcagctgacttgttcttctctgggtttcactgaccacttatggtataggagtaggttgattcgtcagcc
tccaggaagggtctggagtggctgacacacatttgggtgaatgataataagtactataacacagccctgaggagccggct
cacaatctccaaggattcctccaacaaccaagtactcctcaagatcgccaatgtggacactgcagataccgccacatactac
tgtctctacggctacacttactggggccaagggactctggctactgtctctgctg atca

Amino acid sequence: (SEQ ID NO: 352)

40 mklpvrlvlmfwipasisdvmtqtplslpvsldqasiscrssqsllysngntylhwylqkpgqspklliyklsnrfs
gvpdrfsgsgsgtdftlkisrveaedlgvyfcsqsthvpwtfgggtkleikdggsgggsgggsgggsggtlkesgpgilq

psqtlsltsfsgfslttygigvgwirppgkglewlthiwwndnkyyntalrsrltiskdssnnqvlkianvdtadtaty
yclgytywggqtlvtvsad

65. UCHL-1 VH I11SL12S

5 Nucleotide sequence: (SEQ ID NO: 353)

gggagctctcagattactctgaaagagtctggccctgggactctgcagccctcccagaccctcagtctgacttgttttctct
gggtttcactgaccacttatggtataggagtaggttgattctgcagcctccaggaagggtctggagtggctgacacaca
tttggtggaatgataataagtactataacacagccctgaggagccggctcacaatctcaaggattctccaacaaccaagt
actcctcaagatcgccaatgtggacactgcagataccgccacatactactgtctctacggctacacttactggggccaaggg
10 actctggtcactgtctctgctgatca

Amino acid sequence: (SEQ ID NO: 354)

(gss)qitlkesgpgssqpsqtlsltsfsgfslttygigvgwirppgkglewlthiwwndnkyyntalrsrltiskdssn
nqvlkianvdtadtatyyclgytywggqtlvtvsad

15 **66. UCHL-1 scFv VH L11S**

Nucleotide sequence: (SEQ ID NO: 355)

gttgtaagcttgccgccatgaagttgcctgttaggctgttggtgctgatgttctggattcctgcttccatcagtgatgttgatg
acccaaactccactctccctgcctgtcagctctggagatcaggcctccatctctgcagatctagtcagagccttctttacagta
atggaaacacctatttacattggtacctgcagaagccaggccagctctccaaactcctgatctacaaactttccaaccgatttt
20 ctgggggtcccagacaggttcagtggcagtggtacagggacagatttcacactcaagatcagcagagtggaggctgaggat
ctgggagtttatttctgctctcaaagtacacatgttccgtggacgttcggtggaggcaccaagctggaaatcaaagatggcg
gtggctcgggcggtggtggatctggaggaggtgggagctctcagattactctgaaagagtctggccctgggagctcccag
ccctcccagaccctcagtctgacttgttcttctctgggttttactgaccacttatggtataggagtaggttgattcgtcagcc
tccaggaagggtctggagtggctgacacacatttggtggaatgataataagtactataacacagccctgaggagccggct
25 cacaatctccaaggattcctccaacaaccaagtactcctcaagatcgccaatgtggacactgcagataccgccacatactac
tgtctctacggctacacttactggggccaagggactctggtcactgtctctgct gatca

Amino acid sequence: (SEQ ID NO: 356)

mklpvrllvlmfwipasisdvmtqtplslpvsldqasiscrssqsllysnntylhwylqkpgqspklliyklsnrfs
gvpdrfsgsgsgtdftlkisrveadlgvyfcsqsthpwtfgggtkleikdggsgggsgggssqitlkesgpgssq
30 psqtlsltsfsgfslttygigvgwirppgkglewlthiwwndnkyyntalrsrltiskdssnnqvlkianvdtadtaty
yclgytywggqtlvtvsad

67. UCHL-1 scFv (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 357)

gttgtaagcttgccgccatgaagttgcctgttaggctgttggtgctgatgttctggattcctgcttccatcagtgatgttgatg
acccaaactccactctccctgcctgtcagctctggagatcaggcctccatctctgcagatctagtcagagccttctttacagta
atggaaacacctatttacattggtacctgcagaagccaggccagctctccaaactcctgatctacaaactttccaaccgatttt
ctgggggtcccagacaggttcagtggcagtggtacagggacagatttcacactcaagatcagcagagtggaggctgaggat
ctgggagtttatttctgctctcaaagtacacatgttccgtggacgttcggtggaggcaccaagctggaaatcaaagatggcg
40 gtggctcgggcggtggtggatctggaggaggtgggagctctcagattactctgaaagagtctggccctgggactctgcag
ccctcccagaccctcagtctgacttgttcttctctgggttttactgaccacttatggtataggagtaggttgattcgtcagcc
tccaggaagggtctggagtggctgacacacatttggtggaatgataataagtactataacacagccctgaggagccggct
cacaatctccaaggattcctccaacaaccaagtactcctcaagatcgccaatgtggacactgcagataccgccacatactac

5 tgtctctacggctacacttactggggccaagggactctggctactgtctctgctgatcaggagcccaaatcttctgacaaaac
 tcacacatccccaccgtcctcagcacctgaactcctgggtggaccgtcagcttctcttccccccaaaaccaaggacacc
 ctcatgatctcccgaccctgaggtcacatgcgtgggtggacgtgagccacgaagaccctgaggtcaagttcaactg
 glacgtggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggt
 10 cagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggttccaacaaagccctccca
 gccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccgg
 gatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccaagcgacatcgccgtggagtggga
 gagcaatgggcagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagca
 agtcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccac
 15 tacacgcagaagagcctctc cctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 358)

15 mklpvrllvlmfwipasisdvmtqtplslpvslgdqasiscrssqsllysnngtylhwylqkpgqspklliyklsnrfs
 gvpdrfsgsgsgtdtlkisirveadlgvyfcsqsthpwtfgggtkleikdggsgggsgggssqitlkesgpgilq
 psqtlsltcfsfsgfltygigvgwirppgkglewlthiwwndnkyntalrsrltiskdssnnqvllkianvdtadtaty
 yclgytywggqtlvtvsadqepkssdkthtspssapellggpsvflfppkpkdtlmisrtpvtecvvvdvshedpe
 vkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgpprepqv
 ytlppsrdeltknqvsltclvkfypsdiavewesngqpennykttppvldsdgsttlyskltvdksrwqqgnvfscsv
 mhealhnhytqkslslspgk

20 68. UCHL-1 scFv VHL11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 359)

25 gtgttaagcttgcgccatgaagttgcctgttaggctgttggtgctgatgttctggattcctgcttccatcagtgatgttgatg
 acccaaaactccactctccctgcctgtcagcttggagatcaggcctccatctcttgcagatctagttagagccttctttacagta
 atggaaacacctatttaccattggtacctgcagaagccaggccagctctccaaaactcctgatctacaaactttccaaccgattt
 30 ctgggggtcccagacaggttcagtgagcagtgatcaggagacagatttcaactcaagatcagcagagtgaggctgaggtat
 ctgggagtttatttctgctctcaaagtacacatgttccgtggacgttcggtggaggcaccaagctggaaatcaaatgagcgc
 gtggctcgggcgggtggtgatctggaggaggtgggagctctcagattactctgaaagagtctggccctgggagctcccag
 cctcccagaccctcagctgactgttcttctctgggttttactgaccacttatggtataggagtaggttgattcgtcagcc
 tccagggaagggtctggagtggctgacacacatttgggaatgataataagtactataacacagccctgaggagccggct
 35 cacaatctccaaggattcctccaacaaccaagtactcctcaagatcgccaatgtggacactgcagataaccgccacatactac
 tgtctctacggctacacttactggggccaagggactctgtgctactgtctctgctgatcaggagcccaaatcttctgacaaaac
 tcacacatccccaccgtcctcagcacctgaactcctgggtggaccgtcagcttctcttccccccaaaaccaaggacacc
 ctcatgatctcccgaccctgaggtcacatgcgtgggtggacgtgagccacgaagaccctgaggtcaagttcaactg
 gtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgggtca
 40 gcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggttccaacaaagccctcccagc
 ccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccggga
 tgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccaagcgacatcgccgtggagtgggaga
 gcaatgggcagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaag
 ctaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccacta
 45 cacgcagaagagcctctc cctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 360)

45 mklpvrllvlmfwipasisdvmtqtplslpvslgdqasiscrssqsllysnngtylhwylqkpgqspklliyklsnrfs
 gvpdrfsgsgsgtdftlkisirveadlgvyfcsqsthpwtfgggtkleikdggsgggsgggssqitlkesgpgssq
 psqtlsltcfsfsgfltygigvgwirppgkglewlthiwwndnkyntalrsrltiskdssnnqvllkianvdtadtaty
 yclgytywggqtlvtvsadqepkssdkthtspssapellggpsvflfppkpkdtlmisrtpvtecvvvdvshedpe
 vkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgpprepqv

ytlppsrdeltknqvsitclvkgyfypsdiavewesngqpennykttppvldsdgsttlyskltvdksrwqqgnvfscsv
mhealhnhytqkslsispk

69. 5B9 VH

5 Nucleotide sequence: (SEQ ID NO: 116)

atggctgtcttggggctgctcttctgcctggtgacatttccaagctgtgcctatcccagggtgcagctgaagcagtcaggacc
tggcctagtgcagtcctcacagagcctgtccatcacctgcacagtctctggtttctcattaactacctatgctgtacactgggtt
cgccagctctccaggaaagggctctggagtggctgggagtgatatggagtggggaatcacagactataatgcagctttcatat
ccagactgagcatcaccaaggacgattccaagagccaagttttctttaaataaacagtctgcaacctaatgacacagccat
10 ttattactgtgccagaaatgggggtgataactacccttattactatgctatggactactggggtaaggaacctcagtcaccgt
ctctca

Amino acid sequence: (SEQ ID NO: 118)

mavlgllfclvtfpscvlsqvqlkqsgpglvqssqslsitctvsgfslttyavhwvrqspgkglewlgviwsggitdyna
afisrlsi tkddsksqvttkmnslqpndtaiyyccarnngdnpyyyyamdywgqgtsvtvss

15

70. 5B9 VL

Nucleotide sequence: (SEQ ID NO:120)

atgaggttctctgctcagcttctggggctgcttgctctggatccctggatccactgcagatattgtgatgacgcaggctgca
tttccaatccagtcactcttgaacatcagcttccatctctgcagggtctagtaagagtctcctacatagtaatggcatcactta
20 tttgtattggtatctgcagaagccaggccagctctcctcagctcctgatttatcagatgtccaaccttgctcaggagtcacaga
cagggtcagtagcagtggtgaggaactgatttcacactgagaatcagcagagtggaggctgaggatgtgggtgtttattac
tgtgtcaaaatctagaactccgct cacgttcggtgctgggaccaagctggagctgaaacgg

Amino acid sequence: (SEQ ID NO:121)

mrfsaqllgllvlwipgstadivmtqaafsnpvtlgtasiscrskslhsngitylywylqkpgqspqlliyqmsnlas
25 gvpdrfsssgsgtdflrisrveaedvgvyycanlelpltfagtklelkr

71. 5B9scFv

Nucleotide sequence: (SEQ ID NO: 122)

Aagcttggcgccatgaggttctctgctcagcttctggggctgcttgctctggatccctggatccactgcagatattgtgatg
30 acgcaggctgcatttccaatccagtcactcttgaacatcagcttccatctctgcagggtctagtaagagtctcctacatagt
aatggcatcacttattgtattggtatctgcagaagccaggccagctctcctcagctcctgatttatcagatgtccaaccttgct-
caggagtcccagacaggttcagtagcagtggtgaggaactgatttcacactgagaatcagcagagtggaggctgaggat
gtgggtgtttattactgtgctcaaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacgggggtg
gcgggtggctcgggcgggtgggtgggtcgggtggcggcgggagctctcaggtgcagctgaagcagtcaggacctggcctag
35 tgcagtcctcacagagcctgtccatcacctgcacagtctctggtttctcattaactacctatgctgtacactgggttcgccagtc
tccaggaaagggctctggagtggctgggagtgatatggagtggggaatcacagactataatgcagctttcatatccagactg
agcatcaccaaggacgattccaagagccaagttttctttaaataaacagtctgcaacctaatgacacagccatttattactgt
gccagaaatgggggtgataactacccttattactatgctatggactactggggtaaggaacctcagtcaccgtctcctcag

Amino acid sequence: (SEQ ID NO: 123)

40 mrfsaqllgllvlwipgstadivmtqaafsnpvtlgtasiscrskslhsngitylywylqkpgqspqlliyqmsnlas
gvpdrfsssgsgtdflrisrveaedvgvyycanlelpltfagtklelkr

qssqslsitctvsgfslttyavhwvrqspgkglewlgviwsggitdynaafisrlsitkddsksqvttkmnslqpndtaiy
ycarnggdnyppyya mdywgqgtsvtvss

72. 5B9 VH L11S

5 Nucleotide sequence: (SEQ ID NO: 361)

gggagctctcaggtgcagctgaagcagtcaggacctggctcagtcagtcctcacagagcctgtccatcacctgcacagt
ctctggtttctcattaactacatctgtgtacactgggttcgccagctctccaggaaagggctggagtggtgggagtgatg
gagtggtggaatcacagactataatgcagctttcatatccagactgagcatcaccaaggacgattccaagagccaagtttct
ttaaataaagcagctgtcaacctaataacacagccatttattactgtgccagaaatgggggtgataactacccttattactatgc
10 tatggactactgggggtcaaggaac ctcagtcaccgtctcctcag

Amino acid sequence: (SEQ ID NO: 362)

(gss)qvqlkqsgpgsvqssqslsitctvsgfslttyavhwvrqspgkglewlgviwsggitdynaafisrlsitkddsksqvttk
mnslqpndtaiyyacamggdnyppyyamdywgqgtsvtvss

15 73. 5B9 VH L11S scFv

Nucleotide sequence: (SEQ ID NO: 363)

aagcttgccgcatgaggttctctgtcagcttctggggctgcttgctctggatccctggatccactgcagatattgtgatg
acgcaggtgcattctccaatccagtcactcttggaacatcagcttccatctcctgcaggtctagtaagagtcctacatagt
aatggcatcacttatttattggtatctgcagaagccaggccagctcctcagctcctgatttatcagatgtccaaccttgccctc
20 aggagtcacagacaggttcagtagcagtggtgcaggaaactgatttcacactgagaatcagcagagtgagggtgaggatg
tgggtgtttattactgtgctcaaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacgggggtgg
cggtggctcgggcggtggtgggtcgggtggcggcgggagctctcaggtgcagctgaagcagtcaggacctggctcagt
gcagtcctcacagagcctgtccatcacctgcacagctctctggtttctcattaactacatctgtgtacactgggttcgccagctc
ccaggaaagggctggagtggtgggagtgatggagtggtggaatcacagactataatgcagctttcatatccagactg
25 agcatcaccaaggacgattccaagagccaagtttctttaaataaagcagctgtcaacctaataacacagccatttattactgt
gccagaaatgggggtgataactacccttattactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctcag

Amino acid sequence: (SEQ ID NO: 364)

mrfsaqllgllvlwipgstadivmtqaafsnpvtlgtasiscrskslhsngitylywylkpgqspqlliyqmsnlas
gvpdrf ssssgstdftrsrveaedvgvycaqnlelpltfagtklelkrgggsgggsgggsgssqvqlkqsgpgs-
30 vqssqslsitctvsgfslttyavhwvrqspgkglewlgviwsggitdynaafisrlsitkddsksqvftkmnslqpndtai
yyacamggdnyppyya mdywgqgtsvtvss

73. 5B9 scFv (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 132)

aagcttgccgcatgaggttctctgtcagcttctggggctgcttgctctggatccctggatccactgcagatattgtgatg
acgcaggtgcattctccaatccagtcactcttggaacatcagcttccatctcctgcaggtctagtaagagtcctacatagt
aatggcatcacttatttattggtatctgcagaagccaggccagctcctcagctcctgatttatcagatgtccaaccttgccctc
35 aggagtcacagacaggttcagtagcagtggtgcaggaaactgatttcacactgagaatcagcagagtgagggtgaggatg
tgggtgtttattactgtgctcaaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacgggggtgg
cggtggctcgggcggtggtgggtcgggtggcggcgggagctctcaggtgcagctgaagcagtcaggacctggcctagt
40 gcagtcctcacagagcctgtccatcacctgcacagctctctggtttctcattaactacatctgtgtacactgggttcgccagctc
ccaggaaagggctggagtggtgggagtgatggagtggtggaatcacagactataatgcagctttcatatccagactg
agcatcaccaaggacgattccaagagccaagtttctttaaataaagcagctgtcaacctaataacacagccatttattactgt

gccagaaatgggggtgataactacccttattactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctctga
 tcaggagcccaaatcttctgacaaaactcacatccccaccgtcctcagcacctgaactcctgggtggaccgtcagctctc
 ctcttccccccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtgggtggacgtgagcca
 cgaagaccctgaggtcaagttcaactggtagtgacggcgtggaggtgcataatgccaagacaaagccgcgggagga
 5 gcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaag
 tgcaaggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaacca
 caggtgtacaccctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctat
 ccaagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctgga
 10 ctccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctc
 cgtgatgcatgaggctctgcacaac cactacacgcagaagagcctctccctgtctccgggtaaatgatctagag

Amino acid sequence: (SEQ ID NO: 133)

mrfsaqllgllvlwipgstadivmtqaafsnpvtlgtssisrskslhsngitylywylqkpgqspqlliyqmsnlas
 gvpdrfsssgsgtdflrisrveaedvgvycaqnlelpltfagtklelkrggsgggsgggssqvqlkqsgpglv
 qssqslsitctvsgfsltyavhwvrqspgkglewlgviwsggtdynaafisrlsitkddsksqvftkmnslqpndtaiy
 15 ycarnggdnyppyyamdywgqgtsvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmsirtpetv
 vvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeyckvsnkalpapiektis
 kakgqprepvytlppsrdeltknqvsitclvkgyfypsdiavewesngqpennyktpvldsdgstitlyskltvdksr
 wqqgnvfscsvmhealhnhytqkslslspgk

20 74. 5B9 scFv VHL11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 365)

aagcttgccgccatgaggttctctgctcagcttctggggctgcttgtgctctggatccctggatccactgcagatattgtgatg
 acgcaggtgcattctccaatccagtcactcttgaacatcagcttccatctcctgcaggtctagtaagagtctcctacatagt
 aatggcatcacttatttatttggatctgagaaagccaggccagtcctcagctcctgatttatcagatgtccaaccttgccct
 25 aggagtcacagacaggttcagtagcagtggttcaggaaactgatttcacactgagaatcagcagagtggaggctgaggatg
 tgggtgttattactgtgtcaaaatctagaactccgctcacgttcggtgctgggaccaagctggagctgaaacgggggtg
 cgggtggtcgggcgggtggtgggtcgggtggcggcgggagctctcaggtgcagctgaagcagtcaggacctggctcagt
 gcagtcctcagagacctgtccatcacctgcacagctctctggtttctattaactacctatgctgtacactgggttcgccagct
 ccaggaaagggctgagtggtgggtgagtgatattgagtggtggaatcacagactataatgcagcttccatccagactg
 30 agcatcaccaaggacgattccaagagccaagtttctttaaataaagacagcttgcaacctaataacacagccatttattactgt
 gccagaaatgggggtgataactacccttattactatgctatggactactgggggtcaaggaacctcagtcaccgtctcctctga
 tcaggagcccaaatcttctgacaaaactcacatccccaccgtcctcagcacctgaactcctgggtggaccgtcagctctc
 ctcttccccccaaaaccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtgggtggacgtgagcca
 cgaagaccctgaggtcaagttcaactggtagtgacggcgtggaggtgcataatgccaagacaaagccgcgggagga
 35 gcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaag
 tgcaaggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaacca
 caggtgtacaccctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctat
 ccaagcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctgga
 ctccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctc
 40 cgtgatgcatgaggctctgcacaac cactacacgcagaagagcctctccctgtctccgggtaaatgatctagag

Amino acid sequence: (SEQ ID NO: 366)

mrfsaqllgllvlwipgstadivmtqaafsnpvtlgtssisrskslhsngitylywylqkpgqspqlliyqmsnlas
 gvpdrfsssgsgtdflrisrveaedvgvycaqnlelpltfagtklelkrggsgggsgggsgggssqvqlkqsgpgsv
 qssqslsitctvsgfsltyavhwvrqspgkglewlgviwsggtdynaafisrlsitkddsksqvftkmnslqpndtaiy
 45 ycarnggdnyppyyamdywgqgtsvtvssdqepkssdkthtspssapellggpsvflfppkpkdtlmsirtpetv
 vvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeyckvsnkalpapiektis

kakgqprepqvylppsrdeltnqvslclvkgyfypsdiavewesngqpennykttppvldsdgsttlyskltvdksr
wqqgnvfscsvmhealhnhytqkslsispkg

75. 2H7 scFv (SSS-S)H P238SCH2 WCH3

5 Nucleotide sequence: (SEQ ID NO: 130)

aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcatactccaggggagaagggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctccccaacacctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
10 attactgccagcagtgaggttttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggaggctcg
ggcgggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgctctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
15 gactggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggatcgtcagttctctcttccc
cccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtgggtggtggacgtgagccacgaagac
cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
aacagcacgtaccgtgtggtcagcgtcctaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
20 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
caccctgccccatcccggtgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatcccagcg
acatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
atgaggctctgcacaaccactacacg cagaagagcctctccctgtctccgggtaaatgatctaga

25 Amino acid sequence: (SEQ ID NO: 131)

mdfqvqifslhisasviiargqivlsqspailsaspekvmttcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyycqqsfnpptfagtklelkdggsgsgsgsgssqaylqqsgaelvr
pgasvkmsckasgytftsynmhwwkqtrqglewigaiypngdtsynqkfkakatltvdksstaymqlssltsed
savvyfcarvvyysnsywyfdvwgtgtvtvssdpepkssdkthtspssapellggssvflfppkpkdtlmisrtevt
30 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvlvvlhqdwlngkeykckvsnkalpapiekt
iskakgqprepqvylppsrdeltnqvslclvkgyfypsdiavewesngqpennykttppvldsdgsttlyskltvdks
rwqqgnvfscsvmhealhnhytqkslsispkg

76. 2H7 scFv VH L11S (SSS-S)H P238SCH2 WCH3

35 Nucleotide sequence: (SEQ ID NO: 367)

aagcttgccgccatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcatactccaggggagaagggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtaccagcagaagccaggatcctccccaacacctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
40 attactgccagcagtgaggttttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggaggctcg
ggcgggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
cctcagtgaaagatgctctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctcacaatcagaagttcaagggaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
45 gactggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcaggag
cccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggatcgtcagttctctcttccc

cccaaaaccaagacaccctcatgatctcccgacccctgaggtcacatgcgtgggtgggacgtgagccacgaagac
 cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaagccgcgggaggagcagtac
 aacagcagctaccgtgtggtcagcgtctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
 5 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgcccccatcccggtgatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggttctatcccagcg
 acatcgccgtggagtgaggagagcaatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgacg
 gctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacacg cagaagagcctctcctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 368)

10 mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsgtsysltisrveadaatyccqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
 pgasvkmsckasgytfsynmhvkwqtrpqglewigaiypngdtsynqkfkkatltvdkssstaymqlssltssed
 savyfcarvvyysnsywyfdvwgtgtvtvssdpepkssdkthtspssapellggssvflfppkpkdtlmisrtpevt
 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
 15 skakgqprepqvylppsrdeitknqvsltclvkgfypsdiavewesngqpennyktppvldsdgsttlyskltvdksr
 wqqgnvfscsvmhealhnhytqkslslspgk

77. 2H7 scFv IgA WH WCH2 WCH3 + J Chain

Nucleotide sequence: 2H7 scFv IgA WH WCH2 WCH3 (SEQ ID NO: 61)

20 aagcttcccgcctatggatttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatctgtctgcattctccaggggagaaggtcacaatgacttgcagggccagctcaagtgaag
 ttacatgcactggtaccagcagaagccaggatcctccccaaaccctggattatgccccatccaacctggcttctggagtc
 cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
 attactgccagcagtgaggtttaaccaccacacgttcggtgctgggaccaagctggagctgaaagatggcgggtgctcg
 25 ggcgggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagctggtgagcctgggg
 cctcagtgaaagatgctctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaacgagacacctagaca
 gggcctggaatggattggagctattatccaggaaatggtgatacttctcacaatcagaagttcaagggcaaggccacactg
 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggcttatttctgtgcaa
 gagtggtgtactatagtaacttactggtacttctgatgtctggggcacagggaccacggtcaccgtctcttctgatcagcca
 30 gtccctcaactccactacccatctccctcaactccactacccatctccctcatgctgccacccccgactgtcactgca
 ccgaccggccctcaggagcctgctcttaggttcagaagcgtcctcagctgcacactgaccggcctgagagatgcctcag
 gtgtcaccttcacctggacgccctcaagtgggaagagcgtgtcaaggaccacctgaccgtgacctctgtggctgtaca
 gcgtgtccagtgctctgccgggtgtgccgagccatggaacctgggaagaccttacttgcactgctgcttaccctgagt
 ccaagacccccgtaaccgccacctctcaaaatccggaacacattccggcccagggtccacctgctgccgccgccgtc
 35 ggaggagctggccctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagccccaaaggatgtgctggttcgct
 ggctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccggcaggagcccagccaggggcacc
 accaccttcgctgtgaccagcactgctgcgtggcagccgaggactggaagaagggggacaccttctcctgatggtgg
 gccacgagggcctgccgtggccttcacacagaagaccatcgaccgcttggcgggtaaacccacctatgtaattgtgtct
 gttgtcatggcggaggtggacggcacctgctac tgataatctaga

40 +J Chain (SEQ ID NO: 65)

agatctcaagaagatgaaaggattgttctgttgacaacaaatgtaagtgtgcccggtacttccaggatcatccgttctcc
 gaagatcctaattgaggacattgtggagagaaacatccgaattattgttcctctgaacaacagggagaatatctctgatccac
 ctaccattgagaaccagattgtgtaccattgtgtgacctcagctgtaaaaaatgtgatcctacagaagtggagctggataa
 tcagatagtactgctaccagagcaatatctgtgatgaagacagtgtacagagacctgctacacttatgacagaacaaag
 45 tgctacacagctgtggtccactcgtatatggtggtgagacaaaatggtggaaacagccttaacccagatgcctgctatc
 ctgactaatctaga

Amino acid sequence: 2H7 scFv IgA WH WCH2 WCH3 (SEQ ID NO: 62)

mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyycqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaelvr
pgasvkmsckasgytftslnmhvkwqtrqglewigaiypngdtsynqkfkkgkatltvdksstaymqlssltssd
5 savyfcavvyysnsywyfdvwtgtttvssdqvpstpptpspctchprlsrlhrpaledlllgseailtcti
tglrdasgvtftwtpssgksavqgppdrdlcgcyssvslpgcaepwnhgktfictaaypesktpilatlsksntfrpev
hlpppseelalnelvtltclargfspkdvrvwlqgsqelprekyltwasrqepsqgtttfavtsilrvaaedwkkgdtdfsc
mvghaelplafktidrlagkpthvvnsvvmaevdgtcy
+J Chain (SEQ ID NO: 66)
10 rsqederivlvdnkckcaritsriirssedpnediverniriiplnnrenisdptsplrtrfvyhlsdlsckkkcdpteveldn
qivtatq snicdedsatetcytydmkcytavvplvyggetkmvetaltpdacypd

78. 2H7 scFv VH L11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 369)

15 aagcttgccgccatggatttcaagtgcagatttcagcttcctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattctccaggggagaagggtcacaatgacttcagggccagctcaagttaag
ttacatgactggtagcagcagaagccaggatcctccccaaaccctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctcacaatcagcagagtgagggtgaagatgctgccactt
attactgccagcagtgagtttaaccacccacgttcggtgctgggaccaagctggagctgaaagatggcgggtgctcg
20 ggcgggtggtgagctggaggaggtgggagctctcaggcttatctacagcagctctggggtgagtcggtgaggcctgggg
cctcagtgaaagatgctctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagaca
gggcttggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
actgtagacaaaactcctcagcagacgctacatgcagctcagcagcctgacatctgaagactctgcggcttattctgtgcaa
gagtgggtgtactatagtaactcttactgggtacttcgatgtctggggcagaggaccacggctaccgtctctctgatcaggag
25 cccaaatctctgacaaaactcacacatcccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcttccc
cccaaaacccaaggacaccctcatgatctccggaccctgaggtcacatgcgtgggtggtagctgagccacgaagac
cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtag
aacagcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactgggtgaatggcaaggagtacaagtgaaggt
ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
30 caccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatcccagcg
acatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacg
gtccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
atgaggctctgcacaaccactacacg cagaagagcctctcctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 370)

35 mdfqvqifsfllhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyycqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
pgasvkmsckasgytftslnmhvkwqtrqglewigaiypngdtsynqkfkkgkatltvdksstaymqlssltssd
savyfcavvyysnsywyfdvwtgtttvssdqepkssdkthtspssapellggpsvflfppkpkdtlmisrtpevt
cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
40 skakgcijrepqvtylppsrldeltnqvsltlcvkgfypsdiavewesngcipennykttppvldsdgsttlyskltvdk
rwqqgnvfscsvmhhealnhytqkslslspgk

79. 2H7 scFv VH L11S (CSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 371)

aagcttgccgccaatggattttcaagtgagattttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
 ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttgcagggccagctcaagtgtgag
 ttacatgactggtagcagaagccaggtacctccccaaacctggatttatgccccatcaacctggcttctggagtc
 cctgctcgttcagtgagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
 5 attactgccagcagtgaggatttaacccaccacgttcgggtgctgggaccaagctggagctgaaagatggcgggtggctcg
 ggcgggtggatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagtcggtgaggcctgggg
 cctcagtgaaagatgtcctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagaca
 gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagtcaagggaaggccacactg
 actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggctctatttctgtgcaa
 10 gagtgggtgactatagtaacttactggtacttctgctgctggggcacagggaccacggtcaccgtctcttctgacaggag
 cccaaatcttgtgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcttccc
 cccaaaacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtgggtgggacgtgagccacgaagac
 cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgaggagagcagtagc
 aacagcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
 15 ctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgta
 caccctgccccatccccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttetatcccagcg
 acatcgccgtggagtgaggagagcaatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgacg
 gctccttcttctctacagcaagctaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgc
 atgaggctctgcacaaccactacag cagaagagcctctcctgtctccgggtaaatgatctaga

20 Amino acid sequence: (SEQ ID NO: 372)

mdfqvqifslhisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypsnlasg
 vparfsgsgsytisrveadaatyccqqsfnpptfgagtklelkdggsgggsgggssqaylqqsgaesvr
 pgasvkmsckasgytfsynmhvkwqtrpglewigaiypngdtsynqkfkgtatlvdkssstaymqslsitsed
 savyfcarvvyysnywyfdvwgtgtvtvssdepkscdkhtspssapellggpsvflfppkpkdtlmisrtpevt
 25 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiecti
 skakgciprepqvytlppsrdeitknqvsltlcykgfypsdiavewesngqpennykttppvldsdgsttlyskhvdk
 rwqqgnvfscsvmhhealhnhytqkslslspgk

78. G8-1 scFv VHL11S (SCS-S)H WCH2 WCH3

30 Nucleotide sequence: SEQ ID NO: 373

gttgttaagcttgccgccaatggatccacagctcagttccttgggtgctgctgctgtggcttacaggtggcagatgtgacatc
 cagatgactcagctccagcctcctatctgcatcttgggagagactgtcaccatcacatgtcgaacaagtgaatgtttac
 agttatttggcttggtatcagcagaacagggaatctcctcagctcctggtctcttttcaaaaaccttagcagaaggtgtg
 ccatcaaggttcagtgagcagtgatcagggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagt
 35 atttctgtcaacatattccgataatccgggacgttcgggtggaggcaccgaactggagatcaaaggtggcgggtggctcggg
 cgggtgggtgggtcgggtggcggcggatcgtcagcgggtccagctgcagcagcttgagcctgagtcggaaaagcctggcgc
 ttactgaagatttctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatgaaaga
 gccttgagtggattggaatattgatccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgact
 gtacacaaatcctccagcacagcctacatgcagctcaagagtctgacatctgaggactctgagctctattactgtgcaagat
 40 cggtcggccctatggactactgggtgcaaggaacctcagtcaccgtctcttctgatcaggagcccaatcttctgacaaaa
 tcacacatgccaccgtcctcagcacctgaactcctgggtggaccgtcagcttctcttccccccaaaacccaaggacac
 cctcatgatctcccggaccctgaggtcacatgcgtgggtgggtgagcgtgagccacgaagacctgaggtcaagttcaact
 ggtacgtggacggcgtggaggtgcataatgccaagacaaagccgaggagagcagtagaacagcacgtaccgtggg
 tcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccca
 45 gccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatccccgg
 gatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccaagcagatcgccgtggagtggga
 gagcaatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagga

agctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccac
tacacgcagaagagcctctc cctgtctccgggtaaatgatctagag

Amino acid sequence: SEQ ID NO: 374

5 mvstaqlglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwtfgggteleikgggsgggsgggssavqlqqsgpesekpgas
vkisckasgysftgynmnwvkqnnngkslewnidpyyggtynrkfkgtatltvdksstaymqlksltsedsavy
ycarsvgpmdywgqgtsvtvssdqepkssdkthtccppsapellggpsvflfppkpkdtlmsrtpevtcvvvdvsh
edpevkfnwyvdgvevhnaktkpreeqynstyrsvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqip
10 repqvylppsrdeitknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsttlyskltvdksrwqqgnv
fscsvmhleahnhytqkslslspgk

79.. G28-1 scFv VHL11S (CCS-P)H WCH2 WCH3

Nucleotide sequence: SEQ ID NO: 375

15 gttgtaagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatc
cagatgactcagctccagcctcctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgttta
cagttatttggcttggtatcagcagaaacagggaaaatctcctcagctcctggtctctttgcaaaaacctagcagaaggtgt
gccatcaaggttcagtggtgagtcaggtcagcagacagtttctctgaagatcagcagcctgcagcctgaagattctggaagt
tatttctgtcaacatcattccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcggtggctcgg
gcggtgggtgggtcgggtggcgcggtatcgtcagcgggtccagctgcagcagcttggtgacctgagtcggaaaagcctggcg
20 cttcagtgaaagtctcgaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaaaga
gccttgagtggattggaatattgaccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgact
gtagacaaatcctccagcagcctacatgcagctcaagagtctgacatctgaggactctgcagcttattactgtcaagat
cggtcggccctatggactactgggtgcaaggaacctcagtcaccgtctcttctgatcaggagcccaatettgtgacaaaac
tcacacatgtccaccgtccccagcacctgaactcctgggtggaccgtcagcttctcttccccccaaaaccaaggacac
25 cctcatgatctcccgaccctgaggtcacatgcgtgggtgggtgacgtgagccacgaagacctgaggtcaagttcaact
ggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggagggagcagtacaacagcacgtaccgtgtg
gtcagcgtctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccc
agcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatccc
ggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccaagcgacatcgccgtggagtggg
30 agagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttcttcttacagc
aagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaacca
ctacacgcagaagagcctctc cctgtctccgggtaaatgatctaga

Amino acid sequence: SEQ ID NO: 376

35 mvstaqlglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwtfgggteleikgggsgggsgggssavqlqqsgpesekpgas
vkisckasgysftgynmnwvkqnnngkslewnidpyyggtynrkfkgtatltvdksstaymqlksltsedsavy
ycarsvgpmdywgqgtsvtvssdqepkssdkthtccppsapellggpsvflfppkpkdtlmsrtpevtcvvvdvsh
edpevkfnwyvdgvevhnaktkpreeqynstyrsvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqpr
40 epqvylppsrdeitknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsttlyskltvdksrwqqgnvf
scsvmhleahnhytqkslslspgk

80. G28-1 scFv VH L11S (SCC-P)H WCH2 WCH3

Nucleotide sequence: SEQ ID NO: 377

45 gttgtaagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatc
cagatgactcagctccagcctcctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgttta

cagttatttggcttggtatcagcagaaacagggaatatcctcagctcctggctctctttgcaaaaaccttagcagaaggtgt
gccatcaaggttcagtggtcagtggtatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagt
tatttctgtcaacatcattccgataatccgggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcgg
5 gcggtggtgggtcgggtggcggcgatcgtcagcggtccagctgcagcagcttgacctgagtcggaaaagcctggcg
cttcagtgaagatttctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatgaaaga
gccttgagtggattggaataflgaccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgact
gtagacaaatctccagcacagcctacatgcagctcaagagtctgacatctgaggactctgcagcttattactgtgcaagat
cggtcggccctatggactactgggtcaaggaacctcagtcaccgtctctctgatcaggagcccaaatctctgacaaaac
10 tcacacatgccaccgtgccagcacctgaactcctgggtggaccgtcagcttctctctcccccaaaaacccaaggacac
cctcatgatctcccggaccctgaggtcacatgcgtgggtgggtgagccacgaagacctgaggtcaagttcaact
ggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtg
gtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaagccctccc
agccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccg
15 ggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggttctatccaagcgacatcgccgtggagtggg
agagcaatgggcagccggagaacaactacaagaccacgctcccgtgctggactccgacggctccttctctctacagc
aagctcaccgtggacaagagcaggtggcagcaggggaacgtctctcatgctccgtgatgcatgaggctctgcacaacca
ctacacgcagaagagcctctc cctgtctccgggtaaatgatctaga

Amino acid sequence: SEQ ID NO: 378

mvstaqlfllllwtlgrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
20 sgsgsgtqfslkisslqpedsgsyfcqhhdnpwtfgggteleikggggsgggsgggssavqlqqsgpesekpgas
vkiskcasgysftgynmnwvkqnnkslewnidpnyggttynrktkgkatltvdkssstaymqlsltsedsavy
ycarsvgpmdywgqgtsvtvssdqepkssdkthtccppapellggspsvflfppkpdltmisrtpetvcvvvdvsh
edpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqpr
25 epqvytlppsrdeltknqvslclvkgfypsdiavewesngqpennykttpvldsdgsttlyskltvdksrwqqgnvf
scsvmhealhnhytqkslslspgk

81. G28-1 scFv VH L11S mIgE CH2 CH3 CH4

Nucleotide sequence: SEQ ID NO: 379

aagcttgccgccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
30 gactcagctccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatgtttacagtta
tttggcttggtatcagcagaaacagggaatatcctcagctcctggctcttttgcataaaccttagcagaaggtgtgccatc
aaggttcagtggtcagtggtatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
gtcaacatcattccgataatccgtggacgttcggtggaggcaccgaactggagatcaaaggtggcgggtggctcggcggt
35 ggtgggtcgggtggcggcgatcgtcagcggtccagctgcagcagcttgacctgagtcggaaaagcctggcgcttcag
tgaagatttctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatgaaagagcctt
gagtggattggaatatgtatccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgactgtaga
caaatctccagcacagcctacatgcagctcaagagtctgacatctgaggactctgcagcttattactgtgcaagatcggtc
ggccctatggactactgggtcaaggaacctcagtcaccgtctctctgatcacgttcacctgtcaacatcactgagccca
ccttgagctactccattcactcgcgaccccaatgcattccactccaccatccagctgtactgttcttattatggccacatct
40 aaatgatgtctctgtcagctggctaattggacgatcgggagataactgatacactgcacaaactgttctaataaggaggaa
ggcaactagcctctacctgcagtaactcaacatcactgagcagcaatggatgtctgaaagcaccttcacctgcaaggtc
acctccaaggcgtagactatttggccacactcggagatgccagatcatgagccacggggtgtgattacctacctgat
cccaccagccccctggacctgtatcaaacggtgctcccaagcttacctgtctggtggtggacctggaagcgagaaga
45 atgtcaatgtgactggaaccaagagaagaagacttcagctcagcatccagtggtactaagcaccacaataacgcca
caactagatcacctccatctgcctgtagtgtccaaggactggattgaaggctacggctatcagtgcatagtggaccacct
gattttccaagccattgtgcgttccatcacaagacccaggccagcgtcagccccgaggtatatgtgtccaccac
cagaggaggagagcgaggacaaacgcacactcacctgtttgatccagaacttcttcctgaggatattctgtgcagtggc

tgggggatggcaaaactgatctcaaacagccagcacagtaccacaacaccctgaaatccaatggctccaatcaaggcttct
tcatcttcagtcgcctagaggtcgccaagacactctggacacagagaaaacagttcacctgccaagtatccatgaggcac
ttcagaaaccaggaaactggagaaaacaatatccacaagccttggtaacacctccctccgtccatcctagtaatctagagg

Amino acid sequence: SEQ ID NO: 380

5 mvstaqflglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwfgggteleikggggsgggsgggssavqlqqsgpesekpgas
vkisckasgysftgynmnwvkqnnngkslewnidpyyggtynrkfkghkatltvdkssstaymqlksltsedsavy
ycarsvgpmdywgqgtsvtvssdhvrpvniteptlellhsscdpnafhstiqlycfiyghilndvsvswlmdreitdl
10 aqtvlikeegklascsklniteqqwmsestfickvtsqgvdyahtrrcpdheprgitylippspldlyqngapkltclv
vdleseknvnvtwnqekktvsasqwytkhhimattsitsilpvvakdwiegygyqcivdhdpfkipivrsitktpgq
rsapevyvfppeeesedkrtltcliqnttpedisvqwlgdgklisnshsttllksnsgnqgttisrlevaktlwtqrkq
ftcqihealqkprklektistslgntslrps

82. G28-1 scFv VH L11S mIgA WH WCH2 T4CH3

15 Nucleotide sequence: (SEQ ID NO: 381)

aagcttgccgcatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
gactcagtcaccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaatatgtttacagtta
tttggcttggtatcagcagaacagggaatatcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtgccatc
aaggttcagtgagcagtgatcaggcacacagtttctctgaagatcagcagcctgcagcctgaagattctggaagtatttct
20 gtcaacatcattccgataatccgtggacgttcgggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcgggt
gggtgggtcgggtggcggcgatcgtcagcgggtccagctgcagcagcttgacctgagtcggaagcctggcgcttcag
tgaagatttctgcaaggttctggttactcattcactggctacaatatgaactgggtgaagcagaataatgaaagagcctt
gagtggttggaatatgtatccttattatgggtgtactacctacaaccggaagtcaagggcgaaggccacattgactgtaga
caaactcctccagcacagcctacatgcagctcaagagtctgacatctgaggactctgcagctattactgtgaagatcggtc
25 ggccctatggactactggggtcaaggaacctcagtcaccgtctcttctgatcacatctgttctcctcactactcctcctccac
cttctgccagcccagcctgtcactgcagcggccagctcttgaggacctgctcctgggttcagatgccagcatcacatgtac
tctgaatggcctgagagatcctgagggagctgtcttcacctgggagccctccactgggaaggtgacgtgcagaagaaag
ctgtgcagaattcctcgggctgctacagtggtccagcgtcctgcctgggtgtgctgagcgtggaacagtggcgcacatt
caagtgcacagttaccatcctgagctgtgacaccttaactggcacaattgccaaagtcacagtgaacacctcccacccag
30 gtccacctgtaccgccgccgtcggaggagctggccctgaatgagctcgtgtccctgacatgcctgggtgcgagctttcaac
cctaaagaagtgtgtgctgcatggatgaggaatgaggagctgtccccagaaagctacctagtgttgagcccctaaag
gagccaggcgagggagccaccacctggtgacaagcgtgttgctgtatcagctgaaatctggaacagggtgacc
agtactctgcatggtgggccacgaggccttgcccatgaacttcacccagaagaccatcgaccgtctgtcgggtaaaccca
ccaatgtcagcgtgtctgtgatcatgtcagaggagattgataatctagat

35 Amino acid sequence: (SEQ ID NO: 382)

mvstaqflglillwltggrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslqpedsgsyfcqhhsdnptwfgggteleikggggsgggsgggssavqlqqsgpesekpgas
vkisckasgysftgynmnwvkqnnngkslewnidpyyggtynrkfkghkatltvdkssstaymqlksltsedsavy
ycarsvgpmdywgqgtsvtvssdhicspptppppscqpslsqrpaledllgsdasitctlngrldpegavftwepst
40 gkdavqkkavqnsqgcysvssvlpgaerwnsgasfctvthpesdltltgiakvtvntfppqvhllpppseelalnel
vsltclvrafnpkevlvrwlhgneelspesylvfeplkepegattylvtsvlrvsaeiwwkqgdqyscmvghealpmn
ftqktidrlsgkptnvsvsvimsegd

83. G28-1 scFv Vh L11S hIgE CH2 CH3 CH4

45 Nucleotide sequence: (SEQ ID NO: 383)

aagcttggcccatggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagat
 gactcagttccagcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaaaatgtttacagtta
 tttggcttggatcagcagaaacagggaaaatctcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtccatc
 aaggttcagtggcagtggtacaggcacacagttttcttgaagatcagcagcctgcagcctgaagattctggaagtatttct
 5 gtcaacatcattccgataatccgtggacgttcgggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcggt
 ggtgggtcgggtggcggcggtatcgtagcgggtccagctgcagcagcttgacgtcggaaaagcctggcgcttcag
 tgaagatttctgcaaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatggaaagagcctt
 gagtggattggaatattgatccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgactgtaga
 caaatcctccagcacagcctacatgcagctcaagagtctgacatctgaggactctgcagctctattactgtgaagatcggtc
 10 ggccctatggactactgggtcaaggaaacctcagtcaccgtctcttctgatcacgtctgctccagggaacttcacccgccc
 ccgtgaagatcttacagtcgtctcgtagcggcgggcggtcactccccccgacatccagctcctgtgcctcgtctctgggt
 acacccagggactatcaacatcacctgggtggaggacgggcaggtcatggacgtggactgtccaccgcctctaccacg
 caggagggtgagctggcctccacacaaagcgagctcaccctcagccagaagcactggctgtcagaccgcacctacacct
 gccaggtcacctatcaaggtcacacctttgaggacagcaccaagaagtgtgcagattccaacccgagaggggtgagcgc
 15 ctacctaaagccggccagcccgttcgacctgttaccgcaagtcgccacgatcacctgtctgggtgggtggacctggcacc
 cagcaaggggacctgaacctgacctgggtcccgccagtgaggaaacctgtgaaccactccaccagaaaggaggaga
 agcagcgcaatggcacgttaacctgcacgtccacctgccggtgggcacccgagactggatcgaggggggagacctacc
 agtcaggggtgacccaccccccacctgccagggccctcatgcggtccacgaccaagaccagcgcccgctgctgccc
 cggaagtctatgcgtttgcgacgccggagtggccggggagccgggacaagcgacccctgcctgcctgatccagaactt
 20 catgcctgaggacatctcgggtgcagtggtgcacaacgaggtgcagctcccggacggccggcacagcacgacgcagcc
 ccgcaagaccaagggctccggcttctctgcttccagccgctggaggtgaccagggccgaatgggagcagaaagatgag
 ttcatctgccgtgcagtcctgaggcagcgagccctcacagaccgtccagcgagcggtgtctgtaaatcccggtaaatga
 taatctaga

Amino acid sequence: (SEQ ID NO: 384)

25 mvstaqlfllllwtggrediqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
 sgsgsgtqfslkisslcpedsqsyfchhsdpwtfgggteleikgggsgggsgggssavqlqsgpeseekpga
 svkiskcasgysftgynmnwvkqnnkglewignidpyyggtytnrktkgtatltvdkssstaymqlksltsedsav
 yycarsvgpmdywgqgtsvtvssdhvcsrdftptvkilqsscdggghfpptiqlclvsgytpgtinitwledgqvm
 dvdlstasttqegelastqseltsqkhwlstdrtytcqvtyqghtfedstkkcadsnprgvsaylsrpspfdlfrksptitcl
 30 vvdlapskgtnvltwsrasgkpvnhstrikeekqrmgtltvstltpvgtrdwiegetyqcrvthphlpralmrsttktsgrpr
 aapevyafatpewpgrdkrtlaclqnfmpeidsvqwlhnevqlpdarhsttqprktksgsttvfsrlevtraeweqk
 deficravheaaspsqtvqravsvnpgk

84. G28-1 scFv VH L11S hIgA WH WCH2 T4CH3

35 Nucleotide sequence: (SEQ ID NO: 385)

atggtatccacagctcagttccttgggttgctgctgctgtggcttacaggtggcagatgtgacatccagatgactcagttcca
 gcctccctatctgcatctgtgggagagactgtcaccatcacatgtcgaacaagtgaaaatgtttacagttatttggcttggatc
 agcagaaacagggaaaatctcctcagctcctggctcttttgcaaaaaccttagcagaaggtgtccatcaaggttcagtg
 cagtggtacaggcacacagttttcttgaagatcagcagcctgcagcctgaagattctggaagttamctgtcaacatcattc
 40 cgataatccgtggacgttcgggtggaggcaccgaactggagatcaaaggtggcgggtggctcgggcggtggtgggtcgggt
 ggcgggcggtatcgtagcgggtccagctgcagcagcttgacgtcggaaaagcctggcgcttcagtgaagatttctg
 caaggcttctggttactcattcactggctacaatatgaactgggtgaagcagaataatgaaagagccttgagtggattgga
 aatattgatccttattatggtggtactacctacaaccggaagtcaagggaaggccacattgactgtagacaaatcctccag
 cacagcctacatgcagctcaagagtctgacatctgaggactctgcagctctattactgtgaagatcggtcggccctatggac
 45 tactgggggtcaaggaaacctcagtcaccgtctcttctgatcagccagttccctcaactccacctacccatctccctcaactcca
 cctacccatctccctcatgctgccaccccgactgtcactgcaccgaccggccctcgaggacctgctcttaggttcagaa
 gcgatccctcacgtgcacactgaccggcctgagagatgcctcaggtgtcaccttcacctggacgccctcaagtgggaagag

cgctgttcaaggaccacctgaccgtgacctctgtggctgctacagcgtgtccagtgtcctgccgggctgtgccgagccatg
gaaccatgggaagacct.tcacttgcaactgctgcctaccccagtgccaagacccccgtaaccgccaccctctcaaaatccg
gaaacacattccggcccaggtccacctgctgccgccgccgtcggaggagctggccctgaacgagctggtgacgctga
cgtgcctggcacgtggcttcagccccaggaatgtgctggctcgtggctgcaggggtcacaggagctgccccgcgagaa
5 gtacctgacttgggcatcccggcaggagcccagggcaccaccaccttcgctgtgaccagcatactgcgcgtggca
gccgaggactggaagaagggggacaccttctcctgcatggggtggccacgaggccctgccgctggccttcacacagaag
accatcgaccgcttggcgggtaaacccacccatgtcaatgtgtctgtgtcatggcggaggtggactgataatctaga

Amino acid sequence: (SEQ ID NO: 386)

10 mvstaqlflglllwtlgrcdiqmtqspaslsasvgetvtitcrtsenvysylawyqqkqgkspqllvsfaktlaegvpsrf
sgsgsgtqfslkisslpedsyfcqhhsdnptwfgggteleikgggsgggsgggssavqlqqsgpesekpgas
vkiskcasgysftgynmnwvkqnnngkslewnidpyygttynrkfkkgkatltvdksstaymqklsitsedsavy
ycarsvgpmdywgqg tsvtvssdqvpstptpsptpspschprlslhrraled

85. HD37 VL

15 Nucleotide sequence: (SEQ ID NO: 387)

aagcttgcggccatggagacagacacactcctgctatgggtgctgctgcttgggtccaggctccactggtgacattgtgc
tgaccaatctccagcttcttggctgtgtctctagggcagagggccaccatctcctgcaaggccagccaaagtgttgattat
gatggtgatagtattgaactggtaccaacagattccaggacagccacccaaactcctcatctatgatgcaccaatctagttt
ctgggatcccaccaggttagtggtgagtggtgctgggacagacttcaccctcaacatccatcctgtggagaaggtggatg
20 ctgcaacctatcactgtcagcaaaagtactgaggatccgtggacgttcggtggaggcaccaagctggaaatcaaaggtggc
ggtggctcgggcgggtggtgggtcgggtgg cggcgggagctcg

Amino acid sequence: (SEQ ID NO: 388)

metdtlllwylllwwpgstgdivltqspaslavslgqratisckasqsvdydgsylnwyqqipgqppklliidasnlvs
gipprfsgsgsgtdflnihpvekvdatyhcqqstedpwtfgggtkleikgggsgggsgggsgss

25

86. HD37 VH

Nucleotide sequence: (SEQ ID NO: 389)

gggagctcgcaggttcagctgcagcagctcggggctgagctggtgaggcctgggtcctcagtgaagatttctgcaaggc
ttctggctatgcattcagtagctactggatgaactgggtgaagcagagggcctggacagggcttgagtggattggacagattt
30 ggcttgagatggtgataactacaatggaaagttcaagggtaaagccactctgactgcagacgaatctccagcaca
gcctacatgcaactcagcagcctagcatctgaggactctcgggtctatttctgtgcaagacgggagactacgacggtaggc
cgttattactatgctatggactactg ggtcaaggaacctcagtcaccgtctcctctgatcag

Amino acid sequence: (SEQ ID NO: 390)

35 (gss)qvqlqqsgaelvrpgssvkiiskasyafssywmnvwkqrpqglewigqiwpdgdtnyngkfkkgkatlt
adesstaymqksslasedsavyfcarretttvgryyyamdywgqgtsvtvssdq

87. HD37 scFv

Nucleotide sequence: (SEQ ID NO: 391)

40 aagcttgcggccatggagacagacacactcctgctatgggtgctgctgcttgggtccaggctccactggtgacattgtgc
tgaccaatctccagcttcttggctgtgtctctagggcagagggccaccatctcctgcaaggccagccaaagtgttgattat
gatggtgatagtatttgaactggtaccaacagattccaggacagccacccaaactcctcatctatgatgcaccaatctagttt
ctgggatcccaccaggttagtggtgagtggtgctgggacagacttcaccctcaacatccatcctgtggagaaggtggatg

ctgcaacctatcactgtcagcaaagtactgaggatccgtggacgttcggtggaggcaccaagctggaaatcaaaggtggc
 ggtggctcgggcgggtggtgggtcgggtggcggcgatcgtcacagggtcagctgcagcagctctggggctgagctggtga
 ggcttgggtcctcagtgaagatttctgcaaggcttctggctatgcattcagtagctactggatgaactgggtgaagcagag
 gcctggacaggggtcttgattggacagatttggcctggagatggtgataactacaatggaaagtcaagggtaaa
 5 gccactctgactgcagacgaatcctccagcacagcctacatgcaactcagcagcctagcatctgaggactctcggtctatt
 tctgtgcaagacgggagactacgacggtagggcgttattactatgctatggactactgggggtcaaggaacctcagtcaccgt
 ctctctgatcag

Amino acid sequence: (SEQ ID NO: 392)

metdtlllwlwvllwvpstgdivltqspaslavslgqratiskasqsvdydgdsylnwyqqipgppklliidasnlvs
 10 gipprfsgsgsgtdfilnihpvekvdadyhcqqstcdpwtfgggtkleikgggsgggsgggsgssqvqlqqsgael
 vrpssvkiiskasgyafssywmnwvkqrpqglewigqiwpdgdtnyngkfkkgkatltadessstaymqlssla
 sedsavyfcarretttvgr yyyamdywgqgtsvtvssdq

88. HD37 VHL11S:

15 Nucleotide sequence: (SEQ ID NO: 393)

gggagctcgcagggtcagctgcagcagctctggggctgagtcggtgaggcctgggtcctcagtgaagatttctgcaaggc
 ttctggctatgcattcagtagctactggatgaactgggtgaagcagaggcctggacaggggtcttgattggtgacagatt
 ggcttggagatggtgataactacaatggaaagtcaagggtaaaagccactctgactgcagacgaatcctccagcaca
 gcctacatgcaactcagcagcctagcatctgaggactctcggtctatttctgtgcaagacgggagactacgacggtaggc
 20 cgttattactatgctatggactactg ggtcaaggaacctcagtcaccgtctctctgatcag

Amino acid sequence: (SEQ ID NO: 394)

(gss)qvqlqqsgaesvrpgssvkiiskasgyafssywmnwvkqrpqglewigqiwpdgdtnyngkfkkgkatl
 tadess staymqlsslasedsavyfcarretttvgryyyamdywgqgtsvtvssdq

25 89. HD37 scFv VHL11S:

Nucleotide sequence: (SEQ ID NO: 395)

Aagcttcccgcctaggagacagacacactcctgctatgggtgctgctgctctgggtccaggctccactggtgacattgtg
 ctgacccaatctccagcttcttggctgtgtcttagggcagagggccaccatctctgcaaggccagccaaagtgttgatta
 tgatggtgatagttattgaactggtaccaacagattccaggacagccacccaaactcctcatctatgatgcattcaatctagt
 30 tctgggatccacccaggttttagtggcagtggtctgggacagacttcacctcaacatccatcctgtggagaaggtggat
 gctgcaacctatcactgtcagcaaagtactgaggatccgtggacgttcggtggaggcaccaagctggaaatcaaaggtgg
 cggtggctcgggcgggtggtgggtcgggtggcggcgaggctgcaggttcagctgcagcagctctggggctgagtcggt
 gaggcctgggtcctcagtgaagatttctgcaaggcttctggctatgcattcagtagctactggatgaactgggtgaagcag
 aggcttggacaggggtcttgattggacagatttggcctggagatggtgataactacaatggaaagtcaaggggt
 35 aaagccactctgactgcagacgaatcctccagcacagcctacatgcaactcagcagcctagcatctgaggactctcggtc
 tatttctgtgcaagacgggagactacgacggtagggcgttattactatgctatggactactgggggtcaaggaacctcagtc
 ccgtctctctgatcag

Amino acid sequence: (SEQ ID NO: 396)

metdtlllwlwvllwvpstgdivltqspaslavslgqratiskasqsvdydgdsylnwyqqipgppklliidasnlvs
 40 gipprfsgsgsgtdftlnihpvekvdadyhcqqstcdpwtfgggtkleikgggsgggsgggsgssqvqlqqsgaes
 vrpssvkiiskasgyafssywmnwvkqrpqglewigqiwpdgdtnyngkfkkgkatltadessstaymqlssla
 sedsavyfcarretttvgr yyyamdywgqgtsvtvssdq

90. HD37 scFv IgAH hIgG1 WCH2 T4CH3

Nucleotide sequence (SEQ ID NO: 397)

aagcttgcgccatggagacagacacactcctgctatgggtgctgctgcttgggtccaggctccactggtgacattgtgc
5 tgaccaatctccagcttcttggctgtgtctctagggcagagggccaccatctcctgcaaggccagccaaagtgttgattat
gatggtgatagttatttgaactggtaccaacagattccaggacagccacccaaactcctcatctatgatgcaccaatctagttt
ctgggatcccccagggttagtggcagtggtgctgggacagacttcaccctcaacatccatcctgtggagaaggtggatg
ctgcaacctatcactgtcagcaaagtactgaggatccgtggacgttcgggtggaggcaccaagctggaaatcaaaggtggc
gggtggctcgggcgggtgggtgggtggcggcgaggctgcaggttcagctgcagcagctctggggctgagtcgggtg
aggcctgggtcctcagtgaaatttctgcaaggcttctggctatgcattcagtagctactggatgaactgggtgaagcaga
10 ggccctggacaggggtcttgattggacagatttggcctggagatgggtgataactaactacaatggaagttcaagggtta
aagccactctgactgcagacgaatcctccagcacagcctacatgcaactcagcagcctagcatctgaggactctgcggctt
atttctgtgcaagacgggagactacgacggtaggccgttattactatgctatggactactggggtaaggaaacctcagtcac
cgtctcctctgatcagccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgtgcc
ccccgactgtcactgcaccgaccggccctcaggacctgctcttaggttcagaagcgtcctcacgtgcacactgaccg
15 gcctgagagatgctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgtgttcaaggaccacctgaccgt
gacctctgtggctgctacagcgtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttgc
actgtgcctacccgagtcgaagacccccgtaaccgccacctctcaaatccggaaacacattccggcccaggtcca
cctgtgccgccgccgtcggaggagctggcctgaacgagctggtgacgctgacgtgcctggcacgtggcttcagcccc
aaggatgtgctggttcgctggctgcaggggtcacaggagctccccgcgagaagtacctgacttgggcatcccgccagg
20 agcccagccagggcaccaccaccttcgctgtgaccagcactgcgcgtggcagccgaggactggaagaagggggac
accttctcctgcatgggtgggccacgaggccctgccgtggccttcacacagaagaccatcgaccgcttggcgggtaaac
cacccatgtcaatgtgtctgttgcacatggcggaggtggactgataatctaga

Amino acid sequence (SEQ ID NO: 398)

metdtllllwvllwvpgstgdivltqspaslavslgqratisckasqsvdydgsylnwyqqipgqppklliidasnlvs
25 gipprfsgsgsgtdfilnihpvekvdadyhcqqstcdpwtfgggtkleikgggsggggsggggssqvlqqsgaes
vrpgssvkiackasgyafssywmnwvkqrpqglewigqiwpdgdtnyngkfkgkatltadesstaymqlssla
sedsavyfcarrettvgyryyamdywgqgtsvtvssdqpvpstptpsptpspschprlsihrpaledlllgse
ailtctltglrdasgvtftwtpssgksavqgppdrldgcysvssvlpgaepwnhgtftctaaypeskptlatlsksn
tfrpevhllpppseelalnelvtltclargfspkdvlrwlqgsqelprekyltwasrqpsqgtttfavtsilrvaedwkk
30 gdtfscmvghelpaftqktidrlagkpthvnnsv vmaevd

91. HD37 scFv (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 399)

aagcttgcgccatggagacagacacactcctgctatgggtgctgctgcttgggtccaggctccactggtgacattgtgc
35 tgaccaatctccagcttcttggctgtgtctctagggcagagggccaccatctcctgcaaggccagccaaagtgttgattat
gatggtgatagttatttgaactggtaccaacagattccaggacagccacccaaactcctcatctatgatgcaccaatctagttt
ctgggatcccccagggttagtggcagtggtgctgggacagacttcaccctcaacatccatcctgtggagaaggtggatg
ctgcaacctatcactgtcagcaaagtactgaggatccgtggacgttcgggtggaggcaccaagctggaaatcaaaggtggc
gggtggctcgggcgggtgggtgggtggcggcgatcgtcacaggttcagctgcagcagctctggggctgagctgggtga
40 ggccctgggtcctcagtgaaatttctgcaaggcttctggctatgcattcagtagctactggatgaactgggtgaagcagag
gcctggacaggggtcttgattggacagatttggcctggagatgggtgataactaactacaatggaagtcaagggtaaa
gccactctgactgcagacgaatcctccagcacagcctacatgcaactcagcagcctagcatctgaggactctgcggcttatt
tctgtgcaagacgggagactacgacggtaggccgttattactatgctatggactactggggtaaggaaacctcagtcaccgt
ctcctctgatcaggagcccaaatcttgcacaaactcacatccccaccgtcctcagcacctgaactcctgggtggaccg
45 tcagtcttctcttcccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtgggtggacg
tgagccacgaagacctgaggtcaagttcaactggtagctggacggcgtggaggtgcataatgccaagacaaagccgcg
ggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggag

tacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgag
aaccacaggtgtacaccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaagg
cttctatccaagcgacatcgccgtggagtgggagagcaatgggcagccgggagaacaactacaagaccacgcctcccggtg
ctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctca
5 tgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 400)

metdtlllwlwvllwvpgstgdivltqspaslavslgqratisckasqsvdydgdsylnwyqqipgqppklliydasnlvs
gipprfsgsgsgtdftlnihpvekvdadyhcqqstdepwtfgggtkleikgggsgggsgggsgssqvqlqqsgael
vrpgssvkiskasgyafssywmnwvkqrpqglewigqiwpdgdtnyngkflgkatltadesstaymqlssla
10 sedsavyfcarrettivgryyyamdywgqtsvtvssdqepkssdkthtsppsapellggpsvflfppkpkdtlmsr
tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpa
piektiskakgqprepvytlppsrdeitknqvsitclvkgyfypsdiavewesngqpennyktpvldsdgsttlysklt
vdksrwqqgnvfscsvmhcalnhhytqkslslspgk

15 92. HD37 scFv VH L11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 401)

aagcttcccgcctatggagacagacacactcctgctatgggtgctgctgctctgggttccaggctccactggtgacattgtgc
tgaccaatctccagcttcttggctgtgtcttagggcagagggccaccatctcctgcaaggccagccaaagtgttgattat
gatggatgatgtatttgaactggtagcaacagattccaggacagccacccaaactcctcatctatgatgcatccaatctagttt
20 ctgggatccaccaggttagtgagcagtggtctgggacagacttaccctcaacatccatcctgtggagaaggtggatg
ctgcaacctatcactgtcagcaaaagtactgaggatccgtggacgttcgggtggagggaccaagctggaaatcaaaggtggc
gggtggctcgggcggtgggtgggtcgggtggcgggagctcgcaggttcagctgcagcagctctggggctgagtcggtg
aggcctgggtcctcagtgagatttctgcaaggcttctggctatgcattcagtagctactggatgaactgggtgaagcaga
ggcctggacaggggtcttgagtggattggacagatttggcctggagatggatgataactacaatgaaagttaagggtga
25 aagccactctgactgcagacgaatcctccagcacagcctacatgcaactcagcagcctagcatctgaggactctgcggctt
atttctgtgcaagacgggagactacgacggttagggcgttattactatgctatggactactggggtaaggaacctcagtcac
cgtctcctctgatcaggagcccaatcttctgacaaaactcacatccccaccgtcctcagcacctgaactcctgggggga
ccgtcagctcttcttccccccaaaaccaaggacacccctcatgatctcccggaccctgaggtcaccatgcgtgggtggg
acgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagcc
30 gcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaag
gagtacaagtgaaggtctccaacaaagccctcccagccccatcgagaaaacaatctccaaagccaaagggcagcccc
gagaaccacaggtgtacaccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgtgtaa
aggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccgggagaacaactacaagaccacgcctcc
cgtgctgactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtct
35 tctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctag
a

Amino acid sequence: (SEQ ID NO: 402)

metdtlllwlwvllwvpgstgdivltqspaslavslgqratisckasqsvdydgdsylnwyqqipgqppklliydasnlvs
gipprfsgsgsgtdftlnihpvekvdadyhcqqstdepwtfgggtkleikgggsgggsgggsgssqvqlqqsgaes
40 vrpgssvkiskasgyafssywmnwvkqrpqglewigqiwpdgdtnyngkfkkgkatltadesstaymqlssla
sedsavyfcarrettivgryyyamdywgqtsvtvssdqepkssdkthtsppsapellggpsvflfppkpkdtlmsr
tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpa
piektiskakgqprepvytlppsrdeitknqvsitclvkgyfypsdiavewesngqpennyktpvldsdgsttlysklt
vdksrwqqgnvfscsvmhcalnhhytqkslslspgk

45

91. L6 VL

Nucleotide sequence: (SEQ ID NO: 403)

atggattttcaagtgcagattttcagcttcctgctaatacagtgcttcagtcataatgtccagaggacaaattgttctctcccagttc
ccagcaatcctgtctgcatctccaggggagaaggtcacattgacttgaggccagctcaagtgttaagttcatgaactggt
5 accagcagaagccaggatcctccccaaacctggatttatgccacatccaatttggcttctgagttccctggtcgcttcagt
ggcgagtgggtctgggacctcttactctctcgcaatcagcagagtgagggtgaagatgctgccacttattactgccagcagt
ggaatagtaaccactcacgttcggtgctgggaccaagctggagctgaaagagctctctggtggcgggtggctcgggcggt
ggtgggtcgggtggcggcgatct

Amino acid sequence: (SEQ ID NO: 404)

10 Mdfqvqifslhisasvimsrgqivlsqspailsaspgekvtltcrasssvsfmnwyqqkpgsspkpwiylatsnlasef
pgrfsgewsgtsyslairsveadaatyccqwnsnpltfagtklelkelsgggsgggsgggsgggsg

92. L6 VH

Nucleotide sequence: (SEQ ID NO: 405)

15 cagatccagttggtgcagctctggacctgagctgaagaagcctggagagacagtcagatctcctgcaaggcttctgggtat
accttcacaaactatggaatgaactgggtgaagcaggtccaggaaaggggttaaagtggatgggtggataaacacctac
actggacagccaacatagctgatgacttcaaggacgggttgccttctcttggaaacctctgcctacactgcctatttgcag
atcaacaacctcaaaaatgaggacatggctacatatttctgtcaagatttagctatggaactcacgttacgtgactactgg
ggccaaggcaccactctcacagtct cctctgatca

20 Amino acid sequence: (SEQ ID NO: 406)

qqlvqsgpelkkpgetvkiackasgytftnygmwvkwqapgkglkwmgwintytgqptyaddfkgrfafsletsa
ytaylqinnlknedmatyfcarsygnrsryadywgqgttltyssd

93. L6 scFv

25 Nucleotide sequence: (SEQ ID NO: 407)

atggattttcaagtgcagattttcagcttcctgctaatacagtgcttcagtcataatgtccagaggacaaattgttctctcccagttc
ccagcaatcctgtctgcatctccaggggagaaggtcacattgacttgaggccagctcaagtgttaagttcatgaactggt
accagcagaagccaggatcctccccaaacctggatttatgccacatccaatttggcttctgagttccctggtcgcttcagt
ggcgagtgggtctgggacctcttactctctcgcaatcagcagagtgagggtgaagatgctgccacttattactgccagcagt
30 ggaatagtaaccactcacgttcggtgctgggaccaagctggagctgaaagagctctctggtggcgggtggctcgggcggt
ggtgggtcgggtggcggcgatctctgcagatccagttggtgcagctctggacctgagctgaagaagcctggagagacag
tcaagatctcctgcaaggcttctgggtataccttcacaaactatggaatgaactgggtgaagcaggctccaggaaaggggtt
aaagtggatgggtggataaacctacactggacagccaacatagctgatgacttcaaggacgggttgccttctcttgg
aaacctctgcctacactgcctatttgcagatcaacaacctcaaaaatgaggacatggctacatatttctgtgcaagatttagct
35 atggaactcacgttacgtgactactggggccaaggcaccactctcacagtctctctctgatca

Amino acid sequence: (SEQ ID NO: 408)

mdfvqifslhisasvimsrgqivlsqspailsaspgekvtltcrasssvsfmnwyqqkpgsspkpwiylatsnlasef
pgrfsgewsgtsyslairsveadaatyccqwnsnpltfagtklelkelsgggsgggsgggsgggsglqqlvqsgpel
kkpgetvkiackasgytftnygmwvkwqapgkglkwmgwintytgqptyaddfkgrfafsletsaytaylqinnlk
40 edmatyfcarsygnrsry adywgqgttltyssd

94. L6 VHL11S

Nucleotide sequence; (SEQ ID NO: 409)

ctgcagatccagttggtgcagtcctggacctgagtcgaagaagcctggagagacagtcgaagatctcctgcaaggcttctggg
tataccttcacaaactatggaatgaactgggtgaagcaggctccaggaaagggttaaagtggatgggctggataaacacc
5 tacactggacagccaacatatgctgatgacttcaagggacgggttgccttctctttggaaacctctgcctacactgcctatttgc
agatcaacaacctcaaaaatgaggacatggctacatatttctgtgcaagatttagctatggttaactcacgttacgctgactact
ggggccaaggcaccactctcaca gtctcctctgatca

Amino acid sequence: (SEQ ID NO: 410)

10 qiqlvqsgpeskpkgetvkiskcasgytftnygmnnwvkqapgkglkwmgwintygtqptyaddfkgrfafsletsa
ytaylqi nnlknedmatyfcarfsygnrsryadywgqgttlvtssd

95. L6 VH L11S scFv

Nucleotide sequence; (SEQ ID NO: 411)

15 Aagcttgttattatggattttcaagtcagattttcagcttcctgctaatacagtcgttcagtcataatgtccagaggacaaattgtt
ctctcccagctccagcaatcctgtctgcatctccaggggagaagggtcacattgacttgcagggccagctcaagtgttaagttt
catgaactggtaccagcagaagccaggatcctcccccacccctggatttatgccacatccaatttggcttctgagttccctg
gtcgttcagtggtcagtggtctgggacctctactctctcgcaatcagcagagtgagggtgaagatgctgccacttattac
tccagcagtggaatagtaaccactcacgttcggtgctgggaccaagctggagctgaaagagctctctggtggcgggtgg
20 ctggggcggtggtgggtcgggtggcggtgatctctgcagatccagttggtgcagtcgttcgacctgagtcgaagaagcctg
gagagacagtcgaagatctcctgcaaggcttctgggtataccttcacaaactatggaatgaactgggtgaagcaggctccag
gaaagggttaaagtggatgggtggataaacctacactggacagccaacatatgctgatgacttcaagggacgggttg
ccttctctttggaaacctctgcctacactgcctatttgcagatcaacaacctcaaaaatgaggacatggctacatatttctgtgc
aagatttagctatggttaactcacgttacgctgactactggggccaaggcaccac, tctcacagtcctctctgatca

Amino acid sequence: (SEQ ID NO: 412)

25 mdfqvqifslhisasvimsrgqivisqpaiisaspgekvltlcrasssvsfmnnwyqqkpgsspkpwiysniasef
pgrfsgewsgtsyslaisrveaadaatyycqwnsnphfagtklelkelsgggsgggsgggslqiqlvqsgpes
kpkgetvkiskcasgytftnygmnnwvkqapgkglkwmgwintygtqptyaddfkgrfafsletsaytaylqinnikn
edmatyfcarfsygnrsry adywgqgttlvtssd

30 96. L6 Or L6 VHL11S scFv IgAH hIgGI WCH2 WCH3

Nucleotide sequence: (L6 VHL11S is shown) (SEQ ID NO: 413)

aagcttgttattatggattttcaagtcagattttcagcttcctgctaatacagtcgttcagtcataatgtccagaggacaaattgtt
ctctcccagctccagcaatcctgtctgcatctccaggggagaagggtcacattgacttgcagggccagctcaagtgttaagttt
catgaactggtaccagcagaagccaggatcctcccccacccctggatttatgccacatccaatttggcttctgagttccctg
35 gtcgttcagtggtcagtggtctgggacctctactctctcgcaatcagcagagtgagggtgaagatgctgccacttattac
tccagcagtggaatagtaaccactcacgttcggtgctgggaccaagctggagctgaaagagctctctggtggcgggtgg
ctcggggcggtggtgggtcgggtggcggtgatctctgcagatccagttggtgcagtcgttcgacctgagtcgaagaagcctg
gagagacagtcgaagatctcctgcaaggcttctgggtataccttcacaaactatggaatgaactgggtgaagcaggctccag
gaaagggttaaagtggatgggtggataaacctacactggacagccaacatatgctgatgacttcaagggacgggttg
40 ccttctctttggaaacctctgcctacactgcctatttgcagatcaacaacctcaaaaatgaggacatggctacatatttctgtgc
aagatttagctatggttaactcacgttacgctgactactggggccaaggcaccactctcacagtcctctctgatcagccagttc
cctcaactccacctaccccatctccctcaactccacctaccccatctccctcatcgccacctgaactcctggggggaccgtc
agtcttctcttccccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgctggtggtggacgt
gagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcg

5 ggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaagga
 gtacaagtgaaggtctccaacaagccctcccagccccatcgagaaaacaatctcaaagccaaagggcagccccga
 gaaccacaggtgtacacctgccccatccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtaag
 gcttctatcccagcgacatcgccgtggagtgaggagagcaatgggcagccggagaacaactacaagaccacgcctccgt
 15 gctggactccgacggctccttctctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaacgtctctc
 atgctccgtgatgatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 414)

10 mdfqvqifsfhlhisasvimsrgqivlsqspailsaspgekvtltcrasssvsfmnwyqqkpgsspkpwiya
 pgrfsgewsgtsyslaisrveadaatyycqwnsnpltfagtklelkelsgggsgggsgggslqqlvqsgpes
 kkpgetvkiskasytftnygmnnwvkqapgkglkwmgwintygtqptyaddfkgrfafsletsaytaylqinnlk
 edmatyfcarsygnryadywgqgttlvssdqvpstptpspstptpspsscappellggsplfpkpkdltlmisrt
 pevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvitvqhwdwlngkeykckvsnkalpap
 iektiskakgqprepvytlppsrdeltknqvslclvkgfypsdiavewesngcipennyktppldsdgsttlysklt
 vdkswqqgnvfscsvmhleahnhytqkslslspgk

15

97. L6 scFv VHL11S (SSS-S)H WCH2 WCH3

Nucleotide sequence: (SEQ ID NO: 415)

20 aagcttgttattgattttcaagtgcagatttcagcttctgctaatacagtgcttcagtcataatgtccagaggacaaattgtt
 ctctcccagtcctcagcaatctgtctgcatctccaggggagaaggtcacattgacttcagggccagctcaagtgttaagttt
 catgaactggtagcagcagaagccaggtaccccccaccctggattatgccacatccaatttgcttctgagttccctg
 gtcgcttcagtgaggagtggtctgggaccttactctctcgaatcagcagagtgagggtgaagatgctgccacttattac
 tgccagcagtggaatagtaaccactcacgttcggtgctgggaccaagctggagctgaaagagctctctggtggcggtgg
 ctggggcggtggtgggtcgggtggcgggatctctgcagatccagttggtgcagcttgacctgagtcgaagaagcctg
 gagagacagtcgaatctcctgcaaggctctgggtataccttcacaaactatggaatgaactgggtgaagcaggctccag
 25 gaaagggtttaaagtggatgggtggataaacactacactggacagccaacatagctgatgactcaagggacgggttg
 ctttctcttgaaacctctgcctacactgcctatttgcatcaacaacctcaaaaatgaggacatggctacatattctgtgc
 aagatttagctatggaactcacgttacgctgactactggggccaaggcaccactctcacagctcctctgatcaggagccc
 aaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggaccgtcagcttctcttcccccc
 aaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccct
 30 gaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgaggagagcagtagaac
 agcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctc
 caacaaagccctcccagccccatcgagaaaacaatctcaaagccaaagggcagccccgagaaccacaggtgtacac
 cctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtaaaaggttctatcccagcgaca
 tcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggct
 35 ctttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctctcatgctccgtgatgatg
 aggtctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 416)

40 mdfqvqifsfhlhisasvimsrgqivlsqspailsaspgekvtltcrasssvsfmnwyqqkpgsspkpwiya
 pgrfsgewsgtsyslaisrveadaatyycqwnsnpltfagtklelkelsgggsgggsgggslqqlvqsgpes
 kkpgetvkiskasytftnygmnnwvkqapgkglkwmgwintygtqptyaddfkgrfafsletsaytaylqinnlk
 nedmatyfcarsygnryadywgqgttlvssdqepkssdkthtspssapellggsplfpkpkdltlmisrtpevt
 cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvitvqhwdwlngkeykckvsnkalpapiecti
 skakgqprepvytlppsrdeltknqvslclvkgfypsdiavewesngqpennyktppldsdgsttlyskltvdksw
 wqqgnvfscsvmhleahnhytqkslslspgk

45

98. P238 CH2

Nucleotide sequence: (SEQ ID NO: 91)

5 cctgaactcctgggggatcgtcagtccttcttccccccaaaaccaaggacaccctcatgatctcccggaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctcc
aaagccaaa

Amino acid sequence: (SEQ ID NO:92)

10 pellggssvflfppkpkdtl misrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvl
hqdwlngkeykckvsnkalpapiektiskak

99. P238 CH2 WCH3

a. P238S CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 417)

15 cctgaactcctgggggatcgtcagtccttcttccccccaaaaccaaggacaccctcatgatctcccggaccctgag
gtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcacca
ggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctcc
aaagccaaaaggcagccccgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtc
20 agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagagcaatgggcagccggagaca
actacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggt
ggcagcaggggaacgtcttctcatgctcgtgatgcatgaggtctgcacaaccactacacgcagaagagcctctccctgt
ctccgggtaaatgatctaga

Amino acid sequence: (SEQ ID NO: 418)

25 pellggssvflfppkpkdtl misrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvl
hqdwlngkeykckvsnkalpapiektiskakgqprepvytlppsrdeltknqvsitclvkfypsdiavewesngq
pennykttppvldsdgsttlyskltvdksrwqqgnvfscsvmhealhnhytqkslsispk

b. (SSS-S)H P238S CH2 WCH3

30 Nucleotide sequence: (SEQ ID NO: 419)

tgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctgggggatcgtcagtc
cttctcttccccccaaaaccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgag
ccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcggga
ggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtac
35 aagtgcaaggtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaaggcagccccgagaa
ccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggctt
ctatcccagcgacatcgccgtggagtgaggagagcaatgggcagccggagagaacaactacaagaccacgcctcccgtgct
ggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatg
ctccgtgatgcatgaggtctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

40 Amino acid sequence: (SEQ ID NO: 420)

dqepkssdkhtspssapellggssvflfppkpkdtl misrtpevtcvvvdvshedpevkfnwyvdgvevhnaktk
preeqynstyrvvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvytlppsrdeltknqvsitclv

kgfypsdiavewesngqpennykttppvldsdgsttiyskltdksrwqqgnvfscsvmhealhnhytqkslspsgk*

100. a.CD16-6 low (ED) + NL + (SSS-S)H P238S CH2 WCH3

5 Nucleotide sequence: (SEQ ID NO: 421)

ggagcccaaatcttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggatcgtagcttctctc
tccccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtgggtggtagctgagccacga
agacctgaggtcaagttcaactggtacgtggacggcggtggaggtgcataatgccaagacaaaagccgaggagga
gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga
10 aggtctccaacaaagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacagg
tgtacacctgcccccatcccggtatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatccca
gcgacatcgccgtggagtgaggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctggtgactccg
acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
tgcataagggtctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctaga

15 Amino acid sequence: (SEQ ID NO: 422)

mwqlllptallllvsagmrtdlpkavvflepqwyrvlekdsvtlkcqgayspednstqwfhnslissqassyfidaat
vddsgelyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtylqngkgrkyfhnsdfyi
pkatlkdsysyfcrglvsgknvssetvnititqgladqepkssdkthtspssapellggssvflfppkpkdtlmisrtpv
tcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpapiect
20 iskakgpprepqvylppsrldtknqvslclvkgfypsdiavewesngqpennykttppvldsdgsttiyskltdks
rwqqgnvfscsvmhealhnhytqkslspsgk

101. b.CD16-6 low (ED) + HE4LP + (SSS-S)H P238S CH2 WCH3

Nucleotide sequence: (SEQ ID NO: 423)

25 aagcttgccgccatgcctgcttgcctaggcccgctagccgccgcctcctcctcagcctgctgctgttcggcttcaccct
agtctcaggcaccgggtgcaatgcggactgaagatctccaaaggctgtggtgttcctggagcctcaatggtacagggtgct
cgagaaggacagtgtgactctgaagtgccaggaggcctactccctgaggacaattccacacagtgggttcacaatgaga
gcctcatctcaagccaggcctcgagctacttcatgagctgccacagtcgacgacagtggagagtacaggtgccagaca
aacctctccaccctcagtgaccgggtgcagctagaatccatcggctggctgttgcctcaggccctcggtgggtgttca
30 aggaggaagaccctattcacctgaggtgtcacagctggaagaacactgctctgcataaggtcacatattacagaatggca
aaggcaggaagtattttcatcataattctgacttctacattccaaaagccacactcaaagacagcggtcctacttctgcagg
gggcttgtgggagtaaaaatgtgtcttcagagactgtgaacatcaccatcactcaaggtttggtgatcaggagcccaaat
ttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggatcgtagcttctcttccccccaaaac
ccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtggtagctgagccacgaagacctgaggt
35 caagttcaactggtacgtggacggcggtggaggtgcataatgccaagacaaaagccgaggagagcagtacaacagcac
gtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaaca
aagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgc
ccccatcccggtatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatcccagcgacatgcc
gtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacggctcctct
40 tctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggt
ctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctagaaa

Amino acid sequence: (SEQ ID NO: 424)

mpacrlgplaaaallslfgrtlvsgtgamrtdlpkavvflepqwyrvlekdsvtlkcqgayspednstqwfhnslis
sqassyfidaatvddsgelyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtylqngkgr
45 kyfhnsdfyipkatlkdsysyfcrglvsgknvssetvnititqgladqepkssdkthtspssapellggssvflfppkpk

kdtlmisrtpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckv
snkalpapiektiskakgqprepvytlppsrdeltknqvsltclvkgyfypsdiavewesngqpennykttpvldsdg
sttlyskltvdksrwqqgnvfscsvmh ealhnhytqkslsispk

5 **102. CD16-9 high (ED) (SSS-S)H P238S CH2 CH3**

a. CD16-9 high (ED)NL + (SSS-S)H P238S CH2 CH3

Nucleotide sequence: (SEQ ID NO: 425)

gttgtaagcttgccgcatgtggcagctgctcctccaaactgctctgctacttctagtttcagctggcatgaggactgaagat
ctcccaaaggctgtggtgttctggagcctcaatggtacagggtgctcgagaaggacagtgtagctctgaagtgccaggg
10 agcctactcccctgaggacaattccacacagtggtttcacaatgagagcctcatcgaagccaggcctcagctacttcattg
acgctgccacagtcgacgacagtgagagtagcaggtgccagacaaacctctccaccctcagtgacccgggtgcagctaga
agtccatcgcggtgggtgttgcctcaggccccctcggtgggtgttcaaggaggaagaccctattcacctgaggtgtcacag
ctggaagaacactgctctgcataagggtcacatattacagaatggcaaaggcaggaagtatttcataataattctgacttcta
cattccaaaaggcacactcaagacagcggctcctacttctgcagggggcttttgggagtaaaaatgtgtcttcagagact
15 gtgaacatcaccatcactcaaggtttggctgatcaggagcccaaatcttctgacaaaactcacacatccccaccgtcctcag
cacctgaactcctggggggatcgtcagcttctcttccccccaaaaccaaggacaccctcatgatctccggaccctga
ggtcacatgcgtgggtgggtgacgtgagccacgaagaccctgaggtcaagttcaactgggtacgtggacggcgtggaggtg
cataatgccaaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcacc
aggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccagccccatcgagaaaacaatctc
20 caaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccggtatgagctgaccaagaaccaggt
cagcctgacctgctgtgtaaggttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccgggagaac
aactacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcag
gtggcagcaggggaacgtcttctcatgtccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccct
gtctccgggtaa atgatctagaaa

25 **Amino acid sequence: (SEQ ID NO: 426)**

mwqlllptallllvsagmrtdlpkavvflepqwyrvlekdsvtlkcqgayspednstqwfhnslissqassyfidaat
vddsgelyrcqtnlstldpvqlvhighwillqaprwvfkeedpihlrchswkntalhkvtylqngkgrkyfhnsdfyi
pkatlkdsyfcrglfgsknvsssetvnititqgladqepkssdkthtspssapellggssvflfppkpkdtlmisrtpev
tcvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpapiekt
30 iskakgqprepvytlppsrdeltknqvsltclvkgyfypsdiavewesngqpennykttpvldsdgsttlyskltvdks
rwqqgnvfscsvmh ealhnhytqkslsispk

CD16-9 high (ED) + HE4LP + hIgG1 (SSS-S)H P238S CH2 CH3

Nucleotide sequence: (SEQ ID NO: 427)

aagcttgccgcatgcctgcttgcgcctaggccccgtagccgccgcctcctcctcagcctgctgctgttcggcttcaccct
agtctcaggcaccgggtgcaatgcggactgaagatctcccaaaggctgtggtgttctggagcctcaatggtacagggtgct
cgagaaggacagtgtagctctgaagtgccaggagcctactcccctgaggacaattccacacagtggtttcacaatgaga
gcctcatcgaagccaggcctcagctacttcattgacgtgccacagtcgacgacagtgagagtagcaggtgccagaca
aacctctccaccctcagtgacccgggtgcagctagaagtcacatcggtgggtgttgcctcaggccccctcggtgggtgttca
40 aggaggaagaccctattcacctgaggtgtcacagctggaagaacactgctctgcataagggtcacatattacagaatggca
aaggcaggaagtattttcatcataattctgacttctacattccaaaaggcacactcaagacagcggctcctacttctgcagg
gggcttttgggagtaaaaatgtgtcttcagagactgtgaacatcaccatcactcaaggtttggctgatcaggagcccaatc
ttctgacaaaactcacacatccccaccgtcctcagcacctgaactcctggggggatcgtcagcttctccttccccccaaaac
ccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtgggtggacgtgagccacgaagaccctgaggt
45 caagtcaactggtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtagaacagcac

gtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaaca
aagccctcccagcccccatcgagaaaacaatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgc
ccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtaaaggcttctatcccagcgacatcgcc
gtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttct
5 tctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctatgctccgtgatgatgaggt
ctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaatgatctagaaa

Amino acid sequence: (SEQ ID NO: 428)

mpacrlgplaaallslfllvsgtgamrtedlpkavvflepqwyrvlekdsvtlkcqgayspednstqwfhnslis
sqassyfidaatvddsgeyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtlylqngkgr
10 kyfhnsdfyipkatlkdsgsyfcrglvgsknvssetvnititqgladqepkssdkthtspssapellggssvtlfpkpk
dtlmsirtpvctcvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvs
nkalpapiektiskakgpprepvytlpprsdeltnqvslctlvkgfypsdiavewesngqpennyktpvldsdgs
ttlyskltvdksrwqqgnvfscsvmhe alhnhytqkslslspgk

15 103. a. CD16 ED low (native leader)

Nucleotide sequence: (SEQ ID NO: 429)

aagcttgccgccatgtggcagctgctcctcccaactgctctgctacttctagtttcagctggcatgcggactgaagatctccc
aaaggctgtggtgttctggagcctcaatggtacaggggtgctcgagaaggacagtgtagcttgaagtccaggagcct
actcccctgaggacaattccacacagtggtttcacaatgagagcctcatctcaagccaggcctcgagctacttcattgacgct
20 gccacagtcgacgacagtggagagtacaggtgccagacaaacctctccaccctcagtgacccgggtgcagctagaagtcc
atatcggctggctgttgcctcaggccccctcgggtgggtgttcaaggaggaagaccctattcacctgaggtgtcacagctgga
agaacactgctctgcataaggtcacatatttacagaatggcaaaggcaggaagtatttcatcataattctgacttctacattcc
aaaagccacactcaaagacagcggctcctacttctgcagggggcctgttgggagtaaaaatgtgtcttcagagactgtgaa
catcaccatcactcaaggttggctgatcaaaa

25 Amino acid sequence: (SEQ ID NO: 430)

mwqlllptalllvsagmrtdlpkavvflepqwyrvlekdsvtlkcqgayspednstqwfhnslissqassyfidaat
vddsgeyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtlylqngkgrkyfhnsdfyi
pkatlkdsgsyf crglvgsknvssetvnititqgladq

30 b. CD16 ED low (HE4 leader)

Nucleotide sequence (SEQ ID NO: 431)

aagcttgccgccatgcctgcttgcctaggccccgctagccgccgccctcctcctcagcctgctgctgttcgggttcaccct
agtctcaggcaccgggtgcaatgcggactgaagatctccaaaggctgtggtgttctggagcctcaatggtacaggggtgct
cgagaaggacagtgtagcttgaagtgccaggagcctactcccctgaggacaattccacacagtggtttcacaatgaga
35 gcctcatctcaagccaggcctcgagctacttcattgacgctgccacagtcgacgacagtggagagtacaggtgccagaca
aacctctccaccctcagtgacccgggtgcagctagaagtccatcctggctggctgttgcctcaggccccctcgggtgggtgttca
aggaggaagaccctattcacctgaggtgtcacagctggaagaacactgctctgcataaggtcacatatttacagaatggca
aaggcaggaagtatttcatcataattctgacttctacattccaaaagccacactcaaagacagcggctcctacttctgcagg
gggcttgttgggagtaaaaatgtgtcttcagagactgtgaacatc accatcactcaaggttggctgatcaaa

40 Amino acid sequence (SEQ ID NO: 432)

mpacrlgplaaallslfllvsgtgamrtedlpkavvflepqwyrvlekdsvtlkcqgayspednstqwfhnslis
sqassyfidaatvddsgeyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtlylqngkgr
kyfhnsdfyipkatlkdsgsyfcrglvgsknvssetvnititqgladq

104. a. CD16 ED high (native leader)

Nucleotide sequence: (SEQ ID NO: 433)

gttgtaagcttgcggccatgtggcagctgctcctcccaactgctctgctacttctagtttcagctggcatgcggactgaagat
ctcccaaaggctgtggtgttctggagcctcaatggtacagggtgctcgagaaggacagtgtgactctgaagtgccaggg
5 agcctactcccctgaggacaattccacacagtgggttcacaatgagagcctcatctcaagccaggcctcgagctacttcattg
acgctgccacagtcgacgacagtggagagtacagggtgccagacaaacctctccaccctcagtgacccgggtgcagctaga
agtccatcgcggctgggtgttgcctcaggccccctcggtgggtgttcaaggaggaagaccctattcacctgaggtgtcacag
ctggaagaacactgctctgcataaggtcacatatttacagaatggcaaaggcaggaagtattttcatcataattctgacttcta
cattccaaaagccacactcaaagacagcggctcctacttctgcagggggcttttgggagtaaaaatgtgtcttcagagact
10 gtgaacatcaccatcactcaaggtttggctgatcaaa

Amino acid sequence: (SEQ ID NO: 434)

mwqlllptallllvsagmrtdlpkavvflepqwyrvlekdsvtlkcgayspednstqwfhnslissqassyfidaat
vddsgeyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtylqngkgrkyfhnsdfyi
pkatlkdsysyfcrglfgsknvssetvnititqgladq

15 b. CD16 ED high (HE4 leader)

Nucleotide sequence: (SEQ ID NO: 435)

aagcttgcggccatgcctgcttgcgctaggcccgtagccgcccgcctcctcagcctgctgctgttcggcttcacct
agtctcaggcaccgggtgcaatgcggactgaagatctcccaaaggctgtggtgttcctggagcctcaatggtacagggtgct
cgagaaggacagtgtgactctgaagtgccaggaggcctactcccctgaggacaattccacacagtgggttcacaatgaga
20 gcctcatctcaagccaggcctcgagctacttcattgacgctgccacagtcgacgacagtggagagtacagggtgccagaca
aacctctccacctcagtgacccgggtgcagctagaagtccatcgcggctggctgttgcctcaggccccctcggtgggtgttca
aggaggaagacctattcacctgaggtgtcacagctggaagaacactgctctgcataaggtcacatatttacagaatggca
aaggcaggaagtattttcatcataattctgacttctacattccaaaagccacactcaaagacagcggctcctacttctgcagg
gggcttttgggagtaaaaatgtgtcttcagagactgtgaacatca ccatcactcaaggtttggctgatcaaa

25 Amino acid sequence: (SEQ ID NO: 436)

mpacrlgplaaalllllfgftlvsgtgamrtdlpkavvflepqwyrvlekdsvtlkcgayspednstqwfhnslis
sqassyfidaatvddsgeyrcqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtylqngkgr
kyfhnsdfyipkatlkdsysyfcrglfgsknvssetvnititqgladq

30 105. 2e12 scFv (SSS-S)H P238S CH2 WCH3-hCD80TM/CT

Nucleotide sequence: (SEQ ID NO: 437)

aagcttatggattttcaagtcagattttcagcttctgtaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttcttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaaggtgtgaatatta
tgtcacaagtttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
35 tctggggctcctgccaggttagtggcagtggtgtcgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaatgtaggaaggttccttgagcgttcgggtggaggcaccacagctggaaatcaaacgggggtg
cggtggctcggggcggagggtgggtcgggtggcgcggtatcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatgtgtaaactgggttcgccagcctc
caggaaagggtctggagtggctgggaatgatatgggtgatggaagcacagactataattcagctctcaaatccagactga
40 gcatcacaaggacaactccaagagccaagtttcttaaaaaatgaacagcttgcaactgatgacacagccagatactactg
tgccagagatgggtatagtaactttcattactatgttatggactactgggggtcaaggaacctcagtcaccgtctcctcagatctg
gagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggatcgtcagcttctcctct
tcccccaaaaacccaagacaccctcatgatctcccgaccctgaggtcacatgcgtgggtgggtggacgtgagccacga
agacctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaagacaaaagccgaggaggagca

gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgc
 aggtctccaacaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
 tgtacacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgctggtcaaaggcttctatcca
 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 5 acggctccttctctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
 tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttgaacctgetcc
 catcctggggcattacctaatactcagtaaatggaattttgtgatgtgctgacctactgctttgccccaaagatgcagaga
 gagaaggaggaatgagagattgagaaggga aagtgtacgccctgtataaatcgat

Amino acid sequence: (SEQ ID NO: 438)

10 mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqqkpgppkllisaasn
 vesgvparfsgsgsgtdfslnihpveeddiamefcqqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
 gpglvapsqslsitctvsgfsltygyvnwvrrppgkglewlgmiwgdgstdynsalksrslitkdnksqvflkmnsl
 qtddtaryycardgysnthyyvmdywgqgtsvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmsirt
 pevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
 15 iektiskakgciprepqvytlppsrdeltknqvsltclvkgyfypsdiavewesngcqpennyktpvldsdgsttlyskl
 tvdksrwqqgnvfscsvmhleahnhytqkslslspgkadpsnllpswaitlisvngifviccltycfaprerrrrmerlrr
 esvrpv

106. 10A8 scFv (SSS-S)H P238SCH2 WCH3-hCD80TM/CT

20 Nucleotide sequence: (SEQ ID NO: 439)

aagcttatggattttcaagtgcagatttcagcttctgctaatactgcttcagtcataatgtccagaggagtcgacatccaga
 tgacacagctcccatcctcactgtctgcatctctgggaggcgaagtcaccatcacttgcaaggcaagccaagacattaagaa
 gtatataggttggtaccaacacaagcctggaaaagggtccaggctgctcatatattacacatctacattacagccaggcatcc
 catcaagggttcagtggaagtgggtctgggagagattatccctcagcatcagaaacctggagcctgaagatattgcaactat
 25 tattgtcaacagtatgataatcttcattgacgttcggctcggggacaaaagttgaaataaaacggggtggcgggtggctcgg
 gcgggtgggtgggtcgggtggcggcgatctgatgtacagcttcaggagtcaggacctggcctcgtgaaaccttctcagtcctc
 tctctcaccctgctctgactggctactccatcaccagtggtttctactggaactggatccgacagttccgggaaacaac
 tggatggatgggccaataagccacgacggttaggaataactacaacccatctctcataaatgaatctccatcactcgtga
 cacatctaagaaccagtttttctgaagttgagttctgtgactactgaggacacagctacataattctgtgcaagacactacggt
 30 agtagcggagctatggactactggggtcaaggaacctcagtcaccgtctcctctgatctggagocccaaatctctgacaaaa
 ctacacatccccaccgtccccagcacctgaactcctggggggatcgtcagcttctcctcttcccccaaaacccaaggaca
 cctcatgatctccggaccctgaggcacatgcgtggtggtagctgagccacgaagacctgaggtcaagtcaact
 ggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtagaacgacgacgtaccgtgtg
 gtcagcgtcctcaccgtcctgcaccaggactgggtgaatggcaaggagtacaagtgaaggtctccaacaagccctccc
 35 agcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccg
 ggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatcccagcgacatcgccgtggagtggg
 agagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagc
 aagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctgcacaacca
 ctacacgcagaagagcctctccctgtctccgggtaaagcgatccttgaacctgtcccatcctgggccattaccttaatt
 40 cagtaaatggaattttgtgatgtgctgacctactgctttgccccaaagatgcagagagagaaggaggaatgagagatt
 gagaagggaagtgtacgccctgtataaatcgat

Amino acid sequence: (SEQ ID NO: 440)

mdfvqvifsfllhisasvimsrgvdiqmtspsslsaslggkvitckasqdikkyigwyqhkpkgkprlliyttstkipg
 iprsfsgsgsgrdyslsimlepediatyyccqydnlpitfgsgtkleikrgggsgggsgggsgsdvqlqesgpglvkp
 45 sqslsltsvtygysitgfywnwirqfpgriklewmghishdgrnynpslinrisitrdtsknqtlklssvttedatfyfa
 rhygssgamdywgqgtsvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmsirtpevtcvvvdvshe
 dpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqpre

pqvylppsrldeltnqvsitclvkgfypsdiavewesngqpennykttpvldsdgsttlyskltvdksrwqqgnv fs
csvmhealhnhytqkslspsgkadsnlp swa itlisvngifviccltycfaprcrerrrnerlrresvrp

107. 40.2.36 scFv (SSS-P)H P238SCH2 WCH3-hCD80TM/CT

5 Nucleotide sequence:

Amino acid sequence:

108. 2H7 scFv VHL11S (SSS-P)H P238SCH2CH3-hCD80TM/CT

10 Nucleotide sequence: (SEQ ID NO: 441)

aagcttgccgcatggatttcaagtgcagatttcagcttctgctaatactgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcattccaggggagaaggtcacaatgacttcagggccagctcaagtgaag
ttacatgcactggtagcagaagccaggatcctcccccacccctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgaggagctgaagatgctgccactt
15 attactgccagcagtgagtttaacccacccacgttcggtgctgggaccaagctggagctgaaagatggcggaggctcg
ggcggagggtgatctggaggagtgaggagctctcaggcttatctacagcagctctggggctgagtcggtaggcctgggg
cctcagtgaaatgtctgcaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctattatccaggaaatggtgatacttctacaatcagaagttcaagggcaaggccacactg
actgtagacaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggcttatttctgtgcaa
20 gagggtgtactatagtaacttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatctggag
cccaaatcttctgacaaaactcacatccccaccgtcccagcacctgaactcctggggggatcgtcagcttctcttccc
cccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtggtagcgtgagccacgaagac
cctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtac
aacagcacgtaccgtgtggtcagcgtcctaccgtctgcaccaggactgggtgaatggcaaggagtacaagtgaaggt
25 ctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgta
caccctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctgtaaaaggcttctatcccagcg
acatcgcctggtgagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacg
gtccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctctcatgctccgtgatgc
atgaggctctgcacaaccactacacgcagaagagcctctcctgtctcgggtaaaagcgatccttgaacctgctcccat
30 cctgggccattaccttaatctcagtaaatggaattttgtgatatgctgcctgacctactgcttgccccaagatgcagagaga
gaaggaggaatgagagattgagaagggaagtgtacgccctgtataaatcgat

Amino acid sequence: (SEQ ID NO: 442)

mdfqvqifslhisasviiargqivlsqpailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwi yapsnlasg
vparfsgsgsgtsysltisrveadaatyccqqsfnpptfagtklelkdggsgggsgggssqaylqqsgaesvr
35 pgasvkmsckasgytfts ynmhvwkqtpqglewigaiypngdtsynqkfk gkatltvdksstaymqlsslt sed
savvyfcarvvyysnsywyfdvwtgtvtvssdlepkssdkthtsp spapellggssvflfpkp kdfilmisrtp evt
cvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyr vsvltvlhqdwlngkeykckvsnkalpapi ekti
skakgqprepvylppsrldeltnqvsitclvkgfypsdiavewesngqpennykttpvldsdgsttlyskltvdksr
wqqgnvfscsvmhealhnhytqkslspsgkadsnlp swaitlisvngifviccltycfaprcrerrrnerlrresvrp
40 v

109. G19-4 scFv (SSS-P)H P238SCH2 WCH3-hCD80TM/CT

Nucleotide sequence: (SEQ ID NO: 443)

aagcttatggattttcaagtgcagattttcagcttctgctaatactgcttcagtcataatgtccagaggagtcgacatccaga
 tgacacagactacatcctcctgtctgcctctctgggagacagagtcaccatcagttgcagggcaagtcaggacattcgca
 attatttaaactggatcagcagaaaccagatggaactgttaaactcctgatctactacacatcaagattacactcaggagtc
 catcaagggtcagtggcagtggtctggaacagattattctcaccattgccaacctgcaaccagaagatattgccacttact
 5 tttgccaacagggtaatacgttccgtggacgttcggtggagggaccaaactggttaaccaaacgggagctcgggtggcggt
 ggctcggggcggtgggtgggtcgggtggcggggagctatcagatgaggtccagctgcaacagctcggacctgaactggtga
 agcctggagcttcaatgtcctgcaaggcctctggttactcattcactggctacatcgtgaactggctgaagcagagccatgg
 aaagaaccttgagtggattggacttattaatccatacaaaaggtcttactacctaaccagaaattcaagggaaggccacat
 taactgtagacaagtcatccagcacagcctacatggagctcctcagctgacatctgaagactctgcagcttattactgtgca
 10 agatctgggtactatggtgactcggactggtacttcgatgtctggggcgagggaccacggtcaccgtctcctctgatctgg
 agcccaaatcttctgacaaaactcacacaagcccaccgagcccagcacctgaactcctggggggatcgtcagcttctccttt
 cccccaaaacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtgggtggacgtgagccacgaa
 gacctgaggtcaagttcaactggtacgtggacggcggtggaggtgcataatgccaagacaaagccgcgggaggagcag
 tacaacagcacgtaccgtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgc
 15 ggtctccaacaaagccctcccagccccatcgagaaaacctctccaaagccaaagggcagccccgagaaccacaggt
 gtacacctgcccccatcccggtgatgagctgaccaagaaccaggtcagcctgacctgcctggctcaaaaggcttctatccca
 gcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 acggctccttctcctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtctctcatgctccgtga
 tgcagtaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaagcggatcctcgaacctgtccc
 20 catcctgggccattaccttaatctcagtaaagtgaattttgtgatgtgctgcctgacctactgctttgccccagatgcagaga
 gagaaggaggaatgagagattgagaagggaagtgtacgccctgtataatcgatactcga

Amino acid sequence: (SEQ ID NO: 444)

mdfqvqifsfllisasvmsrgvdiqmtqtsslsaslgdrvtiscrasqdirnylnwyqqkpdgtvkliyytsrlhsgvp
 srfsgsgsgtdysltianlqpediatyfcqqgntlpwtfgggklvtrkrelgggsgggsgggsggsidevqlqqsgpvlv
 25 kpgasmsckasgysftgyivnwlkqshgknlewiglinpykglttynqkflgkatltvdkssstaymellsitsedsav
 yycarsgyygdswyfdvwgagttvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmisrtpvctvv
 vdvshedpevkfnwyvdgvevhnaatkpreeqynstyrvsvltvlhqdwlingkeykckvsnkalpapiektiska
 kgqprepqvytlppsrdeltnqvsltlvkgfypsdiavewesngqpennyktpplvdsdgsfflyskltvdksrwq
 qgnvfscsvmhealnhhytqkslsispkpadpsnllpswaitlisvngifviccltycfaprerrrrerlrrresvrpv
 30

110. 2e12 scFv (SSS-P)H WCH2 WCH3-hCD80TM/CT

Nucleotide sequence: (SEQ ID NO: 51)

aagcttatggattttcaagtgcagattttcagcttctgctaatactgcttcagtcataatgtccagaggagtcgacattgtgct
 caccatcctcagcttcttggctgtgtctctaggtcagagagccaccatcctcgcagagccagtgaaggtgtgaatatta
 35 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
 tctggggtcctgccaggttagtggtcagtggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
 ttgcaatgtattctgtcagcaaaagtaggaaggttccttgacgttcggtggaggaccaagctggaaatcaaacgggtgg
 cggtggctcgggcggaggtgggtcgggtggcgggcggatctcaggtgcagctgaaggagtcaggacctggcctggtgg
 cgccctcacagagcctgtccatcacatgcaccgtctcaggggttcattaaccggctatggtgtaaactgggttcgccagcct
 40 ccaggaaagggtctggagtgggtgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactg
 agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgcaaaactgatgacacagccagatactact
 gtgccagagatggttatagtaacttctactatgttatggactactgggggtcaaggaacctcagtcaccgtctcctcagatct
 ggagcccaaatcttctgacaaaactcacatccccaccgtccccagcacctgaactcctggggggaccgtcagcttctcct
 ctccccccaaaacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtgggtggacgtgagccacg
 45 aagacctgaggtcaagttcaactggtacgtggacggcggtggaggtgcataatgccaagacaaagccgcgggaggagc
 agtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgc
 aaggctctccaacaaagccctcccagccccatcgagaaaacctctccaaagccaaagggcagccccgagaaccacag

gtgtacaccctgccccatcccggtgatgagctgaccaagaaccaggtcagcctgacctgcctggtaaaggcttctatccc
 agcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactcc
 gacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtg
 atgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacctgctc
 5 ccatcctggggccattacctaatactcagtaaatggaattttgtgatatgctgcctgacctactgctttgccccaaagtgcagag
 agagaaggaggaatgagagattgagaagggaagtgtacgcctgtataaatcgat

Amino acid sequence (SEQ ID NO:52)

mdfqvqifsfllisasvimsrgvdivltqspaslavlsgqratiscrasesveyyvtslmqwyyqqkpgppkllisaasn
 vesgvparfsgsgsgtdfslnihpveeddiamyfcqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
 10 gpglvapsqslsitctvsgfsltygvnwvrrppgkglewlgmiwgdgstdynsalksrslitkdnsksqvflkmnsl
 qtddtaryycardgysnfhyvmdywgqgtsvtvssdlepkssdkthtspsspapellggpsvflfppkpkdtlmlsr
 tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpa
 piektiskakgqprepvytlppsrdeitknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsfflyskl
 tvdksrwqqgnvfscsvmhleahnhytqkslslspgkadsnllpswaitlisvngifviccltycfaprerrrrerlrr
 15 esvrpv

111. 2e12 scFv IgA WH WCH2 T4 CH3-hCD80TM/CT

Nucleotide sequence: (SEQ ID NO: 445)

aagcttatggatttcaagtgcagatttcagcttctgctaatactgcttcagtcataatgtccagaggagtcgacattgtgct
 20 caccaatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaagtggtgaatatta
 tgtcacaagttaatactgagtggtaccaacagaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
 tctggggctccctgccaggttagtggtcagtggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
 ttgcaatgtatttctgtcagcaaataggaaggttccttgacgttcgggtggaggcaccagctggaaatcaaacgggggtgg
 cggtggctcggggcggaggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
 25 cgccctcacagagcctgtccatcacatgcaccgtctcaggggtctcattaaccggctatggtgtaaactgggttcgccagcct
 ccaggaaagggtctggagtgggtgggaatgatatgggtgatggaagcacagactataattcagctctcaatccagactg
 agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgaaactgatgacacagccagataactact
 gtgccagagatggttatagtaactttcattactatgttatggactactgggggtcaaggaaacctcagtcaccgtctcctcagatc
 agccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtca
 30 ctgcaccgaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgc
 ctgaggtgtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccacctgacctgacctctgtggctg
 ctacagcgtgtccagtgtcctgccgggtgtgccgagccatggaacctgggaagaccttcacttgactgctgcctacc
 cgagtccaagaccccgtaccgccacctctcaaaatccggaacacattccggcccaggtccacctgctgccgccg
 ccgtcggaggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtgctgg
 35 ttgctggctgcaggggtcacaggagctgccccgcgagaagtacgtgacttgggcatcccgaggagcccagccagg
 gcaccaccaccttcgtgtgaccagcatactgcgcgtggcagccgaggactggaagaagggggacaccttctcctgcat
 ggtgggccacgaggccctgccgtggccttcacacagaagaccatcgaccgcttggcgggtaaaccacccatgtcaat
 gtgtctgttgcattggcggaggtggacgcggatccttcgaacaacctgtcccacctgggccattacctaatactcagtaaa
 tggaaattttgtgatatgctgcctgacctactgctttgccccaaagtgcagagagagaaggaggaatgagagattgagaagg
 40 gaaagtgtacgcctgtataaatcgatac

Amino acid sequence: (SEQ ID NO: 446)

mdfqvqifsfllisasvimsrgvdivltqspaslavlsgqratiscrasesveyyvtslmqwyyqqkpgppkllisaasn
 vesgvparfsgsgsgtdfslnihpveeddiamyfcqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
 45 gpglvapsqslsitctvsgfsltygvnwvrrppgkglewlgmiwgdgstdynsalksrslitkdnsksqvflkmnsl
 qtddtaryycardgysnfhyvmdywgqgtsvtvssdqvpstptpspstptpspcchprlslhrpaledlllgse
 ailtctltglrdasgvftwtpssgksavqppdrldlccysvssvlpgaepwnhgtftctaaypesktpilatlsksn
 tfrpevhllpppseelalnelvtlclargfspkdvlvrlqgsqelprekyltwasrqepsqgtttfavtsilrvaedwkk

gdtfscmvghleaplaftqktidrlagkpthvvnsvvmaevdadpsnnllpswaitlisvngifvicltycfaprerr
rnerlrresvrpv

112. 2e12scFv IgE WCH2WCH3WCH4-hCD80TM/CT

5 Nucleotide sequence: (SEQ ID NO: 181)

aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
caccacatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaagtggtgaatatta
tgtcacaagtttaatgcagtggtaccaacagaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
tctgggggtccctgccaggttagtggtcagtggtctgggacagacttcagcctcaacatccatctgtggaggaggatgata
10 ttgcaatgtatttctgtcagcaagtaggaaggttcttgacgttcggtggaggcaccagctggaaatcaaacgggggtgg
cggtggctcggggcggagggtgggtcggtggcggggatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcct
ccaggaaagggtctggagtggctgggaatgatatgggtgatggaagcacagactataattcagctctcaatccagactg
agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgcaaactgatgacacagccagatactact
15 gtgccagagatggttatagtaactttcattactatgttatggactactggggtaaggaacctcagtcacctctcctcagatc
acgtctgtccagggtcactcaccccgcccaccgtgaagatcttacagtcgtcctgcgacggcgggcagcttcccccg
accatccagctcctgtgcctcgtctctgggtacacccagggtactatcaacatcacctgggtggaggacgggcaggtcatg
gacgtggacttgtccaccgcctctaccacgcaggagggtgagctggcctccacacaaagcgagctcacctcagccaga
agcactggctgtcagaccgcacctacacctgccaggtcacctatcaagggtcacaccttgaggacagcaccaagaagtgt
20 gcagattccaacccgagaggggtgagcgcctacctaagccggccagcccgttcgacgtgttcacccgaagtcgcca
cgatcacctgtctggtgggtggacctggcaccagcaaggggacgtgaacctgacctgggtcccgggccagtggaagc
ctgtgaaccactccaccagaaaggaggagaagcagcgcaatggcacgttaaccgtcacgtccaccctgccggtgggca
cccgagactggatcgagggggagacctaccagtgcagggtgacccacccccacctgccaggggcctcatgcggtcca
cgaccaagaccagcggcccgcgtgctgccccggaagtctatgcgttgcgacgccggagtggccggggagccgggac
25 aagcgcaccctcgcctgcctgatccagaacttcagcctgaggacatctcgtgcagtggtgcacaacgaggtgcagct
cccggacggccggcacagcagcagcagccccgcaagaccaagggtccggcttctcgtcttcagccgcttgagggt
gaccaggggccgaatgggagcagaaagatgagttcatctgccgtgcagtcctatgaggcagcgagccctcacagaccgt
ccagcgagcgggtgtctgtaaatcccggtaaagcggatccttcgaagctcccatcctgggccattacctaactcagtaaag
gaattttgtgatgtgctgcctgacctactgcttggcccaagatgcagagagagaaggaggaatgagagattgagaagg
30 aagtgtagccctgtataatcgata

Amino acid sequence: (SEQ ID NO: 182)

mdfqvqifsfllisasvmsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqqkpgcippkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiamefcqqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
gpglvapsqslsitctvsgfsltygvnwvrpppgkglewlgmiwgdgstdynsalksrsltkdsksqvflkmnsl
35 qtddtaryycardgysnfhyvmdywgqgtsvtvssdhvcsrdftptvkilqsscdggghfpptiqlclvsgytpgt
initwledgqvmvdvlstasttqegelastqseltsqkhwlsdrtytcqvtyqghtfedstkkcadsnprgvsaylsrps
pfdlfrksptitclvvdlapskgtvnltwsrasgkpvnhstrkeekqrngltvtstlpvgtrdwiegetyqcrvthphlpr
almrsttktsgpaaapevyafatpewpgsrdrkrlacliqnfmpevisvqwlhnevqlpdarhsttqprktksgffvfs
rlevtraeweqkdeficravheaaspsqtvqravsvnpgkadpsklpswaitlisvngifvicltycfaprerrnerl
40 rresvrpv

113. mFADD-TM/CT

MFADD scscFv

Nucleotide sequence: (SEQ ID NO: 201)

gttgatgacccctgaacatggacccattcctggctgctgctgactcgtgtccggcagcctgtcgggcaacgatctgatgg
agctcaagttcttgtgccgcgagcgcgtgagcaaacgaaagctggagcgcgtgcagagtggcctggacctgttcacggt
gctgctggagcagaacgacctggagcgcgggcacaccgggctgctgcgcgagttgctggcctcgtgcgccgacacg
atctactgcagcgcctggacgacttcgagggcggggacggcgaccgctgcgccccggggaggcagatctgcaggtg
5 gcatitgacattgtgtgtgacaatgtggggagagactggaaaagactggccccgcgagctgaagggtgtctgaggccaagat
ggatgggattgaggagaagtacccccgaagtctgagtgcgggtaaggagagctgaaagtctggaagaatgctgag
aagaagaacgcctcgggtggccgactggtaaggcgcgtgcggacctgcaggtgaatctggtggctgacctggtggaa
gaagcccaggaatctgtgagcaagagtgaagaatgtccccagtactaagggttcaactgtgtcttctcagaacaccct
gactcgagatcgat

10 Amino acid sequence: (SEQ ID NO: 202)

Vvdpsnmdpflvllhslsgslsgndlmelkflcrervskrlervqsgldlftvllqndlerghtglrellaslrhldlqlrl
ddfeagtataappgeadlqvafdivcdnvgrdwkrlarelkvseakmdgieekyprslservreslkvwknaekkna
svaglvkalrtcrlnlvadlveeaqesvsksenmspvlrdstvsssetp

15 114. 2e12 scFV (SSS-P)H P238S CH2 WCH3-mFADD-TM/CT

Nucleotide sequence: (SEQ ID NO: 447)

aagcttatggattttcaagtgcagattttcagcttctgctaactcagtgcttcagtcataatgtccagaggagtcgacattgtgct
caccacatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgtgaatatta
tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
20 tctggggtccctgccaggttagtgagcagtggtctgggacagacttcagcctcaacatccatctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaatgtaggaaggttcttggacgttcgggtggaggcaccagctggaaatcaaacggggtgg
cggtggctcgggagggtgggtcgggtggcgggatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaactgggttcgccagcct
ccaggaaagggtctggagtgggtgggaatgataggggtgatggaagcacagactataatcagctctcaatccagactg
25 agcatcaccaaggacaactccaagagccaagttttcttaaaatgaacagctctgcaaaactgatgacacagccagatactact
gtgccagagatggttatagtaactttcattactatgttatggactactggggtcaaggaaacctcagtcaccgtctcctcagatct
ggagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctgggtggatcgtcagcttctc
tccccccaaaacccaaggacacctcatgatctcccgacccctgaggtcacatgcgtgggtgggtggacgtgagccacga
agacctgaggtcaagttcaactgggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
30 gtacaacagcacgtaccgtgtggtcagcgtctcaccgtcctgcaccaggactgggtgaatggcaaggagtacaagtgc
agggtccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
tgtacacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctatcca
gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccagcctcccgtgctggactccg
acggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
35 tgcagtagggtctgcacaaccactacacgcagaagagccttccctgtctccgggtaaagcggtaccttcgaacatggacc
cattcctgggtgctgctgcactcgtgtccggcagcctgtcgggcaacgatctgatggagctcaagttcttgtccgcgagcg
cgtgagcaaacgaaagctggagcgcgtgcagagtggcctggacctgttcacggtgctgctggagcagaacgacctgga
gcgcgggcacaccgggtgctgcgcgagttgctggcctcgtgcgccgacacgatctactgcagcgcctggacgacttc
gaggcggggacggcgaccgctgcgccccggggaggcagatctgcaggtggcatttgacattgtgtgtgacaatgtg
40 gggagagactggaaaagactggccccgcgagctgaagggtgtctgaggccaagatggatgggattgaggagaagtacccc
cgaagtctgagtgcgggtaaggagagctgaaagtctggaagaatgctgagaagaagaacgcctcgggtggccgga
ctggtcaaggcgcgtgcggacctgcaggtgaatctggtggctgacctggtggaagaagcccaggaatctgtgagcaaga
gtgagaatatgtccccagtactaagggttcaactgtgtcttctcagaacaccctgactcgagatcgat

Amino acid sequence: (SEQ ID NO: 448)

45 mdfqvqifslisavmsrgvdivltqspaslavlsgqratiscrasesveyyvtslmqwyyqqkpgppkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiameyfcqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
pgplvapsqslsitctvsgfsltgygvnwvrrppgkglewlgmiwdgstdynsalksrslitkdnsksqvflkmnsl

5 qtdtdaryycardgysnfhyvmdywgqgtsvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmisrt
 pevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
 iektiskakgqprepvytlppsrdeltnqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflysklt
 vdksrwqqgnvfscsvmhleahnhytqkslspsgkadsnmdpflvllhslsgslsgndlmelkflcrervskrkler
 vqsgldlftvllleqndlerghtgllrellaslrhdllqrliddfeagtataappgeadlqvafdivcdnvgrdwkrlarelkvs
 eakmdgieekyprslservreslkvwknaekknasvaglvkalrtcrlnlvadlveeaqesvsksenmspvlrdstvss
 setp

115. 2e12 scFv (SSS-P)H WCH2WCH3-mFADD-TM/CT

10 Nucleotide sequence: (SEQ ID NO: 449)

aagcttatggattttcaagtcagattttcagcttctgctaatacagtcgcttcagtcataatgtccagaggagtcgacattgtgct
 caccacatctccagcttcttggctgctctaggtcagagagccaccatctcctgcagagccagtgaaagtgtgaatattatgt
 cacaagtttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaatct
 15 ggggtccctgccaggttttagtggcagtggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgatatt
 gcaatgtatttctgtcagcaaagtaggaaggttccttgacgttcggtggaggaccaagctggaaatcaaacggggtggc
 ggtggctcgggcggaggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtggc
 gccctcacagagcctgtccatcacatgcaccgtctcaggggttctcattaaccggctatggtgtaaactgggttcgccagcctc
 caggaaagggtctggagtgggtgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactga
 gcatcaccaaggacaactccaagagccaagtttcttaaaaaatgaacagcttgcaactgatgacacagccagatactactg
 20 tgccagagatggttatagtaactttcattactatgttatggactactggggtcaaggaacctcagtcaccgtctcctcagatctg
 gagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctgggtggaccgtcagtttctctt
 cccccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacgaa
 gacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcag
 tacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaa
 25 ggtctccaacaaagccctcccagccccatcgagaaaaceatctccaaagccaaagggcagccccgagaaccacaggt
 gtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgacctggtcaaggcttctatccca
 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 acggctccttctcctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
 tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacatggacc
 30 cattctggtgctgctgcactcgtgtccggcagcctgtcgggcaacgatctgatggagctcaagttcttgtgccgcgagcg
 cgtgagcaaacgaaagctggagcgcgtgcagagtggcctggacctgttcacggtgctgctggagcagaacgacctgga
 gcgcgggcacaccgggtgctgcgcgagttgctggcctcgtcgcgccgacacgatctactgcagcgcctggacgacttc
 gaggcggggacggcgaccgtgcgccccgggggaggcagatctgcaggtggcatttgacattgtgtgtgacaatgtg
 gggagagactggaaaagactggcccgagctgaaggtgctgaggccaagatggatgggattgaggagaagtacccc
 35 cgaagtctgagtgagcgggtaaggagagctgaaagtctggaagaatgctgagaagaagaacgcctcgggtggccgga
 ctggtcaaggcgtgcggacctgcaggtgaatctggtggtgacctggtggaagaagcccaggaatctgtgagcaaga
 gtgagaatatgtccccagtactaagggttaactgtgtcttctcagaaacaccctgactcgagatcgat

Amino acid sequence: (SEQ ID NO: 450)

40 mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyqqkpgqppkllisaasn
 vesgvparfsgsgsgtdfslnihpveeddiamyfcqqsrkvpwtfggggtkleikrgggsgggsgggsgvqlkes
 gpglvapsqslsitctvsgfsltygvnwvrpppgkglewlgmwgdgstdynsalksrslsitkdnsksqvflkmnsl
 qtdtdaryycardgysnfhyvmdywgqgtsvtvssdlepkssdkthtspspapellggpsvflfppkpkdtlmisr
 tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpa
 45 piektiskakgqprepvytlppsrdeltnqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskl
 tvdksrwqqgnvfscsvmhleahnhytqkslspsgkadsnmdpflvllhslsgslsgndlmelkflcrervskrkle
 rvqsgldlftvllleqndlerghtgllrellaslrhdllqrliddfeagtataappgeadlqvafdivcdnvgrdwkrlarelkv

seakmdgieekyprslservreslkvwknaekknasvaglvkalrtcrlnlvadlveeaquesvsksenmspvldstvs
ssetp

116. mcasp3-TM/CT

5 Nucleotide sequence: (SEQ ID NO: 451)

Ggaccttcgaacatggagaacaacaaacctcagtgattcaaaatccattaataattttgaagtaaagaccatacatggg
agcaagtcagtggaactctgggatctatctggacagtagttacaaaatggattatcctgaaatgggcatatgcataataataat
aataagaacttcataagagcactggaatgtcatctcgtctggtagcgatggacgcagccaacctcagagagacattc
atgggcctgaaatacaagtcaggaataaaaatgatcttactcgtgaagacattttggaattaatggatagtggttctaaggaa
10 gatcatagcaaaaggagcagcttctgtgtgtgattctaagccatggatgaaggggtcatttatgggacaaatgggcctgt
tgaactgaaaaagttagactagcttctcagaggcgactactgccggagtctgactggaaagccgaaactcttcattcag
gcctgccgggtgacggagctggactgtggcattgagacagacagtgaggactgatgaggagatggcttgcagaagatac
cgggtggaggtgacttctgtatgcttactctacagcacctggtagtacttctggagaaattcaaaggacgggtcgtgttca
tccagtcctttgcagcatgctgaagctgtacgcgcacaagctagaatttatgcacattctcactcgcgttaacaggaaggtg
15 gcaacggaattcagtccttctccctggactccacttccacgaaagaaacagatcccgtgtattgtgcatgctcacgaa
agaactgtactttatcactagctcagatcgatg

Amino acid sequence: (SEQ ID NO: 452)

dpsnmennktsvdsksinnfevktihgsksvdsgiyldssykmdypemgiciiiinnknfhkstgmssrsrgtdvdaa
nlretfmglkyqvmkndltredilelmdsvskedhskrssfvcvilshgdegviygtngpvelkkltstfrrgdyersltg
20 kpklfiiqacrgteldcgietdsgtdeemacqkipveadflyaystapgyyswrnskdgswfiqslcsmklyahklf
mhilrtvrnkratefesfsl dstfhakkqipcivsmkelyfyh

117. 2e12 scFv (SSS-P)H.WCH2WCH3-mcasp3-TM/CT

Nucleotide sequence: (SEQ ID NO: 453)

aagcttatggattttcaagtcagatttcagcttctgtaatcagtgcttcagtcataatgtccagaggagtcgacattgtgct
caccatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgtgaatatta
tgacacaagttaatgcagtggtaccaacagaaaccaggacagccaccaaacctctcatctctgctgcaccaacgtagaa
tctggggtccctgccaggttagtggtcagtggtggtggacagacttcagcctcaacatccatcctgtggaggaggatgata
tlgcaatgtatttctgtcagcaaaagtaggaaggttcttgacgttcgggtggaggcaccagctggaaatcaaacgggggtgg
30 cggtggctcggggcggaggtgggtcgggtggcgggatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaaccggctatggtgtaaactgggttcgccagcct
ccaggaaaggtctggagtggttggaatgatatgggtgatggaagcacagactataattcagctctcaaatccagactg
agcatcaccaaggacaactccaagagccaagttttctaaaaatgaacagctctgcaaacgatgacacagccagatactact
gtgccagagatggttatagtaactttcattactatgttatggactactgggggtcaaggaacctcagtcaccgtctctcagatct
35 ggagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctgggtggaccgtcagttctctc
tccccccaaaaccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtgggtgacgtgagccacga
agacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactgggtgaatggcaaggagtacaagtga
aggtctccaacaaagccctccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
40 gtacacctgccccatcccggtgatgacgaagaaccaggtcagcctgacctgctggtcaaaaggttctatccca
gcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
acggctccttctctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaacgtcttctatgctccgtga
tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttgaacatggaga
acaacaaaacctcagtgattcaaaatccattaataattttgaagtaaagaccatacatgggagcaagtcagtgactctgg
45 gatctatctggacagtagttacaaaatggattatcctgaaatgggcatatgcataataataataagaacttcataagagc
actggaatgtcatctcgtctgtgtacggatgtggacgcagccaacctcagagagacattcatgggcctgaaataccaagtc

actagcttcttcagaggcgactactgccggagtctgactggaaagccgaaactcttcattcaggcctgccggggtacg
gagctggactgtggcattgagacagacagtgggactgatgaggagatggcttgcagaagataccgggtggaggctgactt
cctgtatgcttactctacagcacctggttactattcctggagaaattcaaaggacgggtcgtggttcatccagtcctttgcag
catgctgaagctgtacgcgcacaagctagaatttatgcacattctcactcgcgttaacaggaagggtggcaacggaattcgag
5 tcccttccctggactccactttccacgcaaagaacagatcccgtgtattgtgccatgctcacgaaagaactgtacttttate
actagctcgagatcgatga

Amino acid sequence: (SEQ ID NO: 456)

mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqkpgqppkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiemyfcqsrkvpwtfgggkgleikrgggsgggsgggsgggsgvqlkes
10 gpglvapsqslsitctvsgfsltygvnwvrpppgkglewlgmiwgdgstdynsalksrslitkdnksqvflkmnsl
qtdtdaryycardgysnfhyvmdywgqgtsvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmisrt
pevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
iektiskakgqprepvytlppsrdeitknqvslclvkgyfypsdiavewesngqpennyktpvldsdgsfflysklt
vdksrwqqgnvfscsvmhealhnhytqkslslspgkadpsnsnmennktsvdsksinnfevktihgsksvdsgiy
15 dssykmdypemgicliinnknfhkstgmssrgtdvdaanlretfmglyqvmknldltredilemdsvskedhsk
rssfvcvilshgdegviyngpvelkkltstfrgdycrsltgkpklfiiqacrgteldcgietdsgtdeemacqipvead
flyaystapgyyswrnskdgsfwfqlscsmklyahklefihiltrvnrkvatefesfslstdthakkqipcivsmllkel
yfyh

20 119. mcasp8-TM/CT

Nucleotide sequence: (SEQ ID NO: 457)

ttcgaacatggattccagagtgtctttatgctattgctgaagaactgggcagtgaagacctggctgccctcaagttcctgtg
cttgactacatcccacacaagaagcaggagaccatcgaggatgccagaagctatttctgaggctgcgggaaaagggg
atgttgagggaaggcaatctgtcttctgaaagagctgctttccacatcagtcggtgggacctgctggtcaactcctagac
25 tgcaaccgagaggagatggtgagagagctgcgggatccagacaatgccagatttctccctacagggtcatgctcttaag
ctctcagaagaagtgcgaggtggaattgagatcttttaagttcctttgaacaatgagatcccaaatgtaagctggaagat
gacttgagcctgcttgaatttttagaaatggagaagaggaccatgctggcagaaaataacttgaaaccctaaaatcaat
ctgtgaccaggtcaacaagagcctgctggggaagatcgaggattatgaaagatcaagcacagagagaagaatgagcctt
gaaggaagggaagagtgccaccttcagtttggatgagatgagcctcaaaatggcggaactgtgtgactgccaagaga
30 acaagacagtgcagtcacggacttcagacaaagtattaccaaatgaagaacaaacctcggggatactgtctgatcatcaaaa
tcatgatttcagcaaggcccggaagacataaccaactccgaaaaatgaaggacagaaaaggaacagactgtgataaaa
gaggctctgagtaagaccttaaggagcttcattttgagatagatcttacgacgactgcactgcaaatgaaatccacgagatt
ctagaaggctaccaaaagcgcagaccacaagaacaaagactgcttcatctgctgtatcctatcccacggtgacaagggtgc
gtctatggaacggatgggaaggaggcctccatctatgacctgacatcttacttactggttcaaagtgcccttccctgtctgg
35 gaaaccaagatcttttattcaggccttgccaaggaagtaacttcagaaaggagtgctgatgaggcaggcttcagcaa
cagaaccacactttagaagtggattcatctcacagaactatattccggatgaggcagactttctgctgggaatggctac
ggtgaagaactgcgtttcctaccgagatcctgtgaatggaacctggatattcagtcactttgccagagcctgagggaaga
gtcctcaaggagatgacattcttagcatcctgactggcgtgaactatgacgtgagcaataaagacgacaggaggaacaag
ggaaagcagatgccacagcccaccttcacactacggaagaagctcttcttccctccctaagactcgagatcgatt

40 Amino acid sequence: (SEQ ID NO: 458)

snmdfqsclyiaaelgsedlaalkflcldyiphkkqetiedaqlflrlrekgmleegnlsfkellfhisrwdllvnfldc
nreemvreirpdpnaqispyrvmlfklseevselelrsfkflnneipckkledlslleifvemekrtmlaennletlksi
cdqvnksllgkiedyerssterrmslegreelpsvldemslkmaelcdspreqdsesrtsdkvyqmknkprgyclii
nnhdfskareditqlrkmkdrkgtdcdkealsktfkelhfeivsyddctaneiheilegyqsadhknkdcficcilshgd
45 kgvvvygtdgkeasiydltsyftgskcpslsgkpkiffiqacqgsnfqkgvpdeagfeqqnhtlevdssshknyipdead
fllgmatvkncvsyrdpvnngtwyiqslcqlsrercpqgddilsiltgvnydvsnkddrnkgkqmpqptflrklkffp
p

120. 2e12 scFv hIgG1 (SSS-P)H WCH2 WCH3-mcasp8-TM/CT

Nucleotide sequence: (SEQ ID NO: 459)

aagcttatggattttcaagtgcagattttcagcttcctgctaatacgtgcttcagtcataatgtccagaggagtcgacattgtgct
5 caccacaatctccagcttctttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaagtggtgaatatta
tgtcacaagttaaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
tctgggggtccctgccaggttttagtggcagtggtgtctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtattttgtcagcaaaagtaggaaggttccttgacgttcggtggaggcaccagctggaaatcaaacgggggtgg
cggtggctcgggcggaggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
10 cgccctcacagagcctgtccatcacatgcaccgtctcaggggttcattaaccggctatggtgtaaactgggttcgccagcct
ccaggaaagggctctggagtggttggaatgatatgggtgatggaagcacagactataattcagctctcaaatccagactg
agcatcaccaaggacaactccaagagccaagttttctaaaaatgaacagctctgaaaactgatgacacagccagatactact
gtgccagagatggttatagtaactttcattactatgttatggactactggggtaaggaaacctcagtcaccgtctcctcagatct
ggagcccaaactctctgacaaaactcacatccccaccgtcccagcacctgaactcctgggtggaccgtcagctctcctc
15 tccccccaaaacccaaggacacctcatgatctccggaccttgaggtcacatcgtggtggtggacgtgagccacga
agacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgaggaggagca
gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactgggtgaatggcaaggagtacaagtgc
aggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
tgtacacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctgggtcaaaggcttctatccca
20 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtctctcatgctccgtga
tgcagtaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacatggattt
ccagagttgtctttatgctattgctgaagaactgggcagtgagacacctgggtgccctcaagttcctgtgcttgactacatccc
acacaagaagcaggagaccatcgaggatgccagaagctatttctgaggctgcgggaaaagggtggttgagggaagg
25 caatctgtctttcctgaaagagctgctttccacatcagtcggtgggacctgctggtcaacttctagactgcaaccgagagg
agatggtgagagagctcgggatccagacaatgccagatttctccctacagggctcgtctcttaagctctcagaagaagt
gagcgagttggaattgagatcttttaagtcttttgaacaatgagatcccaaatgtaagctggaagatgacttgagcctgctt
gaaatttttagaaatggagaagaggaccatgctggcagaaaataacttgaaaccctaaaatcaatctgtgaccaggtca
acaagagcctgctggggaagatcgaggattatgaaagatcaagcacagagagaagaatgagccttgaaaggaagggaag
30 agttgccaccttcagtttggatgagatgagcctcaaatggcggaactgtgtgactcgccaagagaacaagacagtgagt
cacggacttcagacaaagttaccaaatgaagaacaaacctcggggatactgtctgatcatcaacaatcatgatttcagcaa
ggcccggaagacataacccaactccgaaaaatgaaggacagaaaagggaacagactgtgataaagaggctctgagtaa
gacctttaaggagcttcattttgagatagtatcttacgacgactgcactgcaaatgaaatccacgagattctagaaggctacca
aagcgcagaccacaagaacaaagactgcttcactgctgtatcctatccacgggtgacaagggtgctgctatggaacgga
35 tgggaaggaggcctccatctatgacctgacatcttacttcactggttcaaagtgcccttccctgcttgggaaacccaagatctt
ttcattcaggcttgccaaggaagtaacttcagaaaggagtgctgatgaggcaggcttcgagcaacagaaccacacttta
gaagtggattcatcatctcacaagaactatattccggatgaggcagactttctgctgggaatggctacgggtgaagaactgcg
tttctaccgagatcctgtgaatggaacctggtatattcagtcactttgccagagcctgagggaaagatgctctcaaggagat
gacattcttagcatcctgactggcgtgaactatgacgtgagcaataaagacgacaggaggaacaagggaagcagatgc
40 cacagcccaccttcacactacggaagaagctcttctccctccctaatactgactcgagatcgatt

Amino acid sequence: (SEQ ID NO: 460)

mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqqkpgppkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiamyfcqsrkvpwtfgggtkleikrgggsgggsgggsgvqlkes
gpglvapsqslsitctvsgfsltygvnwvrpppgkglewlgmwgdgstdynsalksrslsitkdnsksqvflkmnsl
45 qtddtaryycardgysnfhyvmdywgqtsvtvssdlepkssdkthtspsspapellggpsvflfppkpkdtlmlsr
tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpa
piektiskakgqprepvytlppsrdeltnqvslclvkgfypsdiavewesngqpennykttpvldsdgsfflyskl
tvdksrwqqgnvfscsvmhealnhhtqkslslspgkadsnmdfqsclyaieelgsedlaalkflclidyiphkkqe

tiedaqlflrlrekgmleegnlsflkellfhisrwdllvnfldcnreemvrelrdpdnaqispyrvmlfkiseevselelrs
 fkflInneipckkleddlsleifvemekrtmlaennletlksicdqvnksllgkiedyersstermslegreelpsvlde
 mslkmaelcdspreqdsesrtdskvyqmknkprgycliinnhdfskareditqlrkmdrkgtdcdkealsktfkelh
 feivsyddctaneiheilegyqsadhknkdcficcilshgdkgvvygtdgkeasydltsyftgskcpslsqkpkiffiqa
 5 cqgsnfqkgvpdeagfeqqnhtlevdssshknyipdeadflgmatvkncvsyrdpvngrwyiqslcqlsrercpqg
 ddilsiltgvnydvsnkddrmkgkmpqptflrklfipp

121. 2e12 scFv hIgG1 (SSS-P)H P238SCH2 WCH3-mcasp8-TM/CT

Nucleotide sequence: (SEQ ID NO: 461)

10 aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
 caccacatctccagcttctttggctgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgttgatattatg
 tcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaatc
 tgggggtccctgccaggtttagtggtcagtggttctgggacagacttcagcctcaacatccatcctgtggaggaggatgatatt
 gcaatgtatttctgtcagcaaagtaggaaggtccttggtgacgttcggtggaggaccaagctggaaatcaaacggggtggc
 15 ggtggctcgggcggaggtgggtcgggtggcggcgatctcaggtgcagctgaaggagtcaggacctggcctgggtggc
 gccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcctc
 caggaaagggtctggagtggctgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactga
 gcatccaagagcaactccaagagccaagttttcttaaaaaatgaacagcttgcaactgatgacacagccagatactactg
 tgccagagatggttatagtaactttcattactatgttatggactactggggtcaaggaacctcagtcaccgtctcctcagatctg
 20 gagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctgggtggatcgtcagcttctcctt
 cccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacgtgagccacgaa
 gacctgaggtcaagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcag
 tacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcac
 ggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggt
 25 gtacacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctgtaaaggcttctatccca
 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 acggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
 tgcagtaggctctgcacaaccactacacgcagaagagcctctcctctgtctccgggtaaagcggatccttcgaacatggattt
 ccagagtgtctttatgctattgctgaagaactgggcagtgagacacctggctgccctcaagtctctgtgcttgactacatccc
 30 acacaagaagcaggagaccatcgaggatgccagaagctatttctgaggctgcgggaaaaggggatgttgagggaagg
 caatctgtcttctgaaagagctgctttccacatcagtcggtgggacctgctggtcaacttctagactgcaaccgagagg
 agatgggtgagagagctcgggatccagacaatgccagatttctcctacagggctatgctctttaagctctcagaagaagt
 gagcgagttggaattgagatcttttaagttctttgaacaatgagatcccaaatgtaagctggaagatgacttgagcctgctt
 gaaatttttagaaatggagaagaggaccatgctggcagaaaaataacttgaaaccctaaaatcaatctgtgaccaggtca
 35 acaagagcctgctggggaagatcaggattatgaaagatcaagcacagagagaagaatgagcctgaagggaagggaag
 agttgccaccttcagttttggatgagatgagcctcaaatggcggaactgtgtgactcgccaagagaacaagacagtgagt
 cacggacttcagacaaagttaccaaatgaagaacaaacctcggggatactgtctgatcatcaacaatcatgatttcagcaa
 ggcccgggaagacataacccaactccgaaaaatgaaggacagaaaagggaacagactgtgataaagggtctgtagtaa
 gaccttaaggagcttcattttgagatagtatcttacgacgactgcactgcaaatgaaatccacgagattctagaaggctacca
 40 aagcgcagaccacaagaacaaagactgcttcacatgctgtatcctatcccacgggtgacaagggtgctgctatggaacgga
 tgggaaggaggcctccatctatgacctgacatcttacttactggttcaagtgccttccctgtctgggaacccaagatctt
 ttctattcaggcttgccaagggaagtaactccagaaggaagtgcctgatgaggcaggcttcgagcaacagaaccacattta
 gaagtggattcatcatctcacaagaactatattccggatgaggcagactttctgctgggaatggctacggtgaagaactgcg
 ttctctaccgagatcctgtgaatggaacctgggtatattcagtcactttgccagagcctgagggaaagatgctcctaaggagat
 45 gacattcttagcatcctgactggcgtgaactatgacgtgagcaataaagacgacagaggagaacaagggaagcagatgc
 cacagcccaccttcacactacggaagaagctcttcttccctccc taatgactcgagatcgattc

Amino acid sequence: (SEQ ID NO: 462)

mdfqvqifsflisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyqqkpgppkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiamyfcqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
gpglvapsqslsitctvsgfsltgygvnwvrqppgkglewlgmiwgdgstdynsalksrsltkdnksqvflkmnsl
qtdtdaryycardgysnfhyvmdywggqtsvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmsrt
5 pevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
iektiskakgqprepvytlppsrdeltknqvsitclvkgfypsdiavewesngqpennyktpvldsdgsfflysklt
vdksrwqqgnvfscsvmhealhnhytqkslslspgkadpsnmdfqsclyiaaelgsedlaalkfclldyiphkkqet
iedaqlflrlrekgmleegnslfkellfhisrwdllvnfldcnreemvrelrdpdnaqispyrvmflklseevselelrsf
kflInneipkckleddlsleifvemekrtmlaennletlksicdvnskllgkiedyersstermslegreelppsvlde
10 mslkmaelcdspreqdsesrtsdkvyqmknkprgycliinnhdfskareditqlrkmdrkgtdcdkealsktfkelh
feivsyddctaneiheilegyqsadhknkdcficcilshgdkgvvygtdgkeasydltsyftgskcpslsgkpkiffiqa
cqgsnfqkgvpdeagfeqqnhtlevdssshknyipdeadflgmatvkncvsyrdpvngtwyiqslcqlsrercpqg
ddilsiltgvnydvsnkddrnkgkqmpqptflrkklffpp

15 122. hcasp3-TM/CT

Nucleotide sequence: (SEQ ID NO: 463)

Ggatccttcgaacatggagaacactgaaaactcagtgattcaaaatccattaaaaatttgaaccaaagatcatacatgga
agcgaatcaatggactctggaatatccctggacaacagtataaaatggattatcctgagatgggtttatgtataataattaata
ataagaatttcataaaagcactggaatgacatctcggctctggtacagatgctgatgcagcaaacctcagggaacattcag
20 aaacttgaaatatgaagtcaggaataaaaatgatcttacacgtgaagaaattgtggaattgatgcgtgatgtttctaaagaaga
tcacagcaaaaggagcagtttgtttgtgtgcttctgagccatggtgaagaaggaataattttggaacaaatggacctgtga
cctgaaaaaataacaaacttttcagaggggagcgtttagaagtctaactggaaaacccaaacttttcattattcaggcctg
ccgtggtacagaactggactgtggcattgagacagacagtggtgttgatgatgacatggcgtgtcataaaataccagtggga
ggccgacttcttgtatgcatactccacagcacctggttattattcttggcgaaattcaaaggatggctcctggttcatccagtcg
25 ctttgtgccatgctgaaacagatgccgacaagcttgaatttatgcacattcttaccggggttaaccgaaagggtggcaacaga
atgtgagtccttttctttgacgctacttttcagcaagaacagattccatgtattgtttccatgctcacaaaagaactctatttt
atcactaactcgatcgata

Amino acid sequence: (SEQ ID NO: 464)

dpsnmentensvdsksiknlepkiihgsesmdsgisldnsykmdypemglciiinnknfhkstgmtsrsgtdvdaa
30 nlretfmlkyevrnknldtreeivelmrdvskedhskrssfvcvllshgeegiifgtngpvdlkkitnffrgdrersltgkp
klfiiaqrgtelcgieitdsgvdddmachkipveadflyaystapgyyswrnskdgswfiqslcamlkqyadklef
mhiltrvnrkvatefesfsfd atfhakkqipcivsmllkelyfyh

123. 2e12 scFv hIgG1 (SSS-P)H WCH2 WCH3-hcasp3-TM/CT

35 Nucleotide sequence: (SEQ ID NO: 465)

aagcttatggatttcaagtcagattttcagcttcctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgctc
acccaatctccagcttcttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaaagtgtgaatattat
gtcacaagtttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatcaacgtagaat
ctgggggtccctgccaggttagtggcagtggtctgggacagacttcagcctcaacatccatcctgtggaggaggatgatat
40 tgcaatgtatttctgcagcaaaagtaggaaggttccttggaacttcggtggaggacccaagctggaaatcaaacgggtgg
cggtggtcgggagggtgggtcgggtggcgcgatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctattaaccggctatggtgtaaactgggttcgccagcct
ccaggaaagggtctggagtggttggaatgatatgggtgatggaagcacagactataattcagcttcaaatccagactg
agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgaaactgatgacacagccagataactact
45 gtgccagagatggttatagtaactttcattactatgttatggactactgggtcaaggaacctcagtcaccgtctcctcagatct
ggagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctgggtggaccgtcagcttctcctc

5 ttccccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacga
 agaccctgaggtcaagtccaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
 gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga
 aggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
 10 gtacacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgctggtcaaaggcttctatccca
 gcgacatcgccgtggagtgaggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtctctcatgctccgtga
 tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttgaacatggaga
 aactgaaaactcagtggttcaaaatccattaaaaatttgaaccaaagatcataatggaagcgaatcaatggactctgg
 15 aatatccctggacaacagttataaaatggattatcctgagatgggtttatgtataataattaataagaattttcataaaagca
 ctggaatgacatctcggtctggtacagatgtgatgcagcaaacctcagggaaacattcagaaacttgaaatatgaagtcag
 gaataaaatgatcttacacgtgaagaaattgtggaattgatgcgtgatgtttctaaagaagatcacagcaaaaggagcagtt
 ttgtttgtgtctctgagccatggtgaagaaggaataattttggaacaaatggacctgttgacctgaaaaaataacaaactt
 ttcagaggggatcgttgtagaagtctaactggaaaacccaaacttttcattatcaggcctgccgtggtacagaactggact
 20 gtggcattgagacagacagtggtgttgatgatgacatggcgtgtcataaaataaccagtgagggccgacttctgtatgcatac
 tccacagcacctggttattattcttggcgaaattcaaaggatggctcctggtcatccagtcgctttgtccatgctgaaacagt
 atgccgacaagcttgaatttatgcacattcttaccggggttaaccgaaaggtggcaacagaatttgagtcctttcctttgacgc
 tacttttcatgcaaagaacagattccatgtattgtttccatgctcacaaaagaactctattttatcactaactcgagatcgata

Amino acid sequence: (SEQ ID NO: 466)

20 mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyqqkpgppkllisaasn
 vesgvparfsgsgsgtdfslnihpveeddiemyfcqsrkvpwtfggggleikrgggsgggsgggsgvqlkes
 gpglvapsqslsitctvsgfsltgygvnwvrpppgkglewlgmiwgdgstdynsalksrslsitkdnsksqvflkmnsl
 25 qtddtaryycardgysnfhyvmdywgqgtsvtvssdlepkssdkthtspspapellggpsvflfppkpkdtlmsir
 tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpa
 30 piektiskakgqprepvytlppsrdeitknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsfflyskl
 tvdkswqqgnvfscsvmhéalhnhytqkslslspgkadsnmentensvdsksiknlepkiihgsesmdsgisldn
 sykmdypemglciiiinnknfhkstgmtsrsgtdvdaanlretfmklyevrnkndltreeivelmrdvskedhskrssf
 vcvlshgeegiiifngpvdllkitnffrgdrcltgpklfiiaqrgteldcgetdsgvdddmachkipveadflya
 ystapgyyswrmskdgswfqslcamlkqyadklefmhiltrvnrkvatefesfsfdatfhakkqipicivsmlltkelyf
 yh

123. 2e12 scFv hIgG1 (SSS-P)H P238SCH2 WCH3-hcasp3-TM/CT

Nucleotide sequence: (SEQ ID NO: 467)

35 aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
 caccatctccagcttcttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaagtggtgaatatta
 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcatccaacgtagaa
 tctggggtccctgccaggttagtggtcagtggtgctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
 ttgcaatgtatttctgtcagcaaagtaggaaggttccttgacgttcgggtggaggcaccaagctggaaatcaaacggggtg
 40 cggtggctcgggcggaggtgggtcgggtggcggcgatcaggtgcagctgaaggagtcaggacctggcctgggtgg
 cgccctcacagagcctgtccatcacatgcaccgtctcagggttctattaaccggctatggtgtaaactgggttcgccagcct
 ccaggaaagggtctggagtggtgggaatgatatggggtgatggaagcacagactataattcagctctcaatccagactg
 agcatcaccaaggacaactccaagagccaagttttctaaaaatgaacagctctgaaactgatgacacagccagatactact
 gtgccagagatggttatagtaacttcattactatgttatggactactggggtcaaggaacctcagtcaccgtctcctcagatct
 ggagcccaaatcttctgacaaaactcacatccccaccgtcccagcacctgaactcctgggtggatcgtcagttctctc
 45 ttccccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgagccacga
 agaccctgaggtcaagtccaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
 gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga

aggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
 tgtacacctgccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtaaaggcttctatccca
 ggcacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
 5 tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacatggaga
 aactgaaaactcagtgattcaaaatccattaaaaatttgaaccaaagatcatacatggaagcgaatcaatggactctgg
 aatatccctggacaacagtataaaatggattatcctgagatgggtttatgtataataattaataagaatttccataaaagca
 ctggaatgacatctcggctgtgtacagatgtcgatgcagcaaacctcagggaacattcagaacttgaaatatgaagtacg
 gaataaaaatgatcttacacgtgaagaaattgtggaattgatgcgtgatgttctaaagaagatcacagcaaaaggagcagtt
 10 ttgtttgtgtcttctgagccatggtgaagaaggaataattttgaacaaatggacctgttgacctgaaaaaataacaaactt
 ttccagaggggagctgtgtagaagtctaactggaaaacccaaacttttcattatcaggcctgccgtggtacagaactggact
 gtggcattgagacagacagtggtgttgatgatgacatggcgtgtcataaaataaccagtggaggccgactcttctgtatgcatac
 tccacagcacctggttatttcttggcgaaattcaaaggatggctcctgggtcatccagtcgctttgtccatgctgaaacagt
 atgccgacaagcttgaatttatgcacattcttaccgggttaaccgaaaggtggcaacagaatttgagtcctttcctttgacgc
 15 tacttttcatgcaaagaacagattccatgtattgttccatgctcacaaaagaactctattttatcactaactcgagatcgataa

Amino acid sequence: (SEQ ID NO: 468)

mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyvvtslmqwyqqkpgqppkllisaasn
 vesgvparfsgsgsgtdfslnihpveediamyfcqsrkvpwtfgggkgleikrgggsgggsgggsgggsgvqlkes
 gpglvapsqslsitctvsgfsltygvnwvrqppgkglewlgmiwgdgstdynsalksrlsitkdnsksqvfikmnsi
 20 qtddtaryycardgysnfhyvmdywgqtsvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmiisrt
 pevtevvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
 iektiskakgqprepvytlppsrdeltknqvsltclvkgfypsdiavewesngqpennyktppvldsdgsfflysklt
 vdkswqqgnvfscsvmhcalhnhytqkslslspgkdpnsmentensvdsksiknlepkiihgsesmdsgisldn
 sykmdypemglciiinnknfhkstgmsrgtdvdaanretfmlkyevmknldtreeivelmrdvskedhskrssf
 25 vcvlshgeegiiifgtngpvdllkitnffrgdrersltgkpklfiiqacrgteldcgietdsgvdddmachkipveadflya
 ystapgyyswrnskdgswfqslcamlkqyadklefmhiltrvrnkvatfefsfdatfhakkqipicivsmlltkelyf
 yh

124. hcas8-TM/CT

30 hCaspase8B:

Nucleotide sequence: (SEQ ID NO: 469)

ggatccttcgaacatggacttcagcagaaatctttatgatattggggaacaactggacagtgaagatctggcctccctcaagt
 tcttgagcctggactacattccgcaaaggaaagcaagaacctcaaggatgccttgatgtattccagagactccaggaaa
 agagaatgttgaggagaaagcaatctgtccttctgaaggagctgctctccgaattaatagactggatttctgattacctacc
 35 taacactagaaaggaggagatggaagggaacttcagacaccaggcagggctcaaatctgcctacagggatcatgctc
 tatcagatttcagaagaagtgagcagatcagaattgaggtcttttaagtttcttttgcaaggaggaaatctccaaatgcaaaactg
 gatgatgacatgaacctgctggatattttcatagagatggagaagagggtcatcctgggagaaggaaagtggacatcctg
 aaaagagtctgtgccccaatcaacaagagcctgctgaagataatcaacgactatgaagaattcagcaaaagagagaagcag
 cagccttgaaaggaaagtctgatgaattttcaaatggggaggagttgtgtgggtaatgacaatctcgactctccaagagaa
 40 caggatagtgaatcacagactttggacaaggtttaccaaatgaaaagcaaacctcggggatactgtctgatcatcaacaatc
 acaattttgcaaaagcacgggagaaagtgccaaaacttcacagcattaggagacaggaatggaacacacttgatgcaggg
 gctttgaccacgaccttgaagagcttattttgagatcaagccccacgatgactgcacagtagagcaaatctatgagattttg
 aaaatctaccaactcatggaccacagtaacatggactgctcatctgctgtatcctctccatggagacaagggcacatctat
 ggactgatggacaggaggcccccatctatgagctgacatctcagttcactggttgaaagtgccttcccttctgtgaaaac
 45 ccaaagtgtttttattcaggcttgcagggggataactaccagaaaggtatactgttgagactgattcagaggagcaaccc
 tatttagaatggatttatcaccctcaaacgagatatcccgatgaggtgactttctgctggggatggccactgtgaat
 aactgtgttctaccgaaacctgcagagggaacctggatcatccagtcactttgccagagcctgagagagcgatgtcctc

gaggcgatgatattctcaccatcctgactgaagtgaactatgaagtaagcaacaaggatgacaagaaaaacatggggaaa
cagatgcctcagcctactttcacactaagaaaaaaactgtcttcccttctgattgagcatgcatcgata

Amino acid sequence: (SEQ ID NO: 470)

5 DpsnmDFSrnlydigeqlDsedlaslkfIsldyipqrkQepikdalmlfQrlQekrmleesnlSflkellfrinrldllityln
trkeemerelqtpgraqisayrvmlYqiseevsrslrsfkflQeeiskcklDDdmnlldifiemekrvilgegkldilkr
vcaqinkslIkiindyefskerssslegspdefsngeelcgvmtisdspreqdsesqtlDkvYqmkskprgycliinnh
nfakarekvplhsirdnrthldagaltttfeelhfeikphddctveqiyeilkiYqlmdhsnmDcficciIshgdkgiiyg
tdgqeapiyeltsqftglkcpSlagkpkvffiqacqgdnyqkgipvetdseeqylemdlsspqtryipdeadflIgmata
vnncvsyrnpaegtwyiqslcqlsrercprgddiltiltevnYevsnkddkknmgkmpqptflrkkIvfpSDh

10

hCaspase8C:

Nucleotide sequence: (SEQ ID NO: 471)

15 ggatccttcgaacatggacttcagcagaaatctttatgatattggggaacaactggacagtgaagatctggcctccctcaagt
tcttgagcctggactacattccgcaaaggaagcaagaacccatcaaggatgccttgatgttattccagagactccaggaaa
agagaatgttgaggaaagcaatctgtccttctgaaggagctgctctccgaattaatagactggatttgctgattacctacc
20 taaacactagaaaggaggagatggaaagggaacttcagacaccaggcagggctcaaatttctgcctacagggctcatgctc
tatcagatttcagaagaagtgagcagatcagaattgagcttttaagtttctttgcaagaggaaatctcaaactgaaactgga
tgatgacatgaacctgctggatatttcatagagatggagaagagggctcatcctgggagaaggaaagtggacatcctgaa
aagagtctgtgccccaaatcaacaagagcctgctgaagataatcaacgactatgaagaattcagcaaaggggaggagtgtg
25 gtggggtaatgacaatctcggactctccaagagaacaggatagtgaatcacagactttggacaaagtttaccaaatgaaaa
gcaaacctcggggatactgtctgatcatcaacaatcacaatttgcaaaagcacgggagaaagtgccccaaacttcacagca
ttagggacaggaatggaacacacttgatgcaggggctttgaccacgacctttgaagagcttcatttgatcaagcccca
cgatgactgcacagtagagcaaatctatgagattttgaaaatctaccaactcatggaccacagtaacatggactgcttcatct
gctgtatcctctcccatggagacaagggcatcatctatggcactgatggacaggaggcccccatctatgagctgacatctca
30 gttcactggtttgaagtgcccttcccttgctggaaaacccaaagtgtttttattcaggcttgctcagggggataactaccagaaa
ggtatacctgttgagactgattcagaggagcaaccctatttagaaatggatttatcatcacctcaaacgagatatatcccggat
gaggctgactttctgctggggatggccactgtgaataactgtgttctaccgaaacctgcagagggaacctggatcatcc
agtcaatttgccagagcctgagagagcgatgtcctcagggcgatgatatttcaccatcctgactgaagtgaactatgaagt
aagcaacaaggatgacaagaaaaacatggggaaacagatgcctcagcctactttcacactaagaaaaaaactgtcttccc
35 ttctgattgagcatgca tcgata

Amino acid sequence: (SEQ ID NO: 472)

35 dpsnmDFSrnlydigeqlDsedlaslkfIsldyipqrkQepikdalmlfQrlQekrmleesnlSflkellfrinrldllityln
trkeemerelqtpgraqisayrvmlYqiseevsrslrsfkflQeeiskcklDDdmnlldifiemekrvilgegkldilkr
vcaqinkslIkiindyefskgeelcgvmtisdspreqdsesqtlDkvYqmkskprgycliinnhnfakarekvplhs
irdrnghldagaltttfeelhfeikphddctveqiyeilkiYqlmdhsnmDcficciIshgdkgiiygtgdgqeapiyelts
qftglkcpSlagkpkvffiqacqgdnyqkgipvetdseeqylemdlsspqtryipdeadflIgmatavnncvsyrnpa
egtwyiqslcqlsrercprgddiltiltevnYevsnkddkknmgkmpqptflrkkIvfpSD

125. 2e12 scFv hIgG1 (SSS-P)H WCH2 WCH3-hcasp8-TM/CT

40 Nucleotide sequence: (SEQ ID NO: 473)

aagcttatggattttcaagtgcagatttccagcttctgctaatacagtgccttcagtcataatgtccagaggagtcgacattgtgct
caccacatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaagtggtgaatatta
tgacacaagtttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcacccaacgtagaa
45 tctgggggtccctgccaggttagtggtcagtggtctgggacagacttcagcctcaacatccatctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaagtaggaaggttccttgacgttcggtggaggcaccaagctggaaatcaaacgggggtgg

5 cggaggctcgggagggtgggtcgggtggcgccgatctcaggtgcagctgaaggagtcaggacctggcctgggtg
 cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccgctatggtgtaaacgggttcgccagcct
 ccaggaaagggtctggagtgggtgggaatgatatggggtgatggaagcacagactataattcagctctcaaatccagactg
 agcatcaccaaggacaactccaagagccaagttttctaaaaatgaacagctctgcaaactgatgacacagccagatactact
 10 gtgccagagatggttatagtaactttcattactatgttatggactactggggtcaaggaacctcagtcacctctcctcagatct
 ggagcccaaatcttctgacaaaactcacatccccaccgtccccagcacctgaactcctgggtggaccgtcagttctctc
 ttcccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtgggtgggacgtgagccacga
 agacctgaggtcaagtccaactgggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
 gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga
 15 aggtctccaacaaagccctccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
 tgtacacctgccccatcccggtgatgactgaccaagaaccaggtcagcctgacctgctgggtcaaaagcttctatccca
 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccg
 acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
 tgcattgaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaagcggatccttcgaacatggact
 20 tcagcagaaatctttatgatattgggaacaactggacagtgaagatctggcctccctcaagttcctgagcctggactacatt
 ccgcaaaggaagcaagaacctcaaggtgcttgatgtattccagagactccaggaaaagagaatgttgaggagaaag
 caatctgctctcctgaaggagctgcttctccgaattaatagactggatttgcgtgattacctacctaataactagaaaggagga
 gatggaaagggaaacttcagacaccaggcagggtcaaatctgcctacaggggtcatgctctatcagatttcagaagaagt
 gagcagatcagaattgaggtcttttaagtcttttgcaagaggaaatctccaaatgcaaactggatgatgacatgaacctgct
 25 ggatattttcatagagatggagaagagggtcatcctgggagaaggaaagttggacatcctgaaaagagtctgtgccccaat
 caacaagagcctgctgaagataatcaacgactatgaagaattcagcaaaagagagaagcagcagcctgaaggaagtct
 gatgaattttcaaatggggaggagtgtgtggggtaatgacaatctcgactctcaagagaacaggatagtgaatcacag
 actttggacaaaagtttaccaaaatgaaaagcaaacctcggggatactgtctgatcatcaacaatcacaattttgcaaaagcacg
 ggagaaagtgccaaaacttcacagcattaggagacaggaatggaacacacttggatgcaggggctttgaccacgaccttg
 30 aagagcttcattttgagatcaagccccacgatgactgcacagtagagcaaatctatgagattttgaaaatctaccaactcatg
 gaccacagtaacatggactgcttcatctgctgtatcctctccatggagacaagggcatcatctatggcactgatggacagg
 agggccccatctatgagctgacatctcagttcactggttgaaagtgccttcccttgcgtgaaaacccaaagtgtttttattca
 ggcttgcagggggataactaccagaaaggtatacctgttgagactgattcagaggagcaacctatttagaaatggatttat
 catcacctcaaacgagatatatcccggtgagggtgactttctgctggggatggccactgtgaataactgtgttctaccga
 35 aacctgcagagggaaacctggtacatccagtcacttggcagagcctgagagagcgatgtcctcaggcgatgatattctc
 accatcctgactgaagtgaactatgaagtaagcaacaaggtgacaagaaaacatggggaacagatgcctcagcctac
 ttccactaagaaaaaactgtcttccctctgattgagcatgcatcgata

Amino acid sequence: (SEQ ID NO: 474)

35 mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqqkpgqppkllisaasn
 vesgvparfsgsgsgtdfslnihpveeddiameyfcqsrkvpwtfgggtkleikrgggsgggsgggsgvqlkes
 gpglvapsqslsitctvsgfsltygvnwvrppgkglewlgmiwgdgstdynsalksrslsitkdnsksqvflkmnsl
 qtddtaryycardgysnfhyvmdywgqgtsvtvssdlepkssdkthtsppspapellggpsvflfppkpkdtlmisr
 40 tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpa
 piektiskakgqprepqvylppsrdeitknqvsitclvkgfypsdiavewesngqpennykttppvldsdgsfflyskl
 tvdksrwqqgnvfscsvmhcalhnhytqkslslspgkadsnmdfsmlydigeqldsedlaslkflsldyipqrkqe
 pikdalmlfqlqekrmleesnlsflkellfrinrldlitylnrkeemerelqtpgraqisayrvmllyqiseevsrseisfk
 flqeeiskcklddmnlldifiemekrvilgegkldilkrvcaqinkslkiindyeefskerssslegspdefsngeelc
 gvmtisdspreqdsesqldkvyqmkskprgycliinnhnfakarekvpklhsirdmngthldagaltttfeelhfeikp
 45 hddctveqiyelkiyqlmdhsnmdcficilshgdkgiyygtgdqeapiyeltsqftglkcpslagkpkvffiqacqgd
 nyqkgipvetdseeqpylemdlsspqtryipdeadflgmatvnnvsvrnpaegtwiyqlcqlsrercprgddiltl
 tevnyevsnkddkknmgkqmpqptflrklkvfspd

126. 2e12 scFv hIgG1 (SSS-P)H P238SCH2 WCH3-hcasp8B-TM/CT

(other caspase 8 isoforms are similar)

Nucleotide sequence: (SEQ ID NO: 475)

5 aagcttatggattttcaagtgcagattttcagcttctgctaatactgcttcagtcataatgtccagaggagtcgacattgtgct
caccacatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgttgaatatta
10 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
tctgggggtccctgccagggttagtggtcagtggtgtgggacagacttcagcctcaacatccatcctgtggaggaggatgata
ttgcaatgtatttctgtcagcaaagtaggaaggttccttggtgacgttcggtggaggcaccacagctggaaatcaaacgggggtgg
cggtgggtcggggcggagggtgggtcgggtggcggcggtatcaggtgcagctgaaggagtcaggacctggcctgggtgg
15 cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaaactgggttcgccagcct
ccaggaaaagggtctggagtgggtgggaatgatatgggtgatggaagcacagactataattcagctctcaaatccagactg
agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgaaaactgatgacacagccagatactact
gtgccagagatggttatagtaactttcattactatggtatggactactgggggtcaaggaaacctcagtcaccgtctcctcagatct
ggagcccaaatcttctgacaaaactcacatccccaccgtcccagcacctgaactcctgggtggatcgagcttctcctct
20 tcccccaaaaacccaaggacacctcatgatctccggaccctgaggtcacatgcgtgggtgggtgacgtgagccacga
agacctgaggtcaagttcaactgttacgtggacggcgtggaggtgcataatgccaagacaaagccgaggaggagca
gtacaacagcacgtaccgtgtgtcagcgtcctcaccgtcctgcaccaggactgggtgaatggcaaggagtacaagtgc
agggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
tgtacacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctggtaaaaggcttctatccca
25 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
acggctccttctcctctacagcaagtcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
tgcagtaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaagcggatccttcgaacatggact
tcagcagaaaatctttatgatattggggaacaactggacagtgaagatctggcctccctcaagtctcagcctggactacatt
ccgcaaagggaagcaagaacccatcaaggatgccttgatgttattccagagactccaggaaaagagaatgttgaggaaag
30 caatctgtccttctgaaggagctgctcttccgaattaatagactggatttctgattacctacctaataactagaagaggga
gatggaaagggaaacttcagacaccaggcagggctcaaatcttctgcttacagggtcatgctctatcagatttcagaagaagt
gagcagatcagaattgaggtcttttaagtttcttttgcaagaggaaatctccaaatgcaaactggatgatgacatgaacctgct
ggatattttcatagagatggagaagagggtcatcctgggagaaggaaagtggacatcctgaaaagagtctgtgccccaat
caacaagagcctgctgaagataatcaacgactatgaagaattcagcaaagagagaagcagcagcctgaagggaagtct
35 gatgaattttcaaatggggaggagtgtgtggggtaatgacaatctcggactctccaagagaacaggatagtgaatcacag
actttggacaaagtttacaaatgaaaagcaaacctcggggatactgtctgatcatcaacaatcacaattttgcaaaagcacg
ggagaaagtgccaaaacttcacagcattagggacaggaatggaacacacttggatgcagggggtttgaccacgacctttg
aagagcttcattttgatgaagccccacgatgactgcacagtagagcaaatctatgagattttgaaaatctaccaactcatg
40 gaccacagtaacatggactgcttcatctgctgtatcctctccatggagacaagggcatcatctatggcactgatggacagg
aggccccatctatgagctgacatctcagttcactggttgaagtgccttcccttgctggaaaacccaaagtgtttttattca
ggcttgcagggggataactaccagaaaggtatacctgttgagactgattcagaggagcaaccctatttagaaatggattat
catcacctcaaacgagatatatcccggatgaggctgacttctgtggggatggccactgtgaataactgttttctaccga
aaccttcgagaggggaacctggtacatccagtcactttgccagagcctgagagagcgtgtcctcaggcgatgatattctc
accatcctgactgaagtgaactatgaagtaagcaacaaggatgacaagaaaaacatggggaaacagatgcctcagcctac
45 tttcacactaagaaaaaactgttctcccttctgattgagcatgcatcga taa

Amino acid sequence: (SEQ ID NO: 476)

mdfqvqifsfllisavimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqkpgqppkilisaasn
vesgvparfsgsgsgtdfslnihpveeddiamyfcqsrkvpwtfgggtkleikrgggsgggsgggsgggsgvqlkes
45 gpglvapsqslsitctvsgfsltygvnwvrpppgkglewlgmiwgdgstdynsalksrlsitkdnsksqvflkmnsl
qtdtdaryycardgysnfhyvmdywgqtsvtvssdlepkssdkthtspspapellggssvflfppkpkdltlmisrt
pevtcvvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
iektiskakgqprepqvylppsrdeltnqnsvlclvkgyfypsdiavewesngqpennyktpvpldsdgsfflysklt
vdksrwqqgnvfscsvmhlehnhytqkslslspgkadsnmdfsmlydigeqlsdslaslkflsldyipqrkqep

ikdalmlfqrleqkrmleesnlslkellfrinrldlitylntrkeemerelqtpgraqisayrvmlyqiseevsrslrsfkf
llqeeiskckldddmnlldifiemekrvilgegkldilkrvcaqinkslkiindyefskerssslegspdefsngeelcg
vmtisdspreqdsesqtdkvyqmkskprgycliinnhnfakarekvpklhsirdrngthldagaltttfeelhfeikph
ddctveqiyeilkiyqlmdhsnmddcficclshgdkgiygtgqeapiyeltsqftglkcpslagkpkvffiqacqgdn
5 yqkgipvetdseeqpylemdlsspqtryipdeadflgmatvnnvsvrmpaegtwyiqslcqlsrercprgddiltilte
vnyevsnkddkknmgkmpqptflrkklvfspd

127. hFADD-TM/CT

Nucleotide sequence: (SEQ ID NO: 191)

10 Gttgtggatccttgaacccgttctggtgctgctgactcggtgctgctccagcctgtcgagcagcgagctgaccgagctc
aagttcctatgcctcgggcgcgtgggcaagcgcaagctggagcgcgtgcagagcggcctagacctcttctccatgctgct
ggagcagaacgacctggagcccgggcacaccgagctcctgcgcgagctgctgcctccctgcggcgccacgacctgct
gcggcgctcgacgacttcgagcgggggcgggcgccggggcgccgctggggaagaagacctgtgtgcagcattta
acgtcatatgtgataatgtgggaaagattggagaaggtggctcgtcagctcaaagtctcagacaccaagatcgacagca
15 tcgaggacagatacccccgcaacctgacagagcgtgtgcgggagtcactgagaatctggaagaacacagagaaggaga
acgcaacagtgggccacctggtggggctctcaggtcctgccagatgaacctgggtgctgacctggtacaagaggttcag
caggcccgtgacctccagaacaggagtggggccatgtccccgatgtcatggaactcagacgcactctacctccgaagcgtc
ctgataactcgagatcgataacaac

Amino acid sequence: (SEQ ID NO: 192)

20 Dpsnpflvllhsvssslsssteltklclgrvgkrklervqsgldlfsmlleqndlepghtellrellaslrhdllrrvddfea
gaaagaapgeedlcaafnvicdnvgkdwrrlarqlkvsdtkidsiedryprmltervreslriwkntekenatvahlvga
lrscqmnlvadlvqevqqardlqnrsgamspmswnsdastseas

128. 2e12 scFv-hIgG1 (SSS-P)H WCH2 WCH3-hFADD-TM/CT

25 Nucleotide sequence: (SEQ ID NO: 477)

aagcttatggattttcaagtgagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttcttggctgtgtctctaggtcagagagccaccatctctgcagagccagtgaaagtgtgaatatta
tgtcacaagtttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcaccaacgtagaa
tctggggtccctgccaggttagtggcagtggtgctgggacagacttcagcctcaacatccatcctgtggaggaggatgata
30 ttgcaatgtatttctgtcagaaagtaggaaggttccttgacgttcgggtggaggcaccaagctggaaatcaaacggggtgg
cgggtggctcgggagggtgggtcgggtggcgggatctcaggtgcagctgaaggagtcaggacctggcctggtgg
cgccctcacagagcctgtccatcacatgcaccgtctcagggttctcattaaccggctatggtgtaactgggttcgccagcct
ccaggaaagggtctggagtggctgggaatgatatggggtgatggaagcacagactataattcagctctcaatccagactg
agcatcaccaaggacaactccaagagccaagttttcttaaaaatgaacagctctgaaactgatgacacagccagataactact
35 gtgccagagatggttatagtaactttcattactatgttatggactactggggtcaaggaaacctcagtcaccgtctcctcagatct
ggagcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctgggtggaccgtcagttctctc
ttcccccaaaaaccaaggacacctcatgatctcccgaccctgaggtcacatgcgtggtgggtggacctgagccacga
agacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagca
gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga
40 aggtctcaacaaagccctccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacagg
tgtacacctgccccatccgggatgagctgaccaagaaccagggtcagcctgacctgctggtcaaaggcttctatccca
gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
acggctccttctcctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaactcttctcatgctcctga
tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaagcggatccttcgaacccgttct
45 ggtgctgctgactcggtgtcgtccagcctgtcgagcagcgagctgaccgagctcaagttcctatgcctcgggcgcgtgg
gcaagcgcaagctggagcgcgtgcagagcggcctagacctcttctccatgctgctggagcagaacgacctggagcccc

ggcacaccgagctcctgcgcgagctgctgcctccctgcggcgccacgacctgctgcggcgctgcgacttcgagg
 cggggcgggcgccggggccgcgcctggggaagaagacctgtgtgcagcatttaacgtcatatgtgataatgtggggaa
 agattggagaaggctggctcgtcagctcaaagtctcagacaccaagatcgacagcatcaggacagataccccgcaac
 ctgacagagcgtgtgcgggagtcactgagaatctggaagaacacagagaaggagaacgcaacagtggcccacctggtg
 5 ggggctctcaggtcctgccagatgaacctggtggctgacctggtacaagaggttcagcaggcccgtagctccagaaca
 ggagtggggccatgtccccgatgtcatggaactcagacgcattacctccgaagcgtcctgataactcgagatcgataaca
 ac

Amino acid sequence: (SEQ ID NO: 478)

mdfqvqifsfllisasvimsrgvdivltqspaslavlsgqratiscrasesveyyvtslmqwyyqkpgppkllisaasn
 10 vesgvparfsgsgsgtdfslnihpveeddiamyfcqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
 gpglvapsqslsitctvsgfsltgygvnwvrpppgkglewlgmiwgdgstdynsalksrslsitkdnsksqvflkmnsl
 qtddtaryycardgysnfhyvmdywgqtsvtvssdlepkssdkthtspsspapellgpgsvflfppkpkdtlmsir
 tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpa
 piektiskakgpprepqvytlppsrdeitknqvslclvkgyfypsdiavewesngqpennykttppvldsdgsfflyskl
 15 tvdksrwqqgnvfscsvmhcalhnhytqkslslspgkadsnpflvlhsvssslsseltelkflclgrvgkrklervqs
 gldlfsmlleqndlepghtellrellaslrhdllrrvddfeagaaagaapgeedlcaafnvicdnvgkdwrrlarqlkvsd
 tkidsiedryprnltervreslriwkntekenatvahlvgalrscqmnlvadlvqevqqardlqnrsgamspmswnsd
 astseas

20 129. 2e12 scFv-hIgG1 (SSS-P)H P238SCH2 WCH3-hFADD-TM/CT

Nucleotide sequence: (SEQ ID NO: 479)

aagcttatggattttcaagtgcagatttcagcttcctgctaatacgtgcttcagtcataatgtccagaggagtcgacattgtgct
 caccacatctccagctctttggctgtgtctctaggtcagagagccaccatctcctgcagagccagtgaagtggtgaatattat
 gtcacaagtttaatgcagtggtaccaacagaaaccaggacagccacccaaactcctcatctctgctgcattcaacgtagaat
 25 ctgggggtccctgccaggttagtggtcagtggtgctgggacagacttcagcctcaacatccatcctgtggaggaggatgatat
 tgcaatgtatttctgcagcaaaagtaggaaggttccttgacgttcggtggagggaccaagctggaaatcaaacgggggtgg
 cggtggctcgggcgagggtgggtcgggtggcgggatctcaggtgcagctgaaggagtcaggacctggcctgggtgg
 cgccctcacagagcctgtccatcacatgcaccgttcagggttcattaaaccggctatggtgtaaactgggttcgccagcct
 ccaggaaagggtctggagtggtgggaatgatatgggtgatggaagcacagactataattcagctctcaaatccagactg
 30 agcatcaccaaggacaactccaagagccaagtttcttaaaaatgaacagctctgaaaactgatgacacagccagataactact
 gtgccagagatggttatagtaactttcattactatgttatggactactggggtaaggaacctcagtcaccgtctcctcagatct
 ggagcccaaatcttctgacaaaactcacatccccaccgtcccgacacctgaactcctgggtggatcgtcagcttctctc
 tccccccaaaacccaaggacacctcatgatctccggaccttgaggtcacatgcgtggtggtggagctgagccacga
 agacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaaagccgaggaggagca
 35 gtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtga
 aggtctccaacaaagccctcccagccccatcgagaaaacctctccaaagccaaagggcagccccgagaaccacagg
 tgtacacctgccccatcccgggatgagctgaccaagaaccagggtcagcctgacctgcctggtcaaaggcttctatccca
 gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
 acggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtga
 40 tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttaaagcggatccttcgaacccgttct
 ggtgctgctgcactcgggtgctgtccagcctgtcagcagcgagctgaccgagctcaagttcctatgcctcgggcgcgtgg
 gcaagcgcaagctggagcgcgtgcagagcggcctagacctcttctccatgctgctggagcagaacacctggagccccg
 ggcacaccgagctcctgcgcgagctgctgcctccctgcggcgccacgacctgctgcggcgctgcgacttcgagg
 cggggcgggcgccggggccgcgcctggggaagaagacctgtgtgcagcatttaacgtcatatgtgataatgtggggaa
 45 agattggagaaggctggctcgtcagctcaaagtctcagacaccaagatcgacagcatcaggacagataccccgcaac
 ctgacagagcgtgtgcgggagtcactgagaatctggaagaacacagagaaggagaacgcaacagtggcccacctggtg
 ggggctctcaggtcctgccagatgaacctggtggctgacctggtacaagaggttcagcaggcccgtagctccagaaca

ggagtggggccatgtccccgatgcatggaactcagacgcattacctccgaagcgtcctgataactcgagatcgataaca
ac

Amino acid sequence: (SEQ ID NO: 480)

5 mdfqvqifsflisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyyqqkpgppkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiamyfcqqsrkvpwtfgggkgleikrgggsgggsgggsgvqlkes
pgplvapsqslsitctvsgfsltygvnwvrpppgkglewlgmiwgdgstdynsalksrlsitkdnsksqvfikmnsi
10 qtddtaryycardgysnfhyvmdywgqtsvtvssdlepkssdkthtsppspapellggssvflfppkpkdtlmisrt
pevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpap
iektiskakgqprepvytlppsrdeltknqvsitclvkgfypsdiavewesngqpennyktpvldsdgsfflysklt
10 vdkswqqgnvfscsvmhcalhnhytqkslslspgkadsnpflvlhsvssslsssteltelclgrvgkrklervqsg
ldlfsmilleqndlepghtellrellaslrhdllrrvddfeagaaagaapgeedlcaafnvicdnvgkdwrllarqlkvsdt
kidsiedrypmltervreslriwkntekenatvahlvgalrscqmnlvadlvqevqqardlqnrgamspmswnsda
stseas

15 130. 1D8 scFv hIgG1 (SSS-P)H P238S CH2 WCH3-hCD80TMCT

Nucleotide sequence (SEQ ID NO: 108)

aagcttatggatttcaagtgcagatttccagcttctgtaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctcagtgtaagta
catgtactggtaccagcagaagtcaggcgccctcccctaaactctggattatgacacatccaagctggcttctggagtcca
20 atcgcttcagtggtcagtggtctgggacctcttattctctcgaatcaacaccatggagactgaagatgctgccacttattact
gtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacggggtggcggtggctcggg
cggtgggtgggtcgggtggcgcgatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
ccctgtccctcatgactgtctctgggtctcattaaccagcagtggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcagg
25 acacctcaagagccaagtttcttaaaaatcaacagctcgaactgatgacacagccatgtattactgtccagaatccac
tttgattactggggccaaggagtcaggtcagctcctctgatctggagcccaaatcttctgacaaaactcacatcccc
aecgtccccagcacctgaactcctggggggatcgtcagcttctctccccccaaaaccaaggacacctcatgatctcc
cggaccctgaggtcacatgcgtggtggtgacgtgagccacgaagacctgaggtcaagtcaactggtacgtggacg
gcgtggagggtgcataatgccaaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtctca
30 ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaacaaagccctccagccccatcga
gaaaaccatctccaaagccaaaggcagccccgagaaccacaggtgtacacctgccccatcccggtgagctgac
caagaaccagggtcagcctgacctgctggtcaaaaggcttctatccagcgacatcgccgtggagtgaggagcaatggg
cagccggagaacaactacaagaccacgcctcccgctggtgactccgacggctccttctctctacagcaagctcaccgtg
gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctgcacaaccactacacgcaga
35 agagcctctccctgtctccgggtaaaagcgatccttcgaactgctccatcctgggccattaccttaattctcagtaaagga
attttgtgatatgctgcctgacctactgcttggcccaagatgcagagagagaaggagggaatgagagattgagaaggga
agtgtacgccctgtataaatcgat

Amino acid sequence: (SEQ ID NO: 109)

40 mdfqvqifsflisasvimsrgvdivltqspptiaaspgekvitcrassvsymywyqqksgasplwiydtsklasgv
pnrfsgsgsgtsyslaintmetedaatyycqqsstpltfsggkgleikrgggsgggsgggsgvqlkeagpglvqp
tqtlsitctvsgfsltsdgvhwirpppgkglewmgiiydgtdynsaiksrlsirdtsksqvfikinslqtdtamyyc
arihfdywgqgmvtvssdlepkssdkthtsppspapellggssvflfppkpkdtlmisrtpevtcvvvdvshedpev
kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
tlppsrdeltknqvsitclvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdkswqqgnvfscsvm
45 healhnhytqkslslspgkadsnllpswaitlisvngifviccltycfaprerrrrerlrresvrpv

131.1D8 scFv hIgG1 (SSS-P)H WCH2 WCH3-hCD80TM/CT

Nucleotide sequence (SEQ ID NO: 481)

5 aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cacccaatctccagcttctttggctgtgtcttaggtcagagagccaccatctctgcagagccagtgaagtggtgaatatta
10 tgtcacaagttaatgcagtggtaccaacagaaaccaggacagccaccaactcctcatctctgctgcatccaacgtagaa
tctggggtcctgccaggttttagtggcagtggtctgggacagacttcagcctcaacatccatctgtggaggaggatgata
ttgcaatgtattttctgacagaaagtaagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataat
gtccagaggagtcgacattgtgtcactcagctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctg
ccgtgccagctccagtgttaagtacatgtactggtaggcagagaagtcaggcgctcccctaaactctggatttatgacacat
15 ccaagctggcttctggagtccaatcgttcagtggtcagtggtctgggacctctattctctcgcaatcaacaccatggag
actgaagatgctgccacttattactgtcagcagtggttagtactccgctcacgttcgggtctgggaccaagctggagatca
aacgggggtggcgtggctcggcggtgggtgggtcgggtggcggtgatctcaggtgcagctgaaggaggcaggacct
ggcctggtgcaaccgacacagaccctgtccctcacatgcactgtctctgggttctcattaaccagcgatggtgtacactgga
ttcgacagcctccaggaaagggtctggaatggatgggaataatatattatgatggaggcacagattataattcagcaattaa
20 tccagactgagcatcagcaggacacctccaagagccaagtttcttaaaaatcaacagctctgaaaactgatgacacagcc
atgtattactgtgccagaatccactttgattactggggccaaggagtcaggtcacagctcctctgatctggagcccaaatct
tctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagcttctcttcccccaaaaac
ccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggt
caagttaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcac
25 gtaccgtgtgtgcagcgtcctaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaaca
aagccctccagccccatcgagaaaaccatctccaagccaaagggcagccccgagaaccacaggtgtacacctgc
ccccatccgggatgagctgaccaagaaccaggtcagcctgacctgctgtaaaaggcttctatccagcgacatcgcc
gtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttct
tctctacagcaagctaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcagaggct
30 ctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaaaggatccttcgaacctgctccatcctgggcc
attacctaatactcagtaaatggaattttgtgatgtgctgacctactgctttgccccaaagatgcagagagagaaggagg
aatgagagattgagaagggaagtgtacgccctgtataaatcgat

Amino acid sequence: (SEQ ID NO: 482)

30 mdfqvqifsfllisasvimsrgvdivltqspaslavslgqratiscrasesveyyvtslmqwyqqkpgppkllisaasn
vesgvparfsgsgsgtdfslnihpveeddiamyfcqqsklmdfvqifsfllisasvimsrgvdivltqspptiaaspgek
vtitcrasssvsymwyqqksgasplkiwydtkslasgvpnrfsgsgsgtsyslaintmetedaatyycqwsstpltf
gsgtkleikrgggsgggsgggsgqvlkeagpglvqptqlslctvsgfsltsdgvhwirppgkglewmgiyy
dggtdynsaiksrlsrdtsksqvflkinslqddtamyycaihfdywqgvmvtvssdlepkssdkthtspspap
ellggpsvflfppkpkdtlmisrtpvctvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvsvltvlh
35 qdwlngkeykckvsnkalpapiektiskakqpprepqvytlppsrdeitknqvsitclvkgfypsdiavewesngqp
ennyktpvldsdgsfflyskltvdkrsrwqqgnvfscsvmhleahnhytqkslspsgkadpsnllpswaitlisvngi
fviccltycfaprerrrmerlrresvrpv

132.1D8 scFv mIgAT4-hCD80TM/CT

40 Nucleotide sequence (SEQ ID NO: 483)

45 aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctccagtgttaagta
catgtactggtaccagcagaagtcaggcgctcccctaaactctggatttatgacacatccaagctggcttctggagttcca
atcgcttcagtggtcagtggtctgggacctctattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
gtcagcagtggttagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
cggtggtgggtcgggtggcggtgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
ccctgtccctcacatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt

ctggaatggatgggaataatattatgatggaggcacagattataaattcagcaattaaatccagactgagcatcagcaggg
 acacctccaagagccaagttttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtgccagaatccac
 ttgattactggggccaaggagtcaggtcacagtctcctctgatcacatctgttctcctactactcctcctccaccttctg
 ccagcccagcctgtcactgcagcggccagctcttgaggacctgctcctgggttcagatgccagcatcacatgtactctgaat
 5 ggctgagagatcctgagggagctgtcttcacctgggagccctccactgggaaggatgcagtgcagaagaaagctgtgc
 agaattcctgcggctgctacagtgtgtccagcgtcctgacctgggtgtgtgagcgtggaacagtggcgcacattcaagt
 cacagttaccatcctgagctgtgacaccttaactggcacaattgcaaagtcacagtgaacaccttccacccccaggtccac
 ctgctaccgccgccgtcggaggagctggccctgaatgagctcgtgtccctgacatgcctgggtgcgagctttcaaccctaaa
 gaagtgtgtgcatggctgcatggaaatgaggagctgtccccagaaagctacctagtgttgagcccctaaaggagcc
 10 aggcgagggagccaccacctacctgggtgacaagcgtgttgctgtatcagctgaaatctggaaacagggtgaccagtact
 cctgcatggtggccacgaggccttggccatgaacttcaaccagaagaccatgaccgtctgtcgggtaaaccaccaat
 gtcagcgtgtctgtgatcatgtcagaggagaggatccttcgaacaacctgtcccatcctgggccattacctaattctcagt
 aatggaattttgtgatatgtgcctgacctactgctttgccccagatgcagagagagaaggaggaatgagagattgaga
 agggaaagtgtacccctgtataaatcagact

15 Amino acid sequence: (SEQ ID NO: 484)

mdfqvqifsfllisasvimsrgvdivltqspttiaaspgekvitcrasssvsymywyqqksgasplwiydtsklasgv
 pnrfsghsgsytyslaintmetedaatyycqqsstpltfsgtgleikrsgggsgggsgggsgggsgvqlkeagpglvqp
 tqtlstctvsgfsltsdgvhwirpppgkglewmgiiydgtdynsaiksrlsirdtsksqvflkinslqtdtdamyyc
 arihfdywgqgvmvtvssdhicspptppppscqpslsrlrpaedlllgsdasitctlngrdpegavftwepstgkda
 20 vqkkavqnsqgcysvssvlpgcaerwnsgasfktvthpesdltltgiakvtvntfppqvhllpppseelalnelvsltl
 vrafnpkevlvrwlhgneelspeylvfeplkepegattylvtsvlrvsaeiwkqgdqyscmvghealpmnftqkti
 drlsgkptnsvsvimsegedpsnnllpswaitlisv ngifviccltycfaprcrerrrnerlrresvrpv

133.1D8 scFv hIgE-hCD80-TM/CT

25 Nucleotide sequence (SEQ ID NO: 175)

aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
 cactcagctccaacaaccatagctgcatctccaggggagaagggtcaccatcacctgccgtgccagctccagtgttaagtta
 catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
 atcgcttcagtggcagtggtctgggacctcttattctctcgaatcaacaccatggagactgaagatgctgccacttattact
 30 gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcgggtggctcggg
 cgggtgggtgggtcgggtggcggcgatctcaggtgcagtggaaggaggcaggacctggcctgggtgcaaccgacacaga
 cctgtccctcacatgcactgtctctgggttctcattaaccagcagtggtgtacactggattcgacagcctccaggaaagggt
 ctggaatggatgggaataatattatgatggaggcacagattataaattcagcaattaaatccagactgagcatcagcaggg
 acacctccaagagccaagttttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtgccagaatccac
 35 ttgattactggggccaaggagtcaggtcacagtctcctctgatcacgtctgtcctcaggagcttccccgccaccgtga
 agatcttacagtcgtcctgcgacggcggcgggacacttcccccgacctccagctcctgtgcctcgtctctgggtacacc
 cagggactatcaacatcacctggctggaggacgggcaggtcatggacgtggactgtccaccgctctaccacgcagga
 gggtagctggcctccacacaaagcagctcacctcagccagaagcactggctgtcagaccgcacctacacctgccag
 gtcacctatcaaggtcacaccttgaggacagcaccagaagtggtgcagattccaaccgagaggggtgagcgacctacct
 40 aagccggcccagcccgttcgacctgttcacccgaagtcgcccacgatcacctgtctggtggtggacctggcaccagca
 aggggacctgaacctgacctggtcccgggacagtggaagcctgtgaaccactccaccagaaaggaggagaagcag
 cgcaatggcacgttaaccgtcacgtccacctgcccgggtgggacccgagactggatcgagggggagacctaccagtgc
 agggtagccacccccacctgcccagggccctcatgcgtccacgaccaagaccagcgcccgctgtctgccccgga
 agtctatgcgtttgcgacgccggagtggcggggagccgggacaagcgcacctcgcctgcctgatccagaacttcagt
 45 cctgaggacatctcggtgcagtggtgcacaacgaggtgcagctcccgagcggcgccagcagcagcagcagccccg
 caagaccaagggtccggcttctcgtcttcagccgctggaggtgaccagggccgaatgggagcagaaagatgagttc
 atctgccgtgcagtcctgaggcagcgagccctcacagaccgtccagcgagcgggtgtctgtaaatcccggtaaagcgg

atccttcgaagctcccatcctgggccattacctaatactcagtaaatggaattttgtgatatgctgcctgacctactgctttgcc
ccaagatgcagagagagaaggaggaatgagagattgaga- agggaaagtgtacg ccctgtataaatcgata

Amino acid sequence: (SEQ ID NO: 176)

mdfqvqifsflisavimsrgvdivltqspttiaaspgekvtitcrasssvsymywyqqksgasplwiydtsklasgv
5 pnrfsghsgsgtsyslaintmetedaatyccqqsstpltfsggtkleikrgggsgggsgggsgggsgvqlkeagpglvqp
tqtlsltctvsgfsltsdgvhwirpppgkglewmgiiyydggtdynsaiksrlsisrdsksqvflkinslqtdtdamyyc
arihfdywgqgvmvtvssdhvcsrdftpptvkilqsscdggghfpptiqlclvsgytpgtinitwledgqvmdivlsl
asttqegelastqseltsqkhwlsdrtytcqvtyqghtfedstkkcadsnprgvsaylsrpsfdlfrksptitclvdlap
skgtvnltwsrasgkpvnhstrkeekqrmgtltvtstlpvgrdwiegetyqcrvthphlpralmrsttktsgpaaapevy
10 afatpewpgsrdrklaciqnfmpeidsvqwlhnevqlpdarhsttqprktkgsgffvfrlevtraeweqkdeficra
vheaaspsqtvqravsvnpkdpkslpwaitlisvngifv iccltycfaprerrrrnerlrresvrpv

134.1D8 scFv hIgG1 (SSS-P)H WCH2 WCH3-mFADD-TM/CT

Nucleotide sequence (SEQ ID NO: 485)

15 aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaacatagctgcattccaggggagaaggtcaccatcacctgccgtgccagctccagtgtaagtta
catgtactggtagcagaagtcaggcgccctcccctaaactctggattatgacacatccaagctggcttctggagttccaa
atcgcttcagtgagcagtggtctgggacctctattctctcgaatcaacacatggagactgaagatgctgccacttattact
gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
20 cgggtgggtgggtcgggtggcgccgatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
ccctgtccctcatgactgtctctgggttctcattaaccagcagtggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagttttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtccagaatccac
tttgattactggggccaaggagtcagtggtcacagtcctctgatctggagcccaaatcttctgacaaaactcacatcccc
25 accgtccccagcacctgaactcctgggtggaccgtcagcttctctctcccccaaaacccaaggacacctcatgatctcc
cggacccctgaggtcacatgcgtggtgggtgacgtgagccacgaagacctgaggtcaagttcaactggtagctggacg
gcgtggaggtgcataatgccaaagaccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctca
ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcga
gaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatccccgggatgagctgac
30 caagaaccaggtcagcctgacctgcctgggtcaaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatggg
cagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgtg
gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctgcacaaccactacacgcaga
agagcctctccctgtctccgggtaaagcggatccttgaacatggaccattctgtgtgctgctgactcgtgtccggcag
cctgtcgggcaacgatctgatggagctcaagtcttctgccgcgagcgcgtgagcaaacgaaagctggagcgcgtgcag
35 agtggcctggacctgttcacgggtgctgctggagcagaacgacctggagcgcgggcacaccgggctgctgcgcgagttg
ctggcctcgtgcgccgacacgatctactgcagcgcctggacgacttcgaggcggggacggcgaccgctgcgcccccg
ggggaggcagatctgcaggtggcatttgacattgtgtgtgacaatgtggggagagactggaaaagactggccgcgagc
tgaaaggtgtctgaggccaagatggatgggattgaggagaagtacccccgaagtctgagtgagcgggtaaggagagagtct
gaaagtctggaagaatgctgagaagaagacgcctcgtggccggactggtcaaggcgtgcggacctgcaggctgaa
40 tctggtggctgacctggtggaagaagcccaggaatctgtgagcaagagtgagaatatgtccccagtactaagggaattcaac
tgtgtcttctcagaacaccctgactcgagatcgat

Amino acid sequence: (SEQ ID NO: 486)

mdfqvqifsflisavimsrgvdivltqspttiaaspgekvtitcrasssvsymywyqqksgasplwiydtsklasgv
45 pnrfsghsgsgtsyslaintmetedaatyccqqsstpltfsggtkleikrgggsgggsgggsgggsgvqlkeagpglvqp
tqtlsltctvsgfsltsdgvhwirpppgkglewmgiiyydggtdynsaiksrlsisrdsksqvflkinslqtdtdamyyc
arihfdywgqgvmvtvssdlepkssdkhtspspapellggpsvflfppkpkdtlmsrtpevtcvvvdvshedpev
kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeyckvsnkalpapiektiskakgqprepqv

tlppsrdeItknqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvfscsvm
healhnhytqkslsispkdpnmdpflvlhslsgslsgndlmelkflcrervskrklerqvsgldlftvlleqndlrg
htgllrellaslrhldllqrlddfagataappgeadlqvafdivcdnvgrdwkrlarelkvseakmdgieekyprrlsr
vreslkvwknaekknasvaglvkalrtcrlnlvadlveeqesvs ksenmspvldstvsstetp

5

135.1D8 scFv hIgG1 (SSS-P)H P238S CH2 WCH3-mFADD-TM/CT

Nucleotide sequence (SEQ ID NO: 487)

aagcttatggattttcaagtgcagattttcagcttctgctaatactgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctccaacaaccatagctgcattctccaggggagaaggtcaccatcacctgccgtgccagctccagtgaagtta
10 catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagtccaa
atcgcttcagtggtcagtggtctgggacctctattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
gtcagcagtggtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
cggtgggtgggtcgggtggcgggatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
ccctgtccctcacatgcactgtctctgggttctcattaaccagcgtatggtgtactggattcgacagcctccaggaaagggt
15 ctggaatggtgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagtttcttaaaaatcaacagctctgcaactgatgacacagccatgtattactgtccagaatccac
tttgattactggggccaaggagtcaggtcacagctcctctgatctggagcccaaatcttctgacaaaactcacacatcccc
accgtccccagcacctgaactcctgggtggatcgtcagcttctcttcccccaaaacccaaggacacctcatgatctcc
cggaccctgaggtcacatgcgtggtgggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacg
20 gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtcctca
ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaacaaagccctccagccccatcga
gaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccggtgatgagctgac
caagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatggg
cagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagtcaccgtg
25 gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcagggctctgcacaaccactacacgcaga
agagcctctccctgtctccgggtaaaagcggatccttgaacatggaccattctggtgctgctgcactcgtgtccggcag
cctgtcgggcaacgatctgatggagctcaagttctgtgccgcgagcgcgtgagcaaacgaaagctggagcgcgtgcag
agtggcctggacctgttcacgggtgctgctggagcagaacacctggagcgcggggcacaccgggctgctgcgcgagttg
ctggcctcgtgcgccgacacgatctactgcagcgcctggacgacttcgaggcggggacggcgaccgctgcgcccccg
30 ggggaggcagatctgcaggtggcatttgacattgtgtgtgacaatgtggggagagactggaaaagactggcccgagc
tgaaaggtgtctgaggccaagatggatgggattgaggagaagtacccccgaagtctgagtgcgggtaaggagagagtct
gaaagcttggaagaatgctgagaagaagaacgcctcgggtggcggactggtcaaggcgctgcggacctgcaggctgaa
tctggtggctgacctggtggaagaagcccaggaatctgtgagcaagagtgagaatatgtccccagtactaagggttcaac
tgtgtcttctcagaacacctgactcgagatcgat

35 Amino acid sequence: (SEQ ID NO: 488)

mdfqvqifsfllisasvmsrgvdivltqspttiaaspgekvittrasssvsymywyqqksgasplkiwydtsklasgv
pnrfsgsgsgtsyslaintmetedaatyycqqwsstpltfsgstkleikrgggsgggsgggsgqvqlkeagpglvqp
tqtlsltctvsgfsltsdgvhwirpppgklewmgiiyydgtdynsaiksrlsisrdtsksqvfkinslqtdtdamyyc
arihfdywgqgmvtvssdlepkssdkthtspspapellggssvflfppkpkdtlmisrtevtcvvdvshedpev
40 kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
tlppsrdeItknqvsltlvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvfscsvm
healhnhytqkslsispkdpnmdpflvlhslsgslsgndlmelkflcrervskrklerqvsgldlftvlleqndlrg
htgllrellaslrhldllqrlddfagataappgeadlqvafdivcdnvgrdwkrlarelkvseakmdgieekyprrlsr
vreslkvwknaekknasvaglvkalrtcrlnlvadlveeqesvs ksenmspvldstvsstetp

45

136.1D8 scFv hIgG1 (SSS-P)H WCH2 WCH3-mcasp3-TM/CT

Nucleotide sequence: (SEQ ID NO: 489)

aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
cactcagctctccaacaaccatagctgcattctccaggggagaaggtcaccatcacctgccgtgccagctccagtgttaagtta
5 catgtactgggtaccagcagaagtcaggcgccctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
atcgcttcagtggtcagtggtctgggacctcttattctctcgaatcaacacatggagactgaagatgctgccacttattact
gtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacggggtggcggtggctcggg
cggtgggtgggtcgggtggcgccgatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
ccctgtccctcacatgcactgtctctgggttctcattaaccagcagtggtgtacactggattcgacagcctccaggaaagggt
10 ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagttttcttaaaaatcaacagctctgcaaaactgatgacacagccatgtattactgtccagaatccac
tttgattactggggccaaggagtcaggtcacagctctcctctgatctggagcccaaatcttctgacaaaactcacacatcccc
accgtccccagcacctgaactcctgggtggaccgtcagcttctcttccccccaaaacccaaggacacctcatgatctcc
cggaccctgaggtcacatgcgtggtgggtgacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacg
15 gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtaacaacagcacgtaccgtgtggtcagcgtcctca
ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcga
gaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccgggatgagctgac
caagaaccaggtcagcctgacctgcctgggtcaaaagcttctatcccagcgacatcgccgtggagtgaggagcaatggg
cagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcaccgtg
20 gacaagagcaggtggcagcaggggaacgtcttctcatgctcctgatgcatgaggtctgcacaaccactacacgcaga
agagcctctccctgtctccgggtaaaagcggatccttcaaacatggagaacaacaaaacctcagtgattcaaaatccattaa
taattttgaagtaagaccatacatgggagcaagtcagtgactctgggatctatctggacagtagttacaaaatggattatcc
tgaaatgggcatatgcataataataataaagaactccataagagcactggaatgtcatctcgctctggtagcatgtgga
cgcagccaacctcagagagacattcatgggcctgaaataccaagtcaggaataaaaatgatcttactcgtgaagacatttg
25 gaattaatggatagtgtttctaaggaagatcatagcaaaaggagcagctttgtgtgtgattctaagccatggtgatgaagg
ggctatttatgggacaaatgggcctgttgaaactgaaaaagttgactagcttctcagaggcgactactgccggagctgactg
gaaagccgaaactcttcattcaggcctgccggggtacggagctggactgtggcattgagacagacagtgaggactgat
gaggagatggcttgccagaagataccggtggaggtgacttctgtatgcttactctacagcacctggttactattcctggag
aaattcaaaggacgggtcgtggttcacagtcctttgcagcatgctgaagctgtacgcgcacaagctagaatttatgcac
30 attctcactcgcgttaacaggaaggtggcaacggaattcagatccttctccctggactccactttccacgcaagaaacaga
tcccggtgattgtgtccatgctcacgaaagaactgtactttatcactagctcgagatcgatg

Amino acid sequence; (SEQ ID NO: 490)

mdfqvqifsfllisasvimsrgvdivltqspptiaaspgekvititrasssvsymwyqqksgasplkiwydtsklasgv
pnrfsgsgsgtsyslaintmetedaatyycqqsstpltfsgskleikrgggsgggsgggsgggsgvqlkeagpglvqp
35 tqtlstctvsgfsltsdgvhwirppgkglewmgiyydggtdynsaiksrlsrdtsksqvlkinslqtdtdamyyc
arihfdywggqvmvtvssdlepkssdkhtspspapellggsplfpkpkdltlmisrtevtcvvvdvshedpev
kfnwyvdgvevhnaatkpreeqynstyrsvsltlvhqdwlngkeykckvsnkalspapietiskakgqprepqvy
tlppsrdeltknqvsiltclvkgyfypsdiavewesngqpennykttppvldsdgsfflyskltvdkrsrwqqgnvfscsvm
healnhhytqkslslspgkdpnsnmennktsvdsksinnfevktihgksvdsdgiyldssykmdypemgiciniinnk
40 nfhkstgmssrsgtdvdaanlretfmglyqvrnkndltredilelmdsvskedhskrssfvcvilshgdegviygtng
pvelkkltssfrgdyrcsltgkpklfiiqacrgtelcdgietsgtdeemacqkipveadflyaystapgyyswrnskdgs
wfqslcsmklkyahklefmhiltrvrnkvatfefslsdstfhakkqipciv smltkelyfyh

137.1D8 scFv hIgG1 (SSS-P)H P238S CH2 WCH3-mcasp3-TM/CT

45 Nucleotide sequence: (SEQ ID NO: 491)

Aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgc
tcactcagctctccaacaaccatagctgcattctccaggggagaaggtcaccatcacctgccgtgccagctccagtgttaagtta

catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
 atcgcttcagtgagtggtctgggacctctattctctcgaatcaacaccatggagactgaagatgctgccacttattact
 gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
 cggtgggtgggtcgggtggcgcggtatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
 5 cctgtccctcacatgcactgtctctgggttctcattaaccagcgaatgggtgtacactggattcgacagcctccaggaaaggg
 ctggaatggatgggaataatattatgatggaggcacagattataaattcagcaattaaatccagactgagcatcagcaggg
 acacctccaagagccaagttttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtgccagaatccac
 ttgtattactggggccaaggagtcaggtcacagtctcctctga

Amino acid sequence: (SEQ ID NO: 492)

10 mdfqvqifsfllisasvimsrgvdivltqspptiaaspgekvttitrasssvsymywyqqksgasplkiwydtsklasgv
 pnrfsghsgtsyslaintmetedaatyccqwsstpltfsgskleikrgggsgggsgggsgggsgvqlkeagpglvqp
 tqtlstctvsgfsltsdgvhwirpppgkglewmgiyydggtdynsaiksrlsrdtsksqvlkinslqtdtdamyy
 arihfdywgqgvmvtvssdlepkssdkhtspspapellggssvflfpkpkdtlmistrpevtcvvdvshedpev
 kfnwyvdgvevhnaktkpreeqynstyrsvslvtlhqdwlngkeykckvsnkalpapiektiskakgpprepqvy
 15 tlppsrdeltknqvsiltclvkgyfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvfscsvm
 healnhhytqkslslspgkdpnsnmenntksvdsksinnfevktihgksvdsgiyldssykmdypemgicini
 nknfhkstgmssrsgtdvdaanlretfmglkyqvrnkndltredilemdsvskedhskrssfvcvilshgdegviygt
 ngpvelkkltssfrgdyrcsltgkpklfiiqacrgtelcgieidsgeemacqkipveadflyaystapgyyswrnskd
 gswfiqslcsmklyahklefmhiltrvrnkvatfefslldstfhakkqipcivsmllkelyfyh

20

138.1D8 scFv hIgG1 (SSS-P)H WCH2 WCH3-mcasp8-TM/CT

Nucleotide sequence (SEQ ID NO: 493)

aagcttatggattttcaagtcagatttccagcttctgctaactcagtgcttcagtcataatgtccagaggagtcgacattgtgct
 cactcagtcctcaacaaccatagctgcattctccaggggagaaggtcaccatcacctgccgtgccagctccagtgaagtta
 25 catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
 atcgcttcagtgagtggtctgggacctctattctctcgaatcaacaccatggagactgaagatgctgccacttattact
 gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
 cggtgggtgggtcgggtggcgcggtatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
 cctgtccctcacatgcactgtctctgggttctcattaaccagcgaatgggtgtacactggattcgacagcctccaggaaaggg
 30 ctggaatggatgggaataatattatgatggaggcacagattataaattcagcaattaaatccagactgagcatcagcaggg
 acacctccaagagccaagttttcttaaaaatcaacagctctgcaaactgatgacacagccatgtattactgtgccagaatccac
 ttgtattactggggccaaggagtcaggtcacagtctcctctgatctggagccaaaatcttctgacaaaactcacacatccc
 accgtccccagcacctgaactcctgggtggaccgtcagcttctcctctccccccaaaacccaaggacacctcatgatctcc
 cggacccctgaggtcacatgcgtgggtgggtgacgtgagccacgaagaccctgaggtcaagttcaactgggtacgtggacg
 35 gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtaaacagcacgtaccgtgtggtcagcgtcctca
 ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagccccatcga
 gaaaaccatctccaaagccaaaggcagccccgagaaccacaggtgtacacctgccccatcccgggatgagctgac
 caagaaccaggtcagcctgacctgctggtcaaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatggg
 cagccggagaacaactacaagaccacgctcccgtgctggactccgacggctccttctctacagcaagctcaccgtg
 40 gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcagaggtctgcacaaccactacacgcaga
 agagcctctcctgtctccgggtaaagcggatccttcgaacatggattccagagttgtcttattgctattgctgaagaactgg
 gcagtgaagacctggctgccctcaagttctgtgcttgactacatccacacaagaagcaggagaccatcgaggatgcc
 cagaagctatttctgaggctgcgggaaaagggtatgttgagggaaggcaatctgtcttctgaaagagctgctttccacat
 cagtcggtgggacctgctggtcaacttctagactgcaaccgagaggagatgggtgagagagctcggggatccagacaat
 45 gccagatttctcctacagggtcatgctcttaagctctcagaagaagtgagcgagttggaattgagatctttaaagttccttt
 gaacaatgagatccccaaatgaagctggaagatgacttgagcctgctgaaattttgtagaatggagaagaggacatg
 ctggcagaaaataactggaaaccctaaaatcaatctgtgaccaggtcaacaagagcctgctggggaagatcaggattat

gaaagatcaagcacagagagaagaatgagccttgaaggaagggaagagttgccaccttcagtttggatgagatgagcct
 caaaatggcggaactgtgtgactcgccaagagaacaagacagttagtcacggacttcagacaaagttaccaaatgaaga
 acaaacctcggggatactgtctgatcatcaacaatcatgatttcagcaaggcccggaagacataaccaactccgaaaaa
 tgaaggacagaaaaggaacagactgtgataaagggtctgagtaagacctttaaggagcttcatttgagatagtatttac
 5 gacgactgcactgcaaataaaccacgagattctagaaggctaccaaaagcgagaccacaagaacaagactgcttcat
 ctgctgtatcctatcccacgggtgacaagggtgtcgtctatggaacggatgggaaggaggcctccatctatgacctgacatct
 tacttcactgggtcaaagtgcccttcctgtctgggaacccaagatcttttcattcaggcttgccaaggaagtaactccaga
 aaggagtgcctgatgagcgaggcttcgagcaacagaaccacactttagaagtggattcatctcacaagaactatattcc
 ggatgaggcagactttctgctgggaatggctacgggtgaagaactgcgtttcctaccgagatcctgtgaatggaactggat
 10 attcagtcactttgccagagcctgagggaaagatgtcctcaaggagatgacattcttagcatcctgactggcgtgaactatga
 cgtgagcaataaagacgacaggaggaacaagggaagcagatgccacagcccaccttcacactacggaagaagctctt
 ctccctccctaatactgactcagatcgatt

Amino acid sequence (SEQ ID NO: 494)

mdfqvqifsflisasvimsrgvdivltqspttiaaspgekvttitrasssvsymywyqqksgasplwiydtsklasgv
 15 pnrfsrgsgtsyslaintmetedaatyycqqsstpltfsggtkleikrgggsgggsgggsgqvlkeagpglvqp
 tqtlstctvsgfsltsdgvhwirpppgklewmgiiydggtdynsaiksrlsisrdtsksqvlkinslqtdtamyy
 arihfdywgqgvmvtvssdlepkssdkthtspspapellggsvglfpkpkdltmisrtpevtcvvvdvshedpev
 kfnwyvdgvevhnaktkpreeqynstyrsvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
 tlppsrdeltknqvslclvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvfscsvm
 20 healnhytqkslslspgkadsnmdfqsclyaiaeelgsedlaalkflclidyiphkkqetiedaqlflrlrekmlleeg
 nlsflkellfhisrwdllvnfldcnreemvreldpnaqispyrvmlflklseevselelrsfkflnneipckckleddlsle
 ifvemekrtmlaennletksicdqvnksllgkiedyerssterrmslegreelppsvldemslkmaelcdspreqdes
 rtsdkvyqmknkprgycliinnhdfskareditqlrkmdrkgtdcdkealsktfkelhfeivsyddctaneiheilegy
 qsadhknkdcficilshgdkgvvygtdgeasiydltsyftgskcpslsgkpkiffiqacqgsnfqkgvpdeagfeqq
 25 nhtlevdssshknyipdeadflgmatvkncvsyrdpvngtwyiqslcqlsrercpqgddilsiltgvnydvsnkddrr
 nkgkqmpqptflrkklffpp

139.1D8 scFv hIgG1 (SSS-P)H P238SCH2 WCH3-mcasp8-TM/CT

Nucleotide sequence (SEQ ID NO: 495)

aagcttatggattttcaagtgcagatttcagcttctgctaatactagtgcttcagtcataatgtccagaggagtcgacattgtgct
 cactcagctccaacaaccatagctgcattccaggggagaagggtaccatcacctgccgtgccagctccagtgaagtta
 catgtactggtaccagcagaagtcaggcgctccctaaactctggatttatgacacatccaagctggctcttgaggttcaa
 atcgcttcagtggcagtggtctgggacctctattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
 gtcagcagtggtgtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcgggtggctcggg
 35 cgggtgggtgggtcgggtggcgggatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
 ccctgtccctcacatgcactgtctctgggttctcattaaccagcagtggtgtacttgattcgacagcctccaggaaagggt
 ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
 acacctccaagagccaagtttcttaaaaatcaacagctctgaaactgatgacacagccatgtattactgtccagaatccac
 ttgattactggggccaaggagtcaggtcagctcctctgatctggagcccaaatcttctgacaaaactcacacatcccc
 40 accgtccccagcacctgaactcctgggtggatcgtcagcttctcttccccccaaaacccaaggacacctcatgatctcc
 cggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactgggtacgtggacg
 gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtctca
 ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaacaaagccctccagccccatcga
 gaaaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacacctgccccatcccggtgatgagctgac
 45 caagaaccaggtcagcctgacctgcctgggtcaaaggcttctatccagcgacatcgccgtggagtgaggagcaatggg
 cagccggagaacaactacaagaccacgcctcccgctggtgactccgacggctccttctctctacagcaagctcaccgtg
 gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcaga

agagcctctccctgtctccgggtaaagcggatccttgaacatggatttccagagttgtctttatgctattgctgaagaactgg
 gcagtgaagacctggctgccctcaagttcctgtgcttggactacatccacacaagaagcaggagaccatcgaggatgcc
 cagaagctatttctgaggctgcgggaaaaggggatgttggaggaaggcaatctgtcttctgaaagagctgctttccacat
 cagtcggtgggacctgctggtcaacttcctagactgcaaccgagaggagatggtagagagctgcgggatccagacaat
 5 gcccagatttctccctacagggatgctctttaaagctctcagaagaagtgcgcgagttggaattgagatctttaaagttccttt
 gaacaatgagatcccaaatgtaagctggaagatgacttgagcctgcttgaatttttagaaatggagaagaggacatg
 ctggcagaaaataacttgaaaccctaaaatcaatctgtgaccaggtcaacaagagcctgctggggaagatcgaggattat
 gaaagatcaagcacagagagaagaatgagcctgaaggaagggaagagttgccaccttcagtttggatgagatgagcct
 caaaatggcggaactgtgtgactcgccaagagaacaagacagtgcacggacttcagacaaagttaccaaatgaaga
 10 acaaacctcggggatactgtctgatcatcaacaatcatgatttcagcaaggcccggaagacataacccaactccgaaaaa
 tgaaggacagaaaaggaacagactgtgataaagaggctctgagtaagacctttaaaggagcttcattttagatagatcttac
 gacgactgcactgcaaataaataccacgagattctagaaggctaccaaaagcgagaccacaagaacaagactgcttcat
 ctgctgtatctatcccacgggtgacaagggtgtcgtctatggaacggatgggaaggaggcctccatctatgacctgacatct
 tacttcactggttcaaagtgccttccctgtctgggaaaccaagatcttttattcaggcttgccaaggaaagtaacttcaga
 15 aaggagtgcctgatgaggcaggttcgagcaacagaaccacatttagaagtgcattcatcatctcacaagaactatattcc
 ggatgaggcagactttctgctgggaatggctacgggtgaagaactgcgttctaccgagatcctgtgaatggaacctggat
 attcagtcactttgccagagcctgagggaaagatgtcctaaggagatgacattcttagcatcctgactggcgtgaactatga
 cgtgagcaataaagacgacaggaggaacaagggaagcagatgccacagcccaccttcacactacggaagaagctctt
 ctccctccctaatgactcgatcgattc

20 Amino acid sequence (SEQ ID NO: 496)

mdfqvqifsfllisasvimsrgvdivltqspttiaaspgekvititcrasssvsymywyqqksgasplkiwydtsklasgv
 pnrfsghsgtsyslaintmetedaatyycqqsstpltfsggtkleikrgggsgggsgggsgggsgvqlkeagpglvqp
 tqtlstctvsgfsltsdgvhwirpppgkglewmgiiyydggtdynsaiksrlsirsdtksqvlkinslqtdtdamyyc
 arihfdywggqvmvtvssdlepkssdkhtspspapellggssvflfppkpkdtlmisrtevtcvvvdvshedpev
 25 kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepqvy
 tlppsrdeltknqvsiltclvkgyfypsdiavewesngqpennnyktpvldsdgsfflyskltvdkrsrwqqgnvfscsvm
 healhnhytqkslslspgkadpsnmddfqsclyaiaeelgsedlaalkflclidyiphkkqetiedaqlflrlrekmlleeg
 nlsflkellfhisrwdllvnfldcnreemvrelrdpnaqispyrvmlfklseevselelrsfkflnneipkckleddlsle
 ifvemekrtmlaennletlksicdqvnskllgkiedyer'ssterrmslegreelpsvldemslkmaelcdspreqdses
 30 rtsdkvyqmknkprgycliinnhdfskareditqlrkmkdrkgtdcdkealsktfkelhfeivsyddctaneiheilegy
 qsadhknkdcficcilshgdkgvvygtdgkeasiydltsyftgskcpslsgkpkiffiqacqgsnfqkgvpdeagfeqq
 nhtlevdssshknyipdeadflgmatvkncvsyrdpvngrtwyiqslcqlsrercpqgddilsiltgvnydvsnkddrr
 nkgkqmpqptflrklffpp

35 **140.1D8 scFv hIgG1 (SSS-P)H WCH2 WCH3-hcasp3-TM/CT**

Nucleotide sequence (SEQ ID NO: 497)

aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacattgtgct
 cactcagctccaacaaccatagctgcactcaggggagaagggtaccatcacctgccgtgccagctcagtgtaagtta
 catgtactggtaccagcagaagtcaggcgccctcccctaaactctggatttatgacacatccaagctggcttctggagttccaa
 40 atcgcttcagtggcagtggtctgggacctctattctctcgaatcaacaccatggagactgaagatgctgccacttattact
 gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
 cggtgggtgggtcgggtggcgccgatctcaggtgcagctgaaggaggcaggacctggcctgggtgcaaccgacacaga
 ccctgtccctcatatgcactgtctctgggttctcattaaccagcagtggtgtacactggattcgacagcctccaggaaagggt
 ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
 45 acacctccaagagccaagtttcttaaaaatcaacagctctgaaactgatgacacagccatgtattactgtccagaatccac
 ttgattactggggccaaggagtcaggtcacagctcctctgatctggagcccaaatcttctgacaaaactcacacatcccc
 accgtccccagcacctgaactcctgggtggaccgtcagcttctcttcccccaaaaccaaggacacctcatgatctcc

cggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacg
 gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctca
 ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaacaaagccctcccagccccatcga
 gaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgac
 5 caagaaccaggtcagcctgacctgcctggtcaaaagcttctatcccagcgacatcgccgtggagtgaggagcaatggg
 cagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagtcaccgtg
 gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcaga
 agagcctctccctgtctccgggtaagcggatccttcaacatggagaacactgaaaactcagtggttcaaaatccattaa
 aaatttgaaccaaagatcatacatggaagcgaatcaatggactctggaatatccctggacaacagttataaaatggattatc
 10 ctgagatgggtttatgtataataataataagaattttcataaaagcactggaatgacatctcggtctggtacagatgtcat
 gcagcaaaccctcagggaaacattcagaacattgaaatatgaagtcaggaataaaaatgatcttacacgtgaagaaattgtg
 aattgatgcgtgatgtttctaaagaagatcacagcaaaaggagcagttttgtttgtgcttctgagccatggtgaagaagga
 ataattttggaacaaatggacctgttgacctgaaaaaataacaaacttttcagaggggacgtttagaagctaaactgga
 aaacccaaacttttcattatcaggcctgccgtgggtacagaactggactgtggcattgagacagacagtggtgttgatgatga
 15 catggcgtgtcataaaataaccagtgaggccgacttctgtatgatactccacagcacctggtattattcttggcgaaattca
 aaggatggctcctggttcatccagtcgctttgtccatgctgaaacagtatgccgacaagcttgaatttatgcacattcttacc
 gggtaaccgaaaggtggcaacagaatttgagtcctttcctttgacgctacttttcatgcaagaaacagattccatgtattgt
 tccatgctcacaagaactctattttatcactaactcgagatcgata

Amino acid sequence: (SEQ ID NO: 498)

20 mdfqvqifsfllisasvimsrgvdivltqspttiaaspgekvititrasssvsymywyqqksgasplwiydtsklasgv
 pnrfsrgsgtsyslaintmetedaatyycqwsstpltfsgtkleikrgggsgggsgggsgggsgvqlkeagpglvqp
 tqtlstctvsgfsltsdgvhwirpppgkglewmgiyydggtdynsaiksrlsrdtsksqvfkinslqddtamyy
 arihfdywgqgvmtvssdlepkssdkthtspspapellggsplfpkpkdtlmisrteptevtcvvvdvshedpev
 kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
 25 .tlppsrdeltknqvsltlvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdksrwqqgnvfscsvm
 healnhhytqkslslspgkadsnmentensvdsksiknlepkiihgsesmdsgislndsykmdypemglciiinnk
 nfhkstmrsrgtdvdnaanlretfnlkyevrnkndltreeivelmrdrvskedhskrssfvcvllshgeegiiifngpv
 dlkkitnffrgdrsrsltgkplfiiqacrgtelcdgietsgdvdddmachkipveadflyaystapgyyswrnskdgs
 fiqslcamlkqyadklefmhiltrvnrkvatefesfsdatfhakkqipcivsmltkelyfyh

30

141. 1D8 scFv hIgG1 (SSS-P)H P238SCH2 WCH3-hcasp3-TM/CT

Nucleotide sequence: (SEQ ID NO: 499)

aagcttatggattttcaagtcagatttcagcttctgtaatacagtgcttcagtcataatgtccagaggagtcgacattgtct
 cactcagctccaacaaccatagctgcactccaggggagaaggtcacatcacctgccgtgccagctccagtgaagtta
 35 catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagtccaa
 atcgcttcagtggcagtggtctgggacctctattctctcgcaatcaacaccatggagactgaagatgctgccacttattact
 gtcagcagtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacggggtggcggtggctcggg
 cggtggtgggtcgggtggcggggatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
 ccctgtccctcacatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt
 40 ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcaggg
 acacctccaagagccaagttttctaaaaatcaacagctctgaaaactgatgacacagccatgtattactgtgccagaatccac
 ttgattactggggccaaggagtcagtggtcacagtctcctctgatctggagccaaatcttctgacaaaactcacacatccc
 accgtcccagcacctgaactcctgggtggatcgtagcttctcttccccccaaaaccaaggacacctcatgatctcc
 cggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacg
 45 gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctca
 ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaacaaagccctcccagccccatcga
 gaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgac

caagaaccaggtcagcctgacctgcctgggtcaaaaggcttctatcccagcgacatcgccgtggagtgaggagagcaatggg
cagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctacagcaagtcaccgtg
gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcaga
agagccttccctgtctccgggtaaagcggatccttcgaacatggagaacactgaaaactcagtggttcaaaatccattaa
5 aaatttggaaacaaagatcatacatggaagcgaatcaatggactctggaatatccctggacaacagtataaaatggattatc
ctgagatgggtttatgtataataataataaagaattttcataaaagcactggaatgacatctcggctctggtacagatgctgat
gcagcaaacctcagggaaacattcagaaacttgaaatatgaagtcaggataaaaatgatcttacacgtgaagaaattgtgg
aattgatgcgtgatgtttctaaagaagatcacagcaaaaggagcagttttgtgtgtcttctgagccatggtgaagaagga
ataatttttggaaacaaatggacctgttgacctgaaaaaataacaaacttttcagaggggagtcgtttagaagctaaactgga
10 aaacccaaacttttcattattcaggcctgccgtgggtacagaactggactgtggcattgagacagacagtggtgtgatgatga
catggcgtgtcataaaataaccagtggaggccgacttctgtatgcatactccacagcacctggttattattcttggcgaaattca
aaggatggctctggttcattccagtcgcttctgccatgctgaaacagtatgccgacaagctgaatttatgcacattcttacc
gggttaaccgaaaggtggcaacagaattgagtcctttccttgacgctacttttcatgcaaagaacagattccatgtattgtt
tccatgctcacaagaactctattttatcactaactcgagatcgataa

15 Amino acid sequence: (SEQ ID NO: 500)

mdfqvqifsfllisasvimsrgvdivltqspttiaaspgekvitcrasssvsymywyqqksgasplwiydtsklasgv
pnrfsgsgsgtsyslaintmetedaatyycqqwsstpltfsgskleikrgggsgggsgggsgggsgvqlkeagpglvqp
tqtlsltctvsgfsltsdgvhwirpppgkglewmgiyydgtdynsaiksrlsirdtsksqvflkinslqtdtdamyyc
arihfdywgqgvmvtvssdlepkssdkhtspspapellggssvflfppkpkdtlmisrtevtcvvdvshedpev
20 kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
tlppsrdeltknqvsltclvkgfypsdiavewesngqpennykttppvldsdgsfflyskltvdksrwqqgnvfscsvm
healnhhytqkslslspgkdpnmentensvdsksiknlepkiihgsgesmdsgisldnsykmdypemglciiinnk
nfhkstgmtsrsgtdvdaanlretfnlkyevrnkndltreeivelmrdvskedhskrssfvcvllshgeegiiifgtngpv
dlkkitnffrgdrsrstgkpklfiiqacrgteldcgietdsgvdddmachkipveadflyaystapgyyswmskdgs
25 fiqslcamlkqyadklefmhiltrvnrkvatefesfsfdatfhakkqipciivsmltkelyfyh

142.1D8 scFv hIgG1 (SSS-S)H WCH2 WCH3-hcasp8-TM/CT

Nucleotide sequence: (SEQ ID NO: 501)

aagcttatggattttcaagtgcagattttcagcttctctgtaaatcagtgcttcagtcataatgtccagaggagtcgacattgtgct
30 cactcagtcctcaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctccagtgttaagta
catgtactggtaccagcagaagtcaggcgcctcccctaaactctggatttatgacacatccaagctggcttctggagtccaa
atcgcttcagtggtcagtggtctgggacctcttattctctcgaatcaacaccatggagactgaagatgctgccacttattact
gtcagcagtggtgagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
cggtgggtgggtcgggtggcgcgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
35 cctgtccctcatatgcactgtctctgggttctcattaaccagcgatggtgtacactggattcgacagcctccaggaaagggt
ctggaatggatgggaataatattatgaggagcacagattataaattcagcaattaaatccagactgagcatcagcaggg
acacctccaagagccaagttttcttaaaaatcaacagtctgcaactgatgacacagccatgtattactgtccagaatccac
tttgattactggggccaaggagtcaggtcacagtctcctctgatctggagcccaaatcttctgacaaaactcacacatccc
accgtccccagcacctgaactcctgggtggaccgtcagcttcttctccccccaaaacccaaggacacctcatgatctcc
40 cggacccctgaggtcatatgcgtggtgggtggacgtgagccacgaagacctgaggtcaagttcaactgggtacgtggacg
gcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtagaacagcacgtaccgtgtggtcagcgtctca
ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgcaaggtctccaacaaagccctccagccccatcga
gaaaaccatctccaaagccaaaggcagccccgagaaccacaggtgtacacctgccccatcccggtgatgactgac
caagaaccaggtcagcctgacctgctggtcaaaaggcttctatcccagcgacatcgccgtggagtgaggagagcaatggg
45 cagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctacagcaagtcaccgtg
gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcaga
agagccttccctgtctccgggtaaagcggatccttcgaacatggacttcagcagaaatctttatgatattggggaacaactg

gacagtgaagatctggcctccctcaagttcctgagcctggactacattccgcaaaggaagcaagaacccatcaaggatgc
 ctgatgttattccagagactccaggaaaagagaatgttgaggaaagcaatctgccttcctgaaggagctgctctccgaa
 ttaatagactggatttgcctgattacctaactagaaaggaggagatggaaagggaacttcagacaccaggcgagg
 ctcaaatttctgcctacagggatgctctatcagatttcagaagaagtgagcagatcagaattgaggtctttaagtttctttg
 5 caagaggaaatctcaaactgcaaactggatgatgacatgaacctgctggatatttcatagagatggagaagagggtcatcc
 tgggagaaggaaagtggacatcctgaaaagagtctgtgccaaatcaacaagagcctgctgaagataatcaacgactat
 gaagaattcagcaaagagagaagcagcagcctgaagggaagtctgatgaatttcaaattggggaggagttgtgtgggt
 aatgacaatctcggactctccaagagaacaggatagtgaatcacagactttggacaaagtttaccaaatgaaaagcaaacct
 cggggatactgtctgatcatcaacaatcacaattttgcaaaagcacgggagaaagtgccaaacttcacagcattaggagac
 10 aggaatggaacacacttggatgcaggggctttgaccacgacctttgaagagcttcattttgagatcaagccccacgatgact
 gcacagtagagcaaactctatgagattttgaaaatctaccaactcatggaccacagtaacatggactgcttcatctgctgtatcc
 tctcccatggagacaagggtcatctatggcactgatggacaggaggccccatctatgagctgacatctcagttcactgg
 tttgaagtgcccttcccttgcctggaaaacccaaagtgtttttattcaggctgtcagggggataactaccagaaaggtatacct
 gttgagactgattcagaggagcaaccctatttagaaatggatttatcatcacctcaaacgagatatatcccgatgaggctga
 15 ctttctgctggggatggccactgtgaataactgttttctaccgaaacccctgcagagggaacctggtacatccagtcacttt
 gccagagcctgagagagcagatgtctcagaggcagatgatattctcacctcctgactgaagtgaactatgaagtaagcaaca
 aggatgacaagaaaaacatggggaaacagatgcctcagcctactttcacactaagaaaaaaactgtcttcccttctgattga
 gcatgcatcgata

Amino acid sequence: (SEQ ID NO: 502)

mdfqvqifsflisavimsrgvdivltqspttiaaspgekvtitcrasssvsymywyqqksgaspkliwydtsklasgv
 pnrfsghsgtsyslaintmetedaatyycqqsstpltfsgstkleikrgggsgggsgggsgggsgvqlkeagpqlvqp
 tqtlstctvsgfsltsdgvhwirppgkglewmgiyydggtdynsaiksrlsirdtsksqvfkinslqtdtdamyyc
 arihfdywgqgvmtvssdlepkssdkthtspspapellggspsvflfppkpkdtlmisrtpvctvvdvshedpev
 kfnwyvdgvevhnaktkpreeqynstyrvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
 25 tlppsrdeltknqvslclvkgfypsdiavewesngqpennyktpplvdsdgsfflyskltvdkswqqgnvfscsvm
 healhnhytqkslslspgkdpnsmdfsmlydigeqldsedlaslkflsldyipqrkqepikdalmlfqlrlekrml
 snlsflkellfrinrldlitylntrkeemerelqtpgraqisayrvmylqiseevsrsefskflqeeiskckldddmlld
 ifiemekrvilgegkldilkrvcaqinkslkiindyeeffskersslegspsdefsngeelcgvmtisdspreqdsesqtd
 kvyqmkskprgycliinnhnfakarekvpklhsirdmngthldagatlttfeelhfeikphddctveqiyeilkiyqlmd
 30 hsnmdcficilshgdkgiyygdgqeapiyeltsqftglkcpslagkpkvffiqacqgdnyqkgipvetdseeqpyle
 mdlsspqtryipdeadflgmatvnnvcvsynpaegtwyiqslcqlrercprgddiltitevnyevsnkddkknmg
 kqmpqptflrkkllvpsd

143.1D8 scFv hIgG1 (SSS-S)H P238SCH2 WCH3-hcasp8-TM/CT

35 Nucleotide sequence: (SEQ ID NO: 503)

aagccttatggattttcaagtgcagattttcagcttctgctaactcagtgcttcagtcataatgtccagaggagtcgacattgtgct
 cactcagctccaacaaccatagctgcatctccaggggagaaggtcaccatcacctgccgtgccagctccagtgaagtta
 catgtactggtagcagagaagtcaggcgcctccctaaactctggatttatgacacatccaagctggcttctggagtccaa
 atcgcttcagtggcagtggtctgggacctcttattctctcgaatcaacaccatggagactgaagatgctgccacttattact
 40 gtcagcagtgtagtagtactccgctcacgttcgggtctgggaccaagctggagatcaaacgggggtggcggtggctcggg
 cggtgggtgggtcgggtggcgcgatctcaggtgcagctgaaggaggcaggacctggcctggtgcaaccgacacaga
 cctgtccctcatgactgtctctgggttctcattaaccagcagtggtgtacactggattcgacagcctccaggaaaggggt
 ctggaatggatgggaataatattatgatggaggcacagattataattcagcaattaaatccagactgagcatcagcagg
 acacctccaagagccaagtcttcttaaaaatcaacagctctgcaaaactgatgacacagccatgtattactgtccagaatccac
 45 ttgattactggggccaaggagtcaggtcacagctcctctgatctggagccaaatcttctgacaaaactcacacatcccc
 accgtccccagcacctgaactcctgggtggatcgtcagcttctcttccccccaaaacccaaggacacctcatgatctcc
 cggacccctgaggtcacatgcgtgggtggacgtgagccacgaagacctgaggtcaagttcaactgggtacgtggacg

gcgtggaggtgcataatgccaaagacaaagccgaggagcagtagacaacagcacgtaccgtgtggtcagcgtctca
ccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaacaaagccctcccagccccatcga
gaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgac
5 caagaaccaggtcagcctgacctgcctggtaaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatggg
cagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagtcaccgtg
gacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcaga
agagcctctccctgtctccgggtaaagcggatccttcgaacatggacttcagcagaaaatctttatgatattggggaacaactg
gacagtgaagatctggcctccctcaagttcctgagcctggactacattccgcaaaggaagcaagaacccatcaaggatgc
10 ctgatgttattccagagactccaggaagagaatgttgaggaaagcaatctgtccttctgaaggagctgcttccgaa
ttaatagactggatttgctgattacctacctaactagaaaggaggagatggaaagggaacttcagacaccaggcaggg
ctcaaatttctgctacagggtcatgctctatcagatttcagaagaagtgagcagatcagaattgaggtctttaagttctttg
caagaggaaatctccaaatgcaaactggatgatgacatgaacctgctggatatttcatagagatggagaagagggtcatcc
tgggagaaggaaagttggacatcctgaaaagagtctgtgccccaaatcaacaagagcctgctgaagataatcaacgactat
gaagaattcagcaagagagaagcagcagccttgaaaggaagtctgatgaattttcaaatggggaggagttgtgtgggt
15 aatgacaatctcggactctccaagagaacaggatagtgaatcacagactttggacaaagttaccaaatgaaaagcaaacct
cggggatactgtctgatcatcaacaatcacaattttgaaaagcacgggagaaagtgccaaacttcacagcattaggagac
aggaatggaacacacttgatgcaggggctttgaccacgacctttgaagacttcattttgatcaagccccacgatgact
gcacagtagagcaaatctatgagattttgaaaatctaccaactcatggaccacagtaacatggactgcttcatctgcttatcc
tctcccatggagacaagggcacatctatggcactgatggacaggaggccccatctatgagctgacatctcagttcactgg
20 ttgaaagtgcccttcccttgctggaacccaaagtgtttttattcaggcttgctcaggggataactaccgaaaggtatacct
gttgagactgattcagaggagcaaccctatttagaaatggatttatcatcacctcaaacgagatatatccgggatgaggctga
ctttctgctggggatggccactgtgaataactgttttctaccgaaacctgcagagggaacctggtacatccagtcacttt
gccagagcctgagagagcgtatgctctcaggcgatgatatttccaccatcctgactgaagtgaactatgaagtaagcaaca
aggatgacaagaaaaacatggggaaacagatgcctcagcctactttcacactaagaaaaaaacttgcttcccttctgattga
25 gcatgcatacgtataa

Amino acid sequence: (SEQ ID NO: 504)

mdfqvqifsfllisasvimsrgvdivltqspttiaaspgekvtitcrasssvsymwyqqksgasplkiwydtsklasgv
pnrfsgsgsgtsyslaintmetedaatyycqqsstpltfsggtkleikrgggsgggsgggsgqvlkeagpglvqp
30 tqtlstlctvsgfsltsdgvhwirpppgkglewmgiiyydggtdynsaiksrlsirdtsksqvlkinslqddtamyy
arihfdywgqgvmtvssdlepkssdkhtspspapellggssvflfppkpkdtlmisrtepvctvvvdvshedpev
kfnwyvdgvevhnaktkpreeqynstyrsvsvltvlhqdwlngkeykckvsnkalpapiektiskakgqprepvy
tlppsrdeltknqvslclvkgfypsdiavewesngqpennyktpvldsdgsfflyskltvdksrwqqgnvfscsvm
healnhhytqkslslspgkdpnmdfsmlydigeqlsdslaslkflsldyipqrkqepikdalmlfqlrlekrml
35 snlsflkellfrinrldlitylnrkeemerelqtpgraqisayrvmlyqiseevsrslrsfllqeeiskckldddmnlld
ifiemekrvilgegkldilkrvcaqinkslkiindyefskersslegspldefsngeelcgvmtdsdspreqdsesqtl
kvyqmkskprgycliinnhnfakarekvpklhsirdrngthldagalttfeelhfeikphddctveqiyeilkiyqlmd
hsnmdecficilshgdkgiiygtgdqeapiyeltsqftglkcpslagkpkvffiqacqgdnyqkgipvetdseeqpyle
mdlsspqtryipdeadflgmatvnnvcsyrnpaegtwyiqslcqlsrercprgddiltilevnyevsnkddkknmg
40 kqmpqptflrklklvpsd

145. hCTLA4 IgAH IgACH2CH3

Nucleotide sequence: (SEQ ID NO: 505)

atggcttgcccttgatttcagcggcacaaggctcagctgaacctggctgccaggacctggccctgcactctcctgtttttctt
45 ctcttcatccctgtcttctgcaaagcaatgcacgtggcccagcctgctgtgtgactggccagcagccgaggtcagccagct
ttgtgtgtgagatgcacatccaggcaaaagccactgaggtccgggtgacagtgttcggcaggtgacagccaggtgactg
aagtctgtcggcaacctacatgacggggaatgagttgaccttctagatgattccatctgcacgggcacctccagtggaa

atcaagtgaacctcactatccaaggactgagggccatggacacgggactctacatctgcaaggtggagctcatgtaccac
cgccataclacctgggcataggcaacggaaccagatttatgtaattgatccagaaccgtgccagattctgatcagccagt
tcctcaactccacctacccccatccctcaactccacctacccccatccctcatgctgccacccccgactgtcactgcacc
gaccggccctcagaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcaggt
5 gtcaccttcacctggacgcccctaagtgggaagagcgctgttcaaggaccacctgacctgacctctgtggctgtacagc
gtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttgcactgctgcctacccccgagtc
aagaccccgctaaccgccaccctctcaaaatccggaacacattccggcccagggtccacctgctgccgccgccgtcgg
aggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtgctggttcgctgg
ctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccggcaggagcccagccagggcaccac
10 caccttcgctgtgaccagcatactgcgcgtggcagccgaggactggaagaagggggacaccttctcctgcatggtgggc
cacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgtcaatgtgtctgt
gtcat ggcggaggtggacggcacctgctactgataatctaga

Amino acid sequence: (SEQ ID NO: 506)

maclgfqrhkaqlnlaartwpctllffllfipvfckamhvaqpavvlassrgiasfvceyaspgkatevrvtlrqadsq
15 tevcaatymtgneltflldsictgtssgnqvnltiqglramdtglyickvelmypppyylgigngtqiyvidpepcpsd
qpvpstpptpspstpptpspschprlsihrpaledlllgseailtctltglrdasgvtftwtpssgksavqgppdrdlcgy
svssvlpgaepwnhgtftctaaypesktpltatlsksntfrpevhllpppseelalnelvtltclargfspkdvlvrwlq
gsqelprekyltwasrqepsqgtttfavtsilrvaedwkkgdtfscmvghealplftqktidrlagkpthvnnsvvma
evdgtcy

20

146. hCTLA4 IgA WH WCH2 T4CH3 (hCTLA4 IgAH IgACH2CH3)

Nucleotide sequence: (SEQ ID NO: 507)

atggcttgcccttgatttcagcggcacaaggctcagctgaacctggctgccaggacctggccctgcactctcctgtttttctt
ctcttcacccctgtcttctgcaaagcaatgcacgtggcccagcctgctgtggtactggccagcagccgaggcatgccagct
25 ttgtgtgtgagatgcatctccaggcaaaagccactgaggctccgggtgacagtgtctcggcaggctgacagccaggtgactg
aagctgtgcggcaacctacatgacggggaatgagttgaccttcctagatgattccatctgcacgggcacctccagtggaa
atcaagtgaacctcactatccaaggactgagggccatggacacgggactctacatctgcaaggtggagctcatgtaccac
cgccataclacctgggcataggcaacggaaccagatttatgtaattgatccagaaccgtgccagattctgatcagccagt
tcctcaactccacctacccccatccctcaactccacctacccccatccctcatgctgccacccccgactgtcactgcacc
30 gaccggccctcagaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcaggt
gtcaccttcacctggacgcccctaagtgggaagagcgctgttcaaggaccacctgacctgtggtgctgtacagc
gtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttcacttgcactgctgcctacccccgagtc
aagaccccgctaaccgccaccctctcaaaatccggaacacattccggcccagggtccacctgctgccgccgccgtcgg
aggagctggccctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtgctggttcgctgg
35 ctgcaggggtcacaggagctgccccgcgagaagtacctgacttgggcatcccggcaggagcccagccagggcaccac
caccttcgctgtgaccagcatactgcgcgtggcagccgaggactggaagaagggggacaccttctcctgcatggtgggc
cacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacccatgtcaatgtgtctgt
gtcat ggcggaggtggactgataatctaga

Amino acid sequence: (SEQ ID NO: 508)

Maclgfqrhkaqlnlaartwpctllffllfipvfckamhvaqpavvlassrgiasfvceyaspgkatevrvtlrqadsq
40 vtevcaatymtgneltflldsictgtssgnqvnltiqglramdtglyickvelmypppyylgigngtqiyvidpepcpsd
dqpvpstpptpspstpptpspschprlsihrpaledlllgseailtctltglrdasgvtftwtpssgksavqgppdrdlcgc
ysvssvlpgaepwnhgtftctaaypesktpltaflsksgntfrpevhllpppseelalnelvtltclargfspkdvlvrwl
qgsqelprekyltwasrqepsqgtttfavtsilrvaedwkkgdtfscmvghealplftqktidrlagkpthvnnsvv
45 maevd

147. hIGA WH WCH2 T18CH3

Nucleotide sequence: (SEQ ID NO: 509)

5 Tgatcagccagttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccga
ctgtcactgcaccgaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgag
agatgcctcaggtgtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccacctgaccgtgaccttg
10 tggctgctacagcgtgtccagtgctctgccgggctgtgccgagccatggaaccatgggaagaccttcacttgactgctgc
ctaccccgagtccaagaccccgtaaccgccacctctcaaatccggaaacacattccggcccgagggtccacctgctgc
cgccgccgtcggaggagctggccctgaacgagctgggtgacgctgacgtgcctggcacgtgggttcagcccaaggatgt
gctgggtcgtggctgcaggggtcacaggagctccccgcgagaagtacctgacttgggcacccggcaggagccag
15 ccagggcaccaccaccttcgctgtgaccagcactatgcgcgtggcagccgaggactggaagaagggggacacctctcc
tgcattggtggccacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaatgataatctag
a

Amino acid sequence: (SEQ ID NO: 510)

15 dqpvpstptpsptpsptpspschprlslhrrpaledlllgseailtctltglrdasgvftwtpssgksavqgppdrdlcgc
ysvssvlpgaepwnhgktfctaaypesktpatltsksntfrpevhllppseelalnelvtlclargfspkdvlrwl
qgsqelprekyl twasrqpsqgtttfavtsilrvaedwkkgdtfscmvghealplaftqktidrlagk

148. 2h7 scFv IgA WH WCH2 T4CH3

Nucleotide sequence: (SEQ ID NO: 70)

20 aagcttgccgccatggatttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctccagcaatcctgtctgcatctccaggggagaaggtcacatgacttcagggccagctcaagtgtaa
ttacatgcactggtaccagcagaagccaggtacctcccccacacctggatttatgccccatccaacctggcttctggagtc
cctgctcgttcagtggcagtggtctgggaccttactctctcacaatcagcagagtggaggtgaagatgctgccactt
attactgccagcagtgagtttaacccaccacgcttcggtgctgggaccaagctggagctgaaagatggcgggtggctcg
25 ggcggtggtgatctggaggaggtgggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctgggg
cctcagtgaaagatgtctgcaaggcttctggctacacattaccagttacaatatgactgggtaaagcagacacctagaca
gggcttgaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagtcaaggggcaaggccacactg
actgtagacaaaatctccagcagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
gagtgggtgactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctcttctgatcagcca
30 gtccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtcactgca
ccgaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcag
gtgtcaccttcacctggacgccctcaagtgggaagagcgctgttcaaggaccacctgaccgtgacctctgtggctgtaca
gcgtgtccagtgctctgccgggctgtgccgagccatggaaccatgggaagaccttcacttgactgctgcctaccccgagt
ccaagaccccgtaaccgccacctctcaaatccggaaacacattccggcccgagggtccacctgctgccgccgccgtc
35 ggaggagctggccctgaacgagctgggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtgctggttcgct
ggctgcaggggtcacaggagctccccgcgagaagtacctgacttgggcatcccgagggagcccagccaggggcacc
accaccttcgctgtgaccagcactatgcgcgtggcagccgaggactggaagaagggggacaccttctcctgcatggtgg
gccacgaggccctgccgctggccttcacacagaagaccatcgaccgcttggcgggtaaacccacctatgtaattgtct
gttgcattggcggaggtgactgataatctaga

40 Amino acid sequence: (SEQ ID NO: 71)

45 mdfqvqifsfllisasviiargqivlsqpailsaspgekvtmtrasssvsymhwyqqkpgsspkpwiypsnlasg
vparfsgsgsgtsysltisrveadaatyccqqswnpftfagtkleldggsgggsgggssqaylqqsgaelvr
pgasvkmsckasgytftsnnhvwkqtrpqglewigaiypngdtsynqkfkakatltvdkssstaymqlssltسد
savycarvvysnsywyfdvwgtgtvtvssdqpvpstptpsptpsptpspschprlslhrrpaledlllgseailtctlt
glrdasgvftwtpssgksavqgppdrdlcgcysvssvlpgaepwnhgktfctaaypesktpatltsksntfrpev

hlpppseelalnelvtltclargfspkdvlvrlqgsqelprekyltwasrqepsqgtttfavtsilrvaaedwkkgdtfsc
mvghéalpláftqktidrlagkpthvnsvvmaevd

149. 2H7 scFv IgA WH WCH2 T18CH3

5 Nucleotide sequence (SEQ ID NO: 75)

aagctlgccgcatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataattgccagaggacaaattg
ttctctcccagctctccagcaatcctgtctgcatctccaggggagaaggtcacaatgacttcagggccagctcaagtgtaa
ttacatgcactggtagcagaagccaggatcctcccccacccctggattatgccccatccaacctggcttctggagtc
cctgctcgttcagtgagcagtggtctgggaccttactctctcacaatcagcagagtgagggtgaagatgctgccactt
10 attactgccagcagtgagtttaacccaccacgttcggtgctgggaccaagctggagctgaaagatggcggtggtcgc
ggcgggtggtgagctggaggagtgaggagctctcaggcttatctacagcagctctggggctgagctggtgaggcctggg
cctcagtgaaagatgctcgaaggcttctggctacacattaccagttacaatatgcactgggtaaagcagacacctagaca
gggcctggaatggattggagctatttatccaggaaatggtgatacttctacaatcagaagttcaaggccaaggccacactg
actgtagacaaaatcctccagcacagcctacatgcagctcagcagcctgacatctgaagactctgcggtctatttctgtgcaa
15 gactggtgtactatagtaactcttactggtacttcgatgtctggggcacagggaccacggtcaccgtctctctgatcagcca
gttccctcaactccacctaccccatctccctcaactccacctaccccatctccctcatgctgccacccccgactgtcactgca
ccgaccggccctcgaggacctgctcttaggttcagaagcgatcctcacgtgcacactgaccggcctgagagatgcctcag
gtgtcaccttcacctggacgccctcaagtgggaagagcgctgttaaggaccacctgaccgtgacctctgtggctgctaca
gcgtgtccagtgtcctgccgggctgtgccgagccatggaacctgggaagaccttacttgcactgctgcctaccccgagt
20 ccaagacccccgctaaccgccaccctctcaaaatccggaaacacattccggcccagggtccacctgctgccgccgccgtc
ggaggagctggcctgaacgagctggtgacgtgacgtgcctggcacgtggcttcagcccaaggatgtgctggttcgct
ggctgcaggggtcacaggagctgccccgcgagaagtacctgactgggcatcccgagcagagcccagccagggcacc
accaccttcgctgtgaccagcactgctgcgtggcagccgaggactggaagaagggggacaccttctctgcatggtgg
gccacgaggccctgccgctggccttc acacagaagaccatcgaccgcttggcgggtaaatgataatctaga

25 Amino acid sequence: (SEQ ID NO: 76)

mdfqvqifsfllisasviiargqivlsqspailsaspgekvtmtcrasssvsymhwyqqkpgsspkpwiypnsnlasg
vparfsgsgsgtsysltisrveadaatyycqqswnpptfgagtklelkdggsgggsgggssqaylqqsgaelvr
pgasvkmsckasgytftsynmhvwkqtrqglewigaiypngdtsynqkfkkgkatltvdkssstaymqlssltsed
savycarvvyysnsywyfdvwgtgtvtvssdqpvpstptpspctchprlslhrpaledlllgseailtcti
30 tglrdasgvftwtpssgksavqgppdrldcgcysvssvlpgcaepwnhgktftctaaypeskpltatlsksnftfrpev
hlpppseelalnelvtltclargfspkdvlvrlqgsqelprekyltwasrqepsqgtttfavtsilrvaaedwkkgdtfsc
mvghéalpláftqktidrlagk

150. G19-4 scFv (SSS-P)WH WCH2 WCH3-hCD80TMCT

35 Nucleotide sequence: (SEQ ID NO: 511)

aagcttatggattttcaagtgcagattttcagcttctgctaatacagtgcttcagtcataatgtccagaggagtcgacatccaga
tgacacagactacatcctcctgtctgctctctgggagacagagtcaccatcagttgcagggcaagtcaggacattcga
attatttaaactggtatcagcagaaaccagatggaactgttaaactcctgatctactacacatcaagattacactcaggagtc
catcaagggtcagtgagcagtggtctggaacagattauctctcaccattgccaacctgcaaccagaagatattgccacttact
40 ttgccaacagggtaatacgttccgtggacgttcggtggaggcaccaaactgtaaccaaacgggagctcggtggcgggt
ggctcggggcgggtgggtgggtggcgcgagctatcagtgagggtccagctgcaacagctctggacctgaactggtga
agcctggagcttcaatgtcctgcaaggcctctggttactcattcactggctacatcgtgaactggctgaagcagagccatgg
aaagaaccttgagtggtgattggacttattaatccatacaaaggcttactacctacaaccagaattcaaggccaaggccacat
taactgtagacaagtatccagcacagcctacatggagctcctcagctctgacatctgaagactctgcagctctattactgtgca
45 agatctgggtactatggtgactcggactggtacttcgatgtctggggcgagggaccacggtcaccgtctcctctgatctgg
agcccaaatcttctgacaaaactcacacatccccaccgtccccagcacctgaactcctggggggaccgtcagcttctcctt

5
10

cccccaaaaccaaaggacacccctcatgatctccggacccctgaggtcacatgctgtggtggtggacgtgagccacgaa
gaccctgagggtcaagtccaactggtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcag
tacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgc
gggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggt
gtacaccctgccccatccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccca
gcgacatcgccgtggagtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccg
acgggtccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctatgctccgtga
tgcattgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaaagcggatccttcgaacctgtcc
catcctgggccattaccttaatctcagtaaatggaattttgtgatatgctgcctgacctactgctttgccccaaagatgcagaga
gagaaggaggaatgagagattgagaagggaaagt gtacgccctgtataaatcga

Amino acid sequence: (SEQ ID NO: 512)

20 **151. 2e12 scFv (CCC-P)WH WCH2 WCH3-hCD80TM/CT**

Nucleotide sequence: (SEQ ID NO: 126)

Amino acid sequence: (SEQ ID NO: 127)

tpevtcvvvdvshedpevkfnwyvdgvevhnaktkpreeqynstyrvvsvltvlhqdwlngkeykckvsnkalpa
piektiskakgqprepqvylppsrdeitknqvslclvkgfypsdiavewesngqpennykttppvldsdgsfflyskl
tvdksrwqqgnvfscsvmhcalhnhytqkslslspgkadsnllpswaitlisvngifviccltycfaprerrrnerlrr
esvrpv

5

HCD16lowFL + NL

Nucleotide sequence: (SEQ ID NO: 513)

10 aagcttgccgccatgtggcagctgctcctcccaactgctctgctacttctagtttcagctggcatgcggactgaagatctccc
aaaggctgtggtgttctggagcctcaatggtacaggggtgctcgagaaggacagtgtgactctgaagtccaggaggcct
actcccctgaggacaattccacacagtgtttcacaatgagagcctcatctcaagccaggcctcagctacttcattgacgct
gccacagtcgacgacagtggagagtacaggtgccagacaaacctctccaccctcagtgaccgggtgcagctagaagtcc
atatcggctggtgttgcctcaggccccctcgggtgggtgttcaaggaggaagaccctattcacctgaggtgtcacagctgga
agaacactgctctgcataaggtcacataattacagaatggcaaaggcaggaagtattttcatcataattctgacttctacattcc
15 aaaagccacactcaagacagcggctcctacttctgcagggggcttgttgggagtaaaaatgtgtcttcagagactgtgaa
catcaccatcactcaagggttggcagtgtaaccatctcatcttccacctgggtaccaagtctcttctgcttggtgatgg
tactccttttgcagtggacacaggactatattctctgtgaagacaacattcgaagctcaacaagagactggaaggacat
aaatttaaatggagaaaggaccctcaagacaaatgaccc

Amino acid sequence: (SEQ ID NO: 514)

20 mwqlllptallllvsagmrtdlpkavvflepqwyrvlekdsvtlkcgayspednstqwfhnslissqassyfidaat
vddsgerycqtnlstldpvqlvhighwlllqaprwvfkeedpihlrchswkntalhkvtylqngkgrkyfhnsdfyi
pkatlkdsgsyfcrglvgsknvssetvnititqglavstissffppgyqvsfclvmvllfavdtglyfsvktnirsstrdwkd
hkfkwrkdpqdk

25 From the foregoing, it will be appreciated that, although specific
embodiments of the invention have been described herein for the purpose of
illustration, various modifications may be made without deviating from the spirit and
scope of the invention. Accordingly, the present invention is not limited except as by
the appended claims.

30 All patents, patent applications, publications, scientific articles, web
sites, and other documents and materials referenced or mentioned herein are
indicative of the levels of skill of those skilled in the art to which the invention
pertains, and each such referenced document and material is hereby incorporated by
reference to the same extent as if it had been incorporated by reference in its entirety
individually or set forth herein in its entirety. Additionally, all claims in this
35 application, and all priority applications, including but not limited to original claims,
are hereby incorporated in their entirety into, and form a part of, the written description
of the invention. Applicants reserve the right to physically incorporate into this
specification any and all materials and information from any such patents,
applications, publications, scientific articles, web sites, electronically available

information, and other referenced materials or documents. Applicants reserve the right to physically incorporate into any part of this document, including any part of the written description, the claims referred to above including but not limited to any original claims.

5 The specific methods and compositions described herein are representative of preferred embodiments and are exemplary and not intended as limitations on the scope of the invention. Other objects, aspects, and embodiments will occur to those skilled in the art upon consideration of this specification, and are encompassed within the spirit of the invention as defined by the scope of the claims. It
10 will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention. The invention illustratively described herein suitably may be practiced in the absence of any element or elements, or limitation or limitations, which is not specifically disclosed herein as essential. Thus, for example,
15 in each instance herein, in embodiments or examples of the present invention, any of the terms "comprising", "consisting essentially of", and "consisting of" may be replaced with either of the other two terms in the specification. Also, the terms "comprising ", "including", containing", *etc.* are to be read expansively and without limitation. The methods and processes illustratively described herein suitably may be
20 practiced in differing orders of steps, and that they are not necessarily restricted to the orders of steps indicated herein or in the claims. It is also that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality (for example, a culture or population) of such host cells, and
25 so forth. Under no circumstances may the patent be interpreted to be limited to the specific examples or embodiments or methods specifically disclosed herein. Under no circumstances may the patent be interpreted to be limited by any statement made by any Examiner or any other official or employee of the Patent and Trademark Office unless such statement is specifically and without qualification or reservation expressly
30 adopted in a responsive writing by Applicants.

 The terms and expressions that have been employed are used as terms of description and not of limitation, and there is no intent in the use of such terms and expressions to exclude any equivalent of the features shown and described or portions

thereof, but it is recognized that various modifications are possible within the scope of the invention as claimed. Thus, it will be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

Other embodiments are within the following claims. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group.

ABSTRACT

The invention relates to novel binding domain-immunoglobulin fusion proteins that feature a binding domain for a cognate structure such as an antigen, a counterreceptor or the like, a wild-type IgG1, IGA or IgE hinge-acting region, *i.e.*,
5 IgE CH2, region polypeptide or a mutant IgG1 hinge region polypeptide having either zero, one or two cysteine residues, and immunoglobulin CH2 and CH3 domains, and that are capable of ADCC and/or CDC while occurring predominantly as polypeptides that are compromised in their ability to form disulfide-linked multimers. The fusion
10 proteins can be recombinantly produced at high expression levels. Also provided are related compositions and methods, including cell surface forms of the fusion proteins and immunotherapeutic applications of the fusion proteins and of polynucleotides encoding such fusion proteins.